



Article Analytic Automorphisms and Transitivity of Analytic Mappings

Zoriana Novosad¹ and Andriy Zagorodnyuk^{2,*}

- ¹ Department of Higher Mathematics and Quantitative Methods 10, Lviv University of Trade and Economics, Tuhan-Baranovsky Street, 79005 Lviv, Ukraine; zoryana@lute.lviv.ua
- ² Faculty of Mathematics and Computer Science, Vasyl Stefanyk Precarpathian National University, 57 Shevchenka Street, 76018 Ivano-Frankivsk, Ukraine
- * Correspondence: azagorodn@gmail.com or andriy.zagorodnyuk@pnu.edu.ua

Received: 16 November 2020; Accepted: 30 November 2020; Published: 7 December 2020



Abstract: In this paper, we investigate analytic automorphisms of complex topological vector spaces and their applications to linear and nonlinear transitive operators. We constructed some examples of polynomial automorphisms that show that a natural analogue of the Jacobian Conjecture for infinite dimensional spaces is not true. Also, we prove that any separable Fréchet space supports a transitive analytic operator that is not a polynomial. We found some connections of analytic automorphisms and algebraic bases of symmetric polynomials and applications to hypercyclicity of composition operators.

Keywords: topologically transitive operator; hypercyclic operators; function space; analytic functions on Banach spaces; symmetric polynomials on Banach spaces

MSC: 47A16; 47H60; 46E50

1. Introduction

Let *X* be a complex topological vector space. A mapping $F: X \to X$ is called an *analytic automorphism* if *F* is analytic, bijective and F^{-1} is analytic. If both *F* and F^{-1} are polynomials, then *F* is a polynomial automorphism. In this paper, we consider polynomial and analytic automorphisms of topological vector spaces and their applications to linear and nonlinear dynamics. We show that in infinite dimension spaces even a triangular polynomial map is not necessary invertible and prove that any separable Fréchet space supports a topologically transitive analytic operator that is not a polynomial. In addition, we found some interesting relations between analytic automorphisms and algebraic bases of symmetric polynomials on Banach spaces and used it for hypercyclic composition operators in spaces of analytic functions.

Let us recall some basic definitions related to analytic mappings of infinitely many variables and chaotic operators.

A function f on a complex topological vector space X with values in the complex plane \mathbb{C} is called *analytic* if it is continuous and the restriction of f to any finitely dimensional subspace is an analytic function of several variables. If Y is a locally convex topological vector space, then a mapping $F: X \to Y$ is *analytic* if $\phi \circ F$ is an analytic function for every continuous linear functional ϕ on Y. An analytic map F_n is an *n*-homogeneous polynomial if $F_n(\lambda x) = \lambda^n F_n(x)$ for all $\lambda \in \mathbb{C}$, $x \in X$. A finite sum of homogeneous polynomials $F_0 + F_1 + \cdots + F_n$, $F_n \neq 0$ is a polynomial of degree n, where F_0 is a constant in Y. All topological vector spaces considered in this paper are locally convex.

A continuous mapping $T: X \to X$ is called *hypercyclic* on a topological vector space X if there is an element $x_0 \in X$ for which the orbit under T, Orb $(T, x_0) = \{x_0, Tx_0, T^2x_0, ...\}$ is dense in X. Every such

 x_0 is called a *hypercyclic element* of *T*. A continuous mapping $T: X \to X$ is called *topologically transitive* (or just *transitive*) if for each pair *U*, *V* of non-empty open subsets of *X* there is some $n \in \mathbb{N}$ with $T^n(U) \cap V \neq \emptyset$. A transitive map is *chaotic* if it has a dense set of periodic points. If *X* is a separable Fréchet space then, according to the Birkhoff transitivity theorem [1] (p. 10), *T* is topologically transitive if and only if it is hypercyclic.

Let *T* and *S* be mapping (not necessary linear) such that $T = F \circ S \circ F^{-1}$ for some continuous bijection *F*. Then *S* is called *conjugate* to *T*. Clearly, that the relation of conjugacy is an equivalence relation and preserves hypercyclicity, transitivity and chaoticity. So, if *S* is a linear transitive (hypercyclic or chaotic) operator and *F* is an analytic automorphism, we can expect that *T* is a nonlinear analytic transitive (hypercyclic or chaotic) operator.

The question about existence of hypercyclic polynomials and analytic mappings on Banach and Fréchet spaces were considered in [2,3]. Transitive operators on topological vector spaces were studied in [4–7].

Analytic automorphisms can be applied, also, for linear dynamics. For a given analytic map $F: X \to X$ we denote by C_F the composition operator on the space H(X) of all analytic functions on the topological vector space X defined by $C_F(f) = f \circ F$. If C_{Φ} is a hypercyclic composition linear operator on H(X) and F is an analytic automorphism of X, then $C_{F \circ \Phi \circ F^{-1}}$ is a hypercyclic composition linear operator which does not commute, in general, with C_{Φ} [8]. Note that the hypercyclicity of C_{Φ} on $H(\mathbb{C})$ for $\Phi: x \mapsto x + a$ was proved by Birkhoff in [9] and generalized in [10–12], for the case $H(\mathbb{C}^n)$ and for some infinite dimensional cases in [8,13–15].

Section 1 is devoted to study basic properties of polynomial and analytic automorphisms on infinite dimensional complex spaces. We show that an injective polynomial map $P: \ell_1 \to \ell_1$ is not necessary an analytic automorphism even if its Jacobi operator $\left(\frac{\partial P_i}{\partial x_j}\right)$ is a continuous bijection. This gives us a contrast with the finite dimensional case, where the Jacobian Conjecture remains to be unsolved. In addition, we introduce some classes of analytic automorphisms on the space of all complex sequences $\mathbb{C}^{\mathbb{N}}$ and show that subspace

$$\mathcal{E} = \{(x_1,\ldots,x_n,\ldots) \in \mathbb{C}^{\mathbb{N}} \colon \sup_n |x_n|^{1/n} < \infty\}$$

is invariant with respect to this class.

In Section 2 we prove that if a Fréchet space (not necessary separable) admits a linear topologically transitive (chaotic) operator *T*, then *X* admits a topologically transitive (chaotic) polynomial map of any degree as well as topologically transitive (chaotic) analytic map that is not polynomial. Some examples are constructed.

In Section 3, using some results on symmetric polynomials on ℓ_1 , we construct some particular example of "nonlinear weighted backward shift" on $\mathbb{C}^{\mathbb{N}}$ and \mathcal{E} , which is transitive.

In Section 4 we prove some applications to linear dynamics. Using the resources in [16] results in the Fréchet algebra of all entire analytic functions $H(\mathcal{E})$ on \mathcal{E} being isometric to the algebra of bounded type symmetric analytic functions $H_{bs}(L_{\infty}[0, 1])$, we show that the composition operator $f(x) \mapsto f(x + a), f \in H(\mathcal{E})$ is hypercyclic in $H(\mathcal{E})$. Some other hypercyclic composition operators on $H(\mathcal{E})$ and $H_{bs}(L_{\infty}[0, 1])$ are constructed.

Detailed information about dynamics of linear operators is given in [1,17]. Symmetric analytic functions on ℓ_p an L_p were investigated in [16,18–22]. For details on analytic mappings on topological vector spaces we refer the reader to [23].

2. Analytic Automorphisms of Subspaces of $\mathbb{C}^{\mathbb{N}}$

A typical example of polynomial automorphism of \mathbb{C}^n to itself is a triangular polynomial map *P* defined by the following way: $P(x_1, ..., x_n) = (y_1, ..., y_n)$, where

$$y_{1} = c_{1}x_{1} + P_{0}$$

$$y_{2} = c_{2}x_{2} + P_{1}(x_{1})$$

$$y_{3} = c_{3}x_{3} + P_{2}(x_{1}, x_{2})$$

$$\dots$$

$$y_{n} = c_{n}x_{n} + P_{n-1}(x_{1}, \dots, x_{n-1}),$$

where all c_1, \ldots, c_n are nonzero constants and P_j are polynomials on \mathbb{C}^j , $j = 1, \ldots, n-1$ and $P_0 = const$. Note that for the case \mathbb{C}^2 every polynomial automorphism is a composition of triangular automorphisms, for the case \mathbb{C}^3 it is not so and for \mathbb{C}^n , n > 3 it is unknown (see [24]). Triangular polynomial automorphisms can be considered in infinite-dimensional spaces. Let *X* be a complex Banach space which can be represented as a direct topological sum of some nontrivial closed subspaces $X = X_1 \oplus \cdots \oplus X_n$. Then we define $P(x) = P(x_1, \ldots, x_n) = (y_1, \ldots, y_n)$, $x_i, y_i \in X_i$ by $y_1 = A_1(x_1)$ and $y_k = A_k(x_k) + P_{k-1}(x_1, \ldots, x_{k-1})$, $1 \le k \le n$, where A_k is a linear isomorphisms of X_k to itself and $P_{k-1}: X_1 \oplus \cdots \oplus X_{k-1}$ is a polynomial. It makes sense to consider the case of an infinity direct sum; however, in this case we have some surprising effects.

Example 1. Let $X = \ell_p, 1 \le p < \infty$ and (e_k) be the standard basis in ℓ_p , that is $e_n = (\underbrace{0, \dots, 0, 1}_k, 0, \dots)$.

Let $P: X \to X$ *be a polynomial map such that* y = P(x), $x = \sum_{k=1}^{\infty} x_k e_k$ *and*

$$y_1 = x_1$$

$$y_2 = x_2 + x_1^2$$

$$\dots$$

$$y_n = x_n + x_{n-1}^2$$

$$\dots$$

It is easy to see that P is continuous and the inverse map on the range of P can be represented by

$$\begin{aligned} x_1 &= y_1 \\ x_2 &= y_2 - y_1^2 \\ x_3 &= y_3 - (y_2 - y_1^2)^2 \\ & \cdots \\ x_n &= y_n - \left(y_{n-1} - \left(y_{n-2} - \cdots - \left(y_2 - y_1^2\right)^2 \cdots\right)^2\right)^2 \\ & \cdots \\ & \vdots \end{aligned}$$

The first observation is that P^{-1} is not a polynomial because $deg(x_n(y)) = 2^{n-1}$ for n > 1. Also, it is interesting that P is not onto. For example, there is no $x \in X$ such that $P(x) = e_1$. Indeed, $P^{-1}(2e_1) = (2, -4, 8, ..., (-1)^{n+1}2^n, ...) \notin X$. Moreover, the range of P is not dense in X. Let us suppose for simplicity that $X = \ell_1$. Then the range of P does not contain the ball of radius r < 1/4, centered at $2e_1$. Indeed, if $||y - 2e_1|| < r$, then $|y_1| > 7/4$ and $\sum_{k>1} |y_k| < 1/4$. So, for n > 1,

$$|x_n| > 49/16 - 1/4 > 2$$
,

that is, $x \notin \ell_1$. The similar argument works for the general case ℓ_p .

The famous Jacobian Conjecture asserts that a polynomial map $P: \mathbb{C}^n \to \mathbb{C}^n$, $P(x) = (P_1(x), \ldots, P_n(x))$ is a polynomial automorphism if and only if the Jacobian

$$\det\left(\frac{\partial P_i}{\partial x_j}\right) = C \neq 0,\tag{1}$$

where *C* is a constant in \mathbb{C} . It is easy to check that if *P* is a polynomial automorphism, then the Jacobian is equal to a nonzero constant. The inverse statement is an open problem for $n \ge 2$. The Jacobian Conjecture was formulated in 1939 by O. H. Keller in [25]. There is a large number of partial results on this conjecture where different approaches are used. In particular, it is known that if a polynomial map $P: \mathbb{C}^n \to \mathbb{C}^n$ is injective, then it is surjective and so is a polynomial automorphism. For more details on this open problem, we refer the reader to the survey paper [26].

For the infinite-dimensional case we can replace condition (1) by the following one: *for every fixed x the operator* $\left(\frac{\partial P_i}{\partial x_j}\right)$ *is invertible;* however, Example 1 shows that this condition does not imply that *P* is a polynomial automorphism. Indeed if *P* is as in Example 1, then

$$\left(\frac{\partial P_i}{\partial x_j}\right) = \left(\begin{array}{cccc} 1 & 0 & 0 & \cdots \\ 2x_1 & 1 & 0 & \cdots \\ 0 & 2x_2 & 1 & \cdots \\ \cdots & \cdots & \cdots & \cdots \end{array}\right)$$

is invertible but *P* is not a polynomial automorphism and it is not even an analytic automorphism.

It is interesting to ask about a "natural" domain of a triangular analytic map, that is, the map should be well defined on a topological linear space and invertible. The following proposition is evident.

Proposition 1. A triangular analytic map $F: (x_1, \ldots, x_n, \ldots) \mapsto (y_1, \ldots, y_n, \ldots)$ of the form

$$y_{1} = c_{1}x_{1} + f_{0}$$

$$y_{2} = c_{2}x_{2} + f_{1}(x_{1})$$

$$y_{3} = c_{3}x_{3} + f_{2}(x_{1}, x_{2})$$

$$\dots$$

$$y_{n} = c_{n}x_{n} + f_{n-1}(x_{1}, \dots, x_{n-1})$$

$$\dots$$

is well-defined and invertible on the space of all sequences $\mathbb{C}^{\mathbb{N}}$, where $c_n \neq 0$ and f_n are analytic functions on \mathbb{C}^n , $n \in \mathbb{N}$ and $f_0 = \text{const.}$

Let us consider a partial case of the mapping in Proposition 1, which can be given by a recurrence formula

$$y_n = a_{n1}x_n + a_{n2}x_{n-1}y_1 + \dots + a_{nk}x_{n+1-k}y_{k-1} + \dots + a_{nn}x_1y_{n-1}, \quad n \in \mathbb{N},$$
(2)

where all numbers $a_{nj} \neq 0, j \in \mathbb{N}$.

Proposition 2. There are constants $b_{i_1...i_n} \in \mathbb{C}$, $n \in \mathbb{N}$ such that the map (2) can be written by

$$y_n = \sum_{i_1 + 2i_2 + \dots + ni_n = n} b_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n}.$$
(3)

Moreover, if $|a_{ij}| \leq c$ *for some positive c and all i, j* $\in \mathbb{N}$ *, then*

$$|y_n| \leq \sum_{k=1}^n c^k \max_{i \leq n} |x_i|^k.$$

Proof. For n = 1 we have $b_1 = a_{11}$. Let us suppose that the proposition is true for every k < n. Then

$$y_n = \sum_{k=1}^n a_{nk} x_{n+1-k} \sum_{i_1+\ldots+(k-1)i_{k-1}=k-1} b_{i_1\ldots i_{k-1}} x_1^{i_1} \cdots x_{k-1}^{i_{k-1}}.$$

Since $i_1 + ... + (k - 1)i_{k-1} + n + 1 - k = k - 1 + n + (k - 1) = n$ for all k < n, the vector y_n is of the form (3). Also, from (2) we can see that

$$|y_n| \le c \max_{i \le n} |x_i| \left(\sum_{k=1}^{n-1} c^k \max_{i \le n} |x_i|^k + 1 \right) = \sum_{k=1}^n c^k \max_{i \le n} |x_i|^k.$$

Corollary 1. Let *F* be an analytic map on $\mathbb{C}^{\mathbb{N}}$ as in Proposition 2 and $|a_{ij}| \leq c$ for some positive *c* and all $i, j \in \mathbb{N}$. If there is a constant d > 0 such that

$$\sup_{n\in\mathbb{N}}|x_n|^{1/n}\leq d,$$

then there is a constant b > 0 such that

$$\sup_{n\in\mathbb{N}}|y_n|^{1/n}\leq b.$$

Proof. From Proposition 2 we have

$$\sup_{n\in\mathbb{N}}|y_n|^{1/n}\leq \sup_{n\in\mathbb{N}}\left(\sum_{k=1}^n c^k\max_{i\leq n}|x_i|^k\right)^{1/n}.$$

If cd = 1, then $\sup_{n \in \mathbb{N}} |y_n|^{1/n} \le \sup_{n \in \mathbb{N}} n^{1/n} < 2$. In the general case we can compute the sum of the geometric series

$$\sup_{n\in\mathbb{N}}|y_n|^{1/n}\leq \sup_{n\in\mathbb{N}}\left(\frac{c^nd^n-1}{cd-1}\right)^{1/n}<\infty$$

because

$$\limsup_{n\to\infty}\left(\frac{c^nd^n-1}{cd-1}\right)^{1/n}=cd.$$

Let us denote $\mathcal{E} = \{ x \in \mathbb{C}^{\mathbb{N}} \colon \sup_{n} |x_n|^{1/n} < \infty \}.$

Theorem 1. Let *F* be an analytic map on $\mathbb{C}^{\mathbb{N}}$ as in Proposition 2 and there are positive constants *c* and *c'* such that $|a_{ij}| \leq c$ and $|a_{i1}|^{-1} \leq c'$, $i, j \in \mathbb{N}$. Then \mathcal{E} is *F*-invariant in $\mathbb{C}^{\mathbb{N}}$ and the restriction F_0 of *F* onto \mathcal{E} is a bijection.

Proof. From Corollary 1, we obtain that the map $F_0: \mathcal{E} \to \mathcal{E}$ is into. The inverse map on the range of F_0 can be written by

$$x_n = -\frac{1}{a_{n1}} \big(-y_n + a_{n2} x_{n-1} y_1 + \dots + a_{nk} x_{n+1-k} y_{k-1} + \dots + a_{nn} x_1 y_{n-1} \big), \quad n \in \mathbb{N}.$$

Since $|a_{n1}|^{-1} \leq c'$, we can apply to this map Corollary 1. So, the range of F_0 is \mathcal{E} and F_0 is onto. Thus F_0 is a bijection. \Box

Note that \mathcal{E} admits function space representations. For every $x = (x_1, ..., x_n, ...)$ we can assign a complex function $f_x(t), t \in \mathbb{C}$ by

$$f_x(t) = \sum_{n=1}^{\infty} x_n t^{n-1}.$$

It is well known that the map $x \mapsto f_x$ is a linear bijection from \mathcal{E} to the space $H(0_{\mathbb{C}})$ of holomorphic germs at the origin in \mathbb{C} . Also, $H(0_{\mathbb{C}})$ is isomorphic to the space of functions of exponential type, $H_{exp}(\mathbb{C})$ via the Borrel transform

$$f_x(t)\mapsto \sum_{n=1}^{\infty}\frac{x_nt^{n-1}}{(n-1)!}.$$

Example 2. Let us consider a special case of mapping (2) on $\mathbb{C}^{\mathbb{N}}$. We denote by $\Theta \colon \mathbb{C}^{\mathbb{N}} \to \mathbb{C}^{\mathbb{N}}$, $\Theta(x) = y$ such that

$$y_n = x_1 y_{n-1} - x_2 y_{n-2} + \dots + (-1)^{n-1} x_n, \quad n \in \mathbb{N}, \ x_0 = y_0 = 0$$

Then $\Theta^{-1} = \Theta$ *, that is,* $\Theta^2(x) = x$ *,* $x \in \mathbb{C}^{\mathbb{N}}$ *and* Θ *satisfies conditions of Theorem* 1*.*

3. The Existence of Hypercyclic Analytic Mappings

Theorem 2. Let X be an infinite dimension Fréchet space (not necessary separable). If X admits a linear topologically transitive (chaotic) operator T, then X admits a topologically transitive (chaotic) polynomial map of any degree as well as topologically transitive (chaotic) analytic map that is not polynomial and is conjugate to T.

Proof. It is enough to construct an analytic automorphism $\Phi: X \to X$ such that $\Phi \circ T \circ \Phi^{-1}$ is nonlinear. We know that there are a lot of ways to define an analytic automorphism. Let X_1 be a 1-dimensional subspace in X and X_2 be a complemented subspace, that is, $X = X_1 \oplus X_2$. Let F be an analytic map on X such that $F(x_1 + x_2) = F(x_1) \in X_2$ for all $x_1 \in X_1$ and $x_2 \in X_2$. We set

 $\Phi = I + F$, where *I* is the identity operator.

Then $\Phi^{-1} = I - F$ and so Φ is an analytic automorphism. Let us write $T(x) = (T(x))_1 + (T(x))_2$, where $(T(x))_1 \in X_1$ and $(T(x))_2 \in X_2$. Then, for every $x = (x_1, x_2) = x_1 + x_2$, $x_i \in X_i$

$$\Phi \circ T \circ \Phi^{-1}(x_1, x_2) = \Phi \circ T(x_1, x_2 - F(x_1))$$
$$= \Phi\Big(\Big(T(x_1, x_2 - F(x_1))\Big)_1, \Big(T(x_1, x_2 - F(x_1))\Big)_2\Big)$$
$$= \Big(T(x_1, x_2 - F(x_1))\Big)_1 + \Big(T(x_1, x_2 - F(x_1))\Big)_2 + F\Big(\Big(T(x_1, x_2 - F(x_1))\Big)_1\Big)$$
$$= T(x - F(x)) + F(T(x) - F(x)) = T(x) + T \circ F(x) - F \circ T(x)$$

since $F(x) \in X_2$ and by the definition of F, F(T(x) - F(x)) = F(T(x)). Thus, to make sure that $\Phi \circ T \circ \Phi^{-1}$ ia an *m*-degree polynomial we have to guaranty that $T \circ F - F \circ T$ is so. Let us suppose that $T(X_1) \subset X_2$ and $F(tz_1) = f(t)z_2$ for some $z_1 \in X_1$, $z_1 \neq 0$ and $z_2 \in X$ such that $T(z_2) \neq 0$, where f is a function of $t \in \mathbb{C}$. Then

$$\Phi \circ T \circ \Phi^{-1}(tz_1) = T(tz_1) + T \circ F(tz_1) - F \circ T(tz_1) = tT(z_1) + f(t)T(z_2)$$

Therefore, if *f* is an *m*-degree polynomial or an analytic function which is not a polynomial, then $\Phi \circ T \circ \Phi^{-1}$ is so. \Box

Note first that if we put instead of f a continuous function on \mathbb{C} which is not analytic, we can get a continuous nonanalytic topologically transitive (or chaotic) map. Also, $(\Phi \circ T \circ \Phi^{-1})^n = \Phi \circ T^n \circ \Phi^{-1}$, and for the map in Theorem 2,

$$\left(\Phi \circ T \circ \Phi^{-1}\right)^n(x) = T^n(x) + T^n \circ F(x) - F \circ T^n(x).$$

It is known (see e.g., [6]) that the left backward shift $T: (x_1, \ldots, x_n, \ldots) = (x_2, \ldots, x_{n+1}, \ldots)$ on $\mathbb{C}^{\mathbb{N}}$ is hypercyclic.

Corollary 2. Let $F: \mathbb{C}^{\mathbb{N}} \to \mathbb{C}^{\mathbb{N}}$ be as in Proposition 1. Then $F \circ T \circ F^{-1}$ is hypercyclic. If the function f_1 is nonlinear, then $F \circ T \circ F^{-1}$ is nonlinear and if f_1 is an analytic function which is not a polynomial, then $F \circ T \circ F^{-1}$ is so.

Proof. The hypercyclisity of $F \circ T \circ F^{-1}$ it follows from the hypercyclisity of *T* and Proposition 1. Let

$$(z_1,\ldots,z_n,\ldots)=F\circ T\circ F^{-1}(x_1,\ldots,x_n,\ldots).$$

Then from the direct calculations we can see that

$$z = \frac{c_1}{c_2} \left(x_2 - f_1 \left(\frac{x_1}{c_1} - f_0 \right) \right) - f_0$$

that completes the proof. \Box

4. Analytic Automorphisms and Symmetric Polynomials

In this section we apply some results on symmetric polynomials of infinite many variables to analytic automorphism. For us it is convenient consider polynomials defined on the complex Banach space ℓ_1 . Let us recall some definitions.

A polynomial $P: \ell_1 \to \mathbb{C}$ is said to be *symmetric* if it is invariant with respect to permutations of basis vectors $e_n, n \in \mathbb{N}$. Let us denote by $\mathcal{P}_s(\ell_1)$ the space of all symmetric polynomials on ℓ_1 . It is well known [20] that polynomials

$$F_k(z) = \sum_{n=1}^{\infty} z_n^k, \quad z = \sum_{n=1}^{\infty} z_n e_n \in \ell_1$$

form an algebraic basis in $\mathcal{P}_s(\ell_1)$, that is, every polynomial in $\mathcal{P}_s(\ell_1)$ can be uniquely represented as a finite algebraic combination of polynomials $\{F_k\}_{k=1}^{\infty}$.

If $\{Q_k\}_{k=1}^{\infty}$ is another algebraic basis of homogeneous polynomials in $\mathcal{P}_s(\ell_1)$, then

$$Q_n = a_{n1}F_n + a_{n2}F_{n-1}Q_1 + \dots + a_{nk}F_{n+1-k}Q_{k-1} + \dots + a_{nn}F_1Q_{n-1}, \quad n \in \mathbb{N},$$
(4)

where all numbers $a_{nj} \neq 0, j \in \mathbb{N}$. Comparing this equation with (2), we can see that there is a bijection between algebraic bases of $\mathcal{P}_s(\ell_1)$ and analytic automorphisms (2) on $\mathbb{C}^{\mathbb{N}}$. We denote by F_Q the map (2) associated with (4).

The basis $\{F_k\}$ is called *the basis of power polynomials*. There some other classical algebraic basis in $\mathcal{P}_s(\ell_1)$, so-called *the basis of elementary symmetric polynomials*

$$G_k(z) = \sum_{n_1 < n_2 < \dots < n_k} z_{n_1} \dots z_{n_k}.$$

By the Newton formulas, the corresponding analytic automorphism F_G can be written as

$$ny_n = \sum_{j=1}^n (-1)^{j+1} x_j y^{n-j}, \quad x, y \in \mathbb{C}^{\mathbb{N}}.$$

In [8] it is proved that if τ_a is a homomorphism of $\mathcal{P}_s(\ell_1)$, defined on the algebraic basis $\{F_k\}$ by

$$F_k(z) \mapsto F_k(z) + F_k(a)$$

for some $a \in \ell_1$, $a \neq 0$, then

$$\tau_a(G_n)(z) = \sum_{j=0}^n G_j(z)G_{n-j}(a), \quad z \in \ell_1, \, n \in \mathbb{N}, \, G_0 = 1.$$
(5)

Proposition 3. Let $y = F_G(x)$ and $y' = F_G(x')$ for some $x, x' \in \mathbb{C}^{\mathbb{N}}$. Then

$$u = F_G(x + x'),$$
 where $u_n = \sum_{j=0}^n y_j y'_{n-j},$ $y_0 = y'_0 = 1.$

Proof. If there are $z, a \in \ell_1$ such that $F_n(z) = x_n$ and $F_n(a) = x'_n$ for all natural n, then $G_n(z) = y_n$ and $G_n(a) = y'_n$, and the statement is true by Formula (5). Let us show that it is true for all $x, x' \in \mathbb{C}^{\mathbb{N}}$.

Let $u = F_G(x + x')$ and we suppose for contrary that $u \neq v$, where

$$v_n = \sum_{j=0}^n y_j y'_{n-j}.$$

Let *m* be the smallest natural number such that $u_m \neq v_m$. We denote by *w* and *b* vectors in ℓ_1 such that $F_n(w) = x_n$ and $F_n(b) = x'_n$ for $1 \leq n \leq m$. Such vectors always exist (see e.g., [18]). Then $G_n(w) = y_n$ for $1 \leq n \leq m$ because y_n depends only on x_1, \ldots, x_n . By the same reason, $G_n(a) = y'_n$ for $1 \leq n \leq m$. So, according to (5), $u_m = v_m$. A contradiction. \Box

In [5], the following result was proved.

Theorem 3. Let X be a locally convex sequence space such that $c_{00} \subset X \subset \mathbb{C}^{\mathbb{N}}$ with continuous inclusions, and such that c_{00} is dense in X. If ω is a strictly positive weight such that the weighted backward shift T_{ω} is continuous on X, then $I + T_{\omega}$ is a transitive operator on X.

Corollary 3. Let ω be a strictly positive weight. The mapping $\Phi \colon \mathbb{C}^{\mathbb{N}} \to \mathbb{C}^{\mathbb{N}}$ defined by

$$\Phi(y) = u$$
, where $u_n = \sum_{j=0}^n \omega_{n-j} y_j y_{n+1-j}$, $n \in \mathbb{N}$

is transitive and so hypercyclic.

Proof. Note that if $x = F_G^{-1}(y)$, then

$$(I+T_{\omega})\circ F_G^{-1}(y)=x+T_{\omega}(x)=(x_1+\omega_1x_2,\ldots,x_n+\omega_nx_{n+1},\ldots).$$

Thus, according to Proposition 3,

$$\Phi(y) = F_G \circ (I + T_\omega) \circ F_G^{-1}(y).$$

By Theorem 3, $I + T_{\omega}$ is transitive. Hence Φ is transitive and so hypercyclic, since $\mathbb{C}^{\mathbb{N}}$ is a Fréchet space. \Box

Let us consider a modified version of F_G , defined by

$$v = F(x), \quad v_n = \sum_{j=1}^{n-1} (-1)^{n+1} \frac{x_j v_{n-j}}{n-j} + (-1)^{n+1} x_n, \quad x \in \mathbb{C}^{\mathbb{N}}, \, n \in \mathbb{N}.$$
(6)

It is easy to see that if we put $y_n = v_n/n$, then $y = F_G(x)$. Using similar arguments as in Proposition 3 we have the following result.

Proposition 4. Let v = F(x) and v' = F(x') for some $x, x' \in \mathbb{C}^{\mathbb{N}}$. Then

$$u = F(x + x'), \quad where \quad u_n = \sum_{j=0}^n \frac{n}{j(n-j)} v_j v'_{n-j}, \quad y_0 = y'_0 = 1$$

The mapping *F* is interesting for us because the subspace $\mathcal{E} \subset \mathbb{C}^{\mathbb{N}}$ is invariant of *F* by Theorem 1. So, we have the following theorem.

Theorem 4. Let ω be a strictly positive weight. The mapping $\Psi \colon \mathbb{C}^{\mathbb{N}} \to \mathbb{C}^{\mathbb{N}}$ defined by

$$\Psi(y) = u$$
, where $u_n = \sum_{j=0}^n \frac{n}{j(n-j+1)} \omega_{n-j} y_j y_{n+1-j}$, $n \in \mathbb{N}$

is hypercyclic. If ω is such that the weighted backward shift T_{ω} is continuous on \mathcal{E} , then the restriction Ψ_0 of Ψ to \mathcal{E} is transitive.

Proof. As in Corollary 3, we can see that $\Psi = F \circ (I + T_{\omega}) \circ F^{-1}$ and if F_0 is the restriction of F to \mathcal{E} , then $\Psi = F_0 \circ (I + T_{\omega}) \circ F_0^{-1}$. \Box

5. Applications to Linear Dynamics

If we have an analytic automorphism F on a topological vector space X, then the composition operator C_F is a linear isomorphism of the space H(X) of analytic functions on X, defined by $C_F(f) = f \circ F$, $f \in H(X)$. It is well known that the translation operator $f(x) \mapsto f(x + a)$ defined on the space of entire functions $H(\mathbb{C}^n)$ is hypercyclic if $a \neq 0$. Infinite-dimensional generalizations of this results can be found in [8,13,15].

Let us denote by $H(\mathcal{E})$ the space of all analytic functions on \mathcal{E} . In [16], it is proved that $H(\mathcal{E})$ is a Fréchet algebra and isomorphic to algebra $H_{bs}(L_{\infty}[0;1])$ of all symmetric analytic functions of bounded type on $L_{\infty}[0;1]$. Let us recall that algebra $H_{bs}(L_{\infty}[0;1])$ consists of all analytic functions on $L_{\infty}[0;1]$ which are bounded on bounded subsets and invariant with respect measure-preserving measurable automorphisms of [0;1]. This algebra is generated by polynomials

$$R_n(z) = \int_{[0;1]} (z(t))^n \, dt, \quad z(t) \in L_\infty[0;1],$$

that is, every polynomial in $H_{bs}(L_{\infty}[0;1])$ is an algebraic combination of polynomials $\{R_n\}$ and the set of polynomials is dense in $H_{bs}(L_{\infty}[0;1])$. The topology of $H_{bs}(L_{\infty}[0;1])$ is the projective topology generated by norms

$$||f||_r = \sup_{||z|| \le r} |f(z)|, \quad f \in H_{bs}(L_{\infty}[0;1]), \quad r \in \mathbb{Q}_+.$$

According to [16], the isomorphism between $H_{bs}(L_{\infty}[0;1])$ and $H(\mathcal{E})$ is given by

$$R_n \mapsto \phi_n, \quad n \in \mathbb{N}$$

where ϕ_n is the coordinate functional, $\phi_n(x) = x_n, x = (x_1, \dots, x_n, \dots) \in \mathcal{E}$.

In [21], it is proved that $H_{bs}(L_{\infty}[0;1])$ is isomorphic to the algebra of symmetric analytic functions of bounded type on $L_{\infty}[0;\infty) \cap L_1[0;\infty)$, which is denoted by $H_{bs}(L_{\infty}[0;\infty) \cap L_1[0;\infty))$. Algebra $H_{bs}(L_{\infty}[0;\infty) \cap L_1[0;\infty))$ is generated by polynomials

$$\mathcal{R}_n(z) = \int_{[0;\infty)} (z(t))^n dt, \quad z(t) \in L_\infty[0;\infty) \cap L_1[0;\infty)$$

and the isomorphism is given by

$$\mathcal{R}_n \mapsto R_n, \quad n \in \mathbb{N}.$$

Let x = x(t) and y = y(t) belong to $L_{\infty}[0, \infty) \cap L_1[0, \infty)$. Let us define $z = x \bullet y$ by

$$z(t) = \begin{cases} x(t-n), & \text{if } 2n \le t < 2n+1, \\ y(t-n-1), & \text{if } 2n+1 \le t < 2n+2, \end{cases}$$

n = 0, 1, 2, ... Clearly that

$$\mathcal{R}_n(x \bullet y) = \mathcal{R}_n(x) + \mathcal{R}_n(y), \quad n \in \mathbb{N}.$$

In [14], it is proved that the operator $f(x) \mapsto f(x \bullet y)$ is hypercyclic in $H_{bs}(L_{\infty}[0;\infty) \cap L_1[0;\infty))$ for any fixed $y \in L_{\infty}[0;\infty) \cap L_1[0;\infty)$, $y \neq 0$. In [16], it is shown that for every $c = (c_1, \ldots, c_n, \ldots) \in \mathcal{E}$ there is a function $y(t) \in L_{\infty}[0;\infty) \cap L_1[0;\infty)$ such that $\mathcal{R}_n(y) = c_n$, $n \in \mathbb{N}$. Combining mentioned results in [14,16,21] we have the following theorem.

Theorem 5. Let $c = (c_1, \ldots, c_n, \ldots) \in \mathcal{E}$, $c \neq 0$. Then operator \mathcal{T}_c on $H(\mathcal{E})$

$$\mathcal{T}_c(f(x)) = f(x+c)$$

is hypercyclic.

Let $\{Q_n\}$ be another algebraic basis of homogeneous polynomials in $L_{\infty}[0;\infty) \cap L_1[0;\infty)$, deg $Q_n = n$. Then

$$\mathcal{Q}_n = \sum_{k=1}^n a_{nk} \mathcal{R}_{n+1-k} \mathcal{Q}_{k-1}.$$
(7)

If the mapping y = F(x) defined by

$$y_n = \sum_{k=1}^n a_{nk} x_{n+1-k} y_{k-1} \tag{8}$$

is an analytic automorphism of \mathcal{E} , then for every $y = (y_1, \ldots, y_n, \ldots)$ there exists $z(t) \in L_{\infty}[0; \infty) \cap L_1[0; \infty)$ such that $y_n = \mathcal{Q}_n(z), n \in \mathbb{N}$. We denote by C_F the composition operator $C_F(f) = f \circ F$, $f \in H(\mathcal{E})$. Since F is an analytic automorphism of \mathcal{E} , then C_F is a continuous isomorphism of $H(\mathcal{E})$.

Corollary 4. Let $\{Q_n\}$ be an algebraic basis of polynomials in $L_{\infty}[0; \infty) \cap L_1[0; \infty)$, defined by (7) such that the mapping (8) is an analytic automorphism of \mathcal{E} . Let u(t) be a function in $L_{\infty}[0; \infty) \cap L_1[0; \infty)$ such that $Q_n(u) \neq 0$ for some n. Then the homomorphism of $H_{bs}(L_{\infty}[0; \infty) \cap L_1[0; \infty))$ defined on $\{Q_n\}$ by

$$\mathcal{Q}_n \mapsto \mathcal{Q}_n + \mathcal{Q}_n(u)$$

is hypercyclic and invertible.

Proof. For every $f(y) \in H(\mathcal{E})$, $y = (y_1, \ldots, y_n, \ldots)$ we claim that $f(\mathcal{Q}_1(z), \ldots, \mathcal{Q}_n(z), \ldots)$ belongs to $H_{bs}(L_{\infty}[0; \infty) \cap L_1[0; \infty))$. Indeed,

$$(\mathcal{Q}_1(z),\ldots,\mathcal{Q}_n(z),\ldots)=F(\mathcal{R}_1(z),\ldots,\mathcal{R}_n(z),\ldots)$$

and so

$$f(\mathcal{Q}_1(z),\ldots,\mathcal{Q}_n(z),\ldots)=C_F(f)(\mathcal{R}_1(z),\ldots,\mathcal{R}_n(z),\ldots)\in H(\mathcal{E}).$$

Since \mathcal{T}_c is hypercyclic for any $c \neq 0$, $C_F^{-1}\mathcal{T}_c C_F$ is hypercyclic. Set $c_n = \mathcal{Q}_n(u)$ for a function $u(t) \in L_{\infty}[0;\infty) \cap L_1[0;\infty)$ such that $\mathcal{Q}_n(u) \neq 0$ for some *n*. Then

$$f(\mathcal{Q}_1(z) + \mathcal{Q}_1(u), \dots, \mathcal{Q}_n(z) + \mathcal{Q}_n(u), \dots) = C_F^{-1} \mathcal{T}_c C_F(f)(\mathcal{R}_1(z), \dots, \mathcal{R}_n(z), \dots).$$

Example 3. Let F_0 be the restriction of the analytic automorphism (6) to \mathcal{E} . That is,

$$y = F(x)$$
, where $y_n = \sum_{j=1}^{n-1} (-1)^{n+1} \frac{x_j y_{n-j}}{n-j} + (-1)^{n+1} x_n$, $x \in \mathbb{C}^{\mathbb{N}}$, $n \in \mathbb{N}$

and $T_{x'}(x) = x + x'$. Then, by Proposition 4,

$$\left(F_0 \circ T_{x'} \circ F_0^{-1}(y)\right)_n = \sum_{i+j=n, i \neq 0} \frac{i+j}{ij} y_j y'_j + y_n + y'_n$$

Thus $C_{F_0 \circ T_{x'} \circ F_0^{-1}(y)} = C_{F_0}^{-1} \mathcal{T}_{x'} C_{F_0}$ is hypercyclic on the Fréchet algebra $H(\mathcal{E})$.

6. Discussion and Conclusions

The paper is an invitation to study analytic automorphisms of infinite dimensional topological vector spaces and possible applications in functional analysis. We show that, by using analytic and polynomial automorphisms, it is possible to construct various examples of nonlinear topological transitive mappings. On the other hand, if *F* is an analytic automorphisms of a topological vector space *X*, then the composition operator C_F is an isomorphism of the algebra of analytic functions H(X). Thus, it is possible to construct new hypercyclic composition operators on spaces of analytic functions. Some applications for linear dynamics on spaces of symmetric analytic functions are proposed.

Note that if we use injective analytic maps with dense ranges $F: Y \to X$, instead of analytic automorphisms, then we can get *quasiconjugate* mappings *S* and *T*, that is, $T \circ F = F \circ S$. The quasiconjugacy is more general than conjugacy and also preserves topological transitivity and chaoticity [1] (pp. 11–12). However, there are some technical difficulties in constructing natural examples of transitive analytic mappings by using the quasiconjugacy, because the closure of the range of an analytic mapping is not necessary a linear subspace in *X*. Our future investigations will study this further.

Author Contributions: Z.N. and A.Z. contributed equally to this work. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the National Research Foundation of Ukraine, 2020.02/0025, 0120U103996.

Acknowledgments: The authors are grateful to anonymous referees for important suggestions and guidelines in order to make this paper more coherent.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Grosse-Erdmann K.G.; Peris Manguillot, A. Linear Chaos; Springer: London, UK, 2011. [CrossRef]
- 2. Bernardes, N.C.; Peris, A. On the Existence of Polynomials with Chaotic Behaviour. *J. Funct. Spaces* 2013, 2013, 320961. [CrossRef]
- 3. Peris, A. Chaotic polynomials on Banach spaces. J. Math. Anal. Appl. 2003, 287, 487–493. [CrossRef]
- 4. Ansari, S. I. Existence of hypercyclic operators on topological vector spaces. *J. Funct. Anal.* **1997**, *148*, 384–390. [CrossRef]
- 5. Bonet, J.; Frerick, L.; Peris, A.; Wengenroth, J. Transitive and hypercyclic operators on locally convex spaces. *Bull. Lond. Math. Soc.* **2005**, *37*, 254–264. [CrossRef]
- 6. Bonet, J.; Peris, A. Hypercyclic Operators on Non-normable Fréchet Spaces. J. Funct. Anal. **1998**, 159, 587–595. [CrossRef]
- 7. Manoussos, A. A Birkhoff type transitivity theorem for non-separable completely metrizable spaces with applications to linear dynamics. *J. Oper. Theory* **2013**, *70*, 165–174. [CrossRef]
- 8. Novosad, Z.; Zagorodnyuk A. Polynomial automorphisms and hypercyclic operators on spaces of analytic functions. *Arch. Math.* **2007**, *89*, 157–166. [CrossRef]
- 9. Birkhoff, G.D. Démonstration d'un théorème élémentaire sur les fonctions entières. *CR Acad. Sci. Paris* **1929**, 189, 473–475.
- 10. Bourdon, P.S.; Shapiro, J.H. Cyclic phenomena for composition operators. *Mem. Amer. Math. Soc.* **1997**, 125, 1–46. [CrossRef]
- 11. Chan, K. C.; Shapiro, J. H. The cyclic behavior of translation operators on Hilbert spaces of entire functions. *Indiana Univ. Math. J.* **1991**, *40*, 1421–1449. [CrossRef]
- 12. Godefroy, G.; Shapiro, J.H. Operators with dense, invariant, cyclic vector manifolds. *J. Funct. Anal.* **1991**, *98*, 229–269. [CrossRef]
- 13. Aron R.; Bès, J. Hypercyclic differentiation operators. Contemp. Math. 1999, 232, 39-46.
- Chernega, I.; Holubchak, O.; Novosad, Z.; Zagorodnyuk, A. Continuity and hypercyclicity of composition operators on algebras of symmetric analytic functions on Banach spaces. *Eur. J. Math.* 2020, *6*, 153–163. [CrossRef]
- 15. Mozhyrovska, Z. H.; Zagorodnyuk, A. V. Hypercyclic behavior of translation operators on spaces of analytic functions on Hilbert spaces. *J. Funct. Spaces* **2015**, 2015, 139289. [CrossRef]
- 16. Galindo, P.; Vasylyshyn, T.; Zagorodnyuk, A. Analytic structure on the spectrum of the algebra of symmetric analytic functions on *L*_∞. *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A Mat.* **2020**, *114*, 56. [CrossRef]
- 17. Bayart, F.; Matheron, E. Dynamics of Linear Operators; Cambridge University Press: New York, NY, USA, 2009.
- Alencar, R.; Aron, R.; Galindo, P.; Zagorodnyuk, A. Algebras of symmetric holomorphic functions on *l_p*. *Bull. Lond. Math. Soc.* 2003, *35*, 55–64. [CrossRef]
- 19. Chernega, I.; Galindo, P.; Zagorodnyuk, A. Some algebras of symmetric analytic functions and their spectra. *Proc. Edinb. Math. Soc.* **2012**, *55*, 125–142. [CrossRef]
- 20. González, M.; Gonzalo, R.; Jaramillo, J.A. Symmetric polynomials on rearrangement-invariant function spaces. *J. Lond. Math. Soc.* **1999**, *59*, 681–697. [CrossRef]
- 21. Vasylyshyn, T. Algebras of entire symmetric functions on spaces of Lebesgue-measurable essentially bounded functions. *J. Math. Sci.* 2020, 246, 264–276. [CrossRef]
- 22. Chernega, I.; Zagorodnyuk, A. Note on Bases in Algebras of Analytic Functions on Banach Spaces. *Carpathian Math. Publ.* **2019**, *11*, 42–47. [CrossRef]
- 23. Dineen, S. *Complex Analysis on Infinite Dimensional Spaces;* Springer: Berlin/Heidelberg, Germany, 1999; p. 543. [CrossRef]
- 24. Shestakov, I.; Umirbaev, U. The tame and the wild automorphisms of polynomial rings in three variables. *J. Am. Math. Soc.* **2004**, *17*, 197–227. [CrossRef]
- 25. Keller, O.H. Ganze Cremonatransformationen Monatschr. Math. Phys. 1939, 47, 229–306.

26. van den Essen, A. *Polynomial Automorphisms and the Jacobian Conjecture;* Birkhäuser: Basel, Switzerland, 2000; p. 329.

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



 \odot 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).