

Table S10. Clinical variables and hematologic toxicity

Characteristics	N	Hematologic toxicity		χ^2	p-value	OR	IC _{95%}
		NO N (%)	YES (Grade 1-4) N (%)				
Gender	101						
Female	52	50(96.2)	2(3.8)				
Male	49	48(98.0)	1(2.0)				
Age diagnosis PS	101	27.1 (18.0-44.3)	29.6 (24.2-34.3)				
Family History of Ps	101						
Yes	52	49 (94.2)	3 (5.8)				
No	49	49 (100.0)	0 (0.0)				
Smoking	101						
Smoker	31	31 (100.0)	0 (0.0)				
Non-smoking	49	46 (93.9)	3 (6.1)				
Former Smoker	21	21 (100.0)	0 (0.0)				
Alcoholic drinking	101						
Drinker	38	38 (100.0)	0 (0.0)				
Non-drinker	61	58 (95.1)	3 (4.9)				
Former Drinker	2	2 (100.0)	0 (0.0)				
Type of Psoriasis	101						
Plaque	74	72(97.3)	2(2.7)				
Pustular	5	5(100.0)	0(0.0)				
Inverse	1	1(100.0)	0(0.0)				
Guttate	5	4(80.0)	1(20.0)				
Plaque and guttate	12	12(100.0)	0(0.0)				
Plaque and inverse	2	2(100.0)	0(0.0)				
Plaque and pustular	1	1(100.0)	0(0.0)				
Plaque, guttate and inverse	1	1(100.0)	1(0.0)				
Localization							
Trunk and limbs	101						
Yes	93	90(96.8)	3(3.2)				
No	8	8(100.0)	0(0.0)				
Scalp and face	101						
Yes	77	74(96.1)	3(3.9)				
No	24	24(100.0)	0(0.0)				
Nails	101						
Yes	58	56(96.6)	2(3.4)				
No	43	42(97.7)	1(2.3)				
Palmoplantar	101						
Yes	19	19(100.0)	0(0.0)				
No	82	79(96.3)	3(3.7)				
Flexures	101						
Yes	28	27 (96.4)	1 (3.6)				
No	73	71 (97.3)	2 (2.7)				
Development of PSA	101						
Yes	31	29(93.5)	2(6.5)				
No	70	69(98.6)	1(1.4)				
Comorbidities	101						

Yes	57	55(96.4)	2(3.5)	-	1*	-	-
No	44	43(97.7)	1(2.3)				
Age of onset of MTX	101	45.77±14.94	40.33±8.50	-	0.385	-	-
MTX therapy duration (months)	101	14.5 (5.0-35.3)	15.0 (8.5-19.5)	-	0.253	-	-
MTX Administration	101						
Oral	47	46 (97.9)	1 (2.1)	-	1*	-	-
Subcutaneous	30	29 (96.7)	1 (3.3)				
Both	24	23 (95.8)	1 (4.2)				
Type of MTX therapy	101						
Monotherapy	93	90 (96.8)	3 (3.2)	-	1*	-	-
Combination Therapy	8	8 (100.0)	0 (0.0)				
Maximum MTX dose (mg/week)	101	12.5 (10.-15.0)	15.0 (10.0-15.0)	-	0.756	-	-
Therapeutic adherence							
Adherent	70	68 (97.1)	2 (2.9)	-	1*	-	-
Intentional non-adherent	30	29 (96.7)	1 (3.3)				
Unintentional non-adherent	1	1 (100.0)	0 (0.0)				

*p-value for the Fisher's test. PS: psoriasis; PSA: psoriatic arthritis