

Processing and Properties Analysis of Grain Foods

Yonghui Li ^{1,*}  and Xiaorong Wu ² ¹ Department of Grain Science and Industry, Kansas State University, Manhattan, KS 66506, USA² Grain Quality and Structure Research Unit, CGAHR, USDA-ARS, 1515 College Ave., Manhattan, KS 66502, USA

* Correspondence: yonghui@ksu.edu; Tel.: +1-785-532-4061; Fax: +1-785-532-7010

Foods from grains and grain-derived ingredients are among the most important energy and nutrient source for humans. Cereals (e.g., wheat, rice, corn) have been the conventional food materials, while interest is growing in utilizing pseudocereals (e.g., buckwheat, quinoa, amaranth), pulses (e.g., dry peas, chickpea, dry beans), oilseeds (soybean, peanut, hempseed), and other grains for the development of various foods and food ingredients. Grain and food processing converts raw grains to functional ingredients and produces palatable and nutritious end-products. Examples of grain-related processes include, but are not limited to, drying, milling, fractionation, hydration, fermentation, extrusion, cooking, baking, frying, steaming, freezing, etc. Various physical and chemical changes and interactions are expected during these processes, which further affect the nutritional, textural, sensory, and many other quality properties of the products. Methods, protocols, and equipment have been developed to process, monitor, and control the processing parameters to achieve the desired end-product quality or functionality. This Special Issue of *Processes* on “Processing and Property Analysis of Grain Foods”, and the associated Special Issue reprint, published 15 research papers from lead scientists and researchers in the area and covered diverse topics related to grain-processing innovations and the effect of both conventional and innovative grain processes on the properties of grain-derived ingredients, intermediates, and end products. The Special Issue is available online at: https://www.mdpi.com/journal/processes/special_issues/Grain_Foods_Processing (accessed on 25 December 2022).

Sorghum (*Sorghum bicolor* (L.) Moench) ranks fifth in global cereal grain production. Starch is the primary constituent of sorghum grain, and its content and properties, especially the amount of amylose content in sorghum starch, influence the suitability of sorghum cultivars for specific end uses. Peiris et al. successfully developed partial least squares NIR (near infrared spectra) models to estimate starch and amylose contents in intact grain sorghum samples [1]. The newly developed calibrations can be used as a rapid screening tool for characterizing sorghum starch composition in segregating populations and to identify germplasms for developing new cultivars and hybrids for specific end uses.

Reducing the particle size of cereal grains is often the first step in the food and feed manufacturing process. Pulses are attracting an increasing interest due to their multiple agronomic and nutritional advantages. Siliveru and colleagues evaluated roller milling to produce chickpea, lentil, and yellow pea flours and characterized the particle and physicochemical properties of the produced flours [2,3]. The findings may assist millers to adapt pulse-milling technologies with minor modifications to their existing wheat milling facilities and provide guidance for more suitable uses of pulses.

Hammermills are among the most commonly used size-reduction equipment in the feed industry because of their high throughput and versatility in grinding different materials. Yellow dent #2 corn is a common grain type used for feed in the U.S. Paulk and colleagues investigated the effects of whole-corn moisture before grinding and hammermill screen size on subsequent ground corn moisture, particle size, and flowability [4]. They further evaluated the effects of hammermill tip speed, assistive airflow, and screen sizes on



Citation: Li, Y.; Wu, X. Processing and Properties Analysis of Grain Foods. *Processes* **2023**, *11*, 95. <https://doi.org/10.3390/pr11010095>

Received: 25 December 2022

Accepted: 27 December 2022

Published: 29 December 2022



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

hammermill throughput and characteristics of ground corn [5]. The studies provide useful references for processes involving grinding operations.

Innovations in food extrusion technology are enabling its rapid expansion and applicability in diverse areas related to bioprocessing and value addition. McGuire et al. compared and characterized the flowability of some common powder ingredients (corn, wheat, and sucrose) in the extrusion process [6]. The study related raw material particulate rheology to the granular flow in a pilot-scale single-screw food extruder, and some findings were visualized using a transparent plexiglass window during extrusion.

Dough mixing is an important process that can affect the quality of end products. Elucidating the mechanisms underlying gluten formation and structure remains challenging. Iwaki et al. employed the front-face fluorescence method to assess changes in hydrophobic interactions among gluten proteins during dough formation by extracting proteins in different concentrations of 1-propanol solution [7].

Pulse flours are commonly added to food products to improve their functional properties, nutritional quality, and health benefits. Zhang et al. investigated the effects of the partial replacement (0–25%) of whole wheat flour with diversified whole pulse flours (yellow pea, green pea, red lentil, and chickpea) on dough properties and bread quality [8]. Among all the tested pulse flours, the composite flour containing yellow pea flour or chickpea flour showed overall better potential for bread-making with good dough-handling properties and product quality. Beans are also important pulse grains for food uses. Mariscal-Moreno et al. investigated the effects of the partial substitution of wheat flour by ayocote bean and black bean in bread production on the thermal properties of composite flours, the bread's proximal composition, in vitro protein digestibility (IPD), as well as color and sensorial parameters of the final products [9]. These studies on composite flours could lead to the development of more nutritious bread products by complementing cereal and pulse ingredients.

Noodles are a staple food in many Asian countries and are widely consumed throughout the world because of their convenience and palatability. Cheng et al. investigated the effects of gaseous chlorine dioxide treatment on the physicochemical properties of buckwheat-based composited flour (buckwheat–wheat–gluten) and shelf-life, textural qualities, and sensory properties of fresh buckwheat noodles [10]. The study reveals the effects of gaseous chlorine dioxide treatment on Tartary buckwheat flour properties and the shelf-life of the noodles. Park et al. explored the noodle-making performance of flour blends with different particle sizes and blending ratios of purple-colored wheat bran and their antioxidant properties [11]. Noodle quality and antioxidant activity were more significantly enhanced by small bran particles at higher blending ratios.

Last but not least, this Special Issue also published interesting papers addressing the physical characteristics of typical maize seeds in a cold area of North China based on principal component analysis [12], the spatiotemporal characteristics of heat resource effectiveness in the southern rice cropping area in China and relationships between heat resource effectiveness and rice potential yield, as well as grain yield reduction rate [13], the effect of different alternative sweeteners that contained sugar alcohols or bulking agents on the physicochemical properties of rolled “sugar” cookie [14], and the effects of different amylose contents of foxtail millet varieties on textural properties of Chinese steamed bread [15].

Author Contributions: Both authors contributed equally. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported in part by the USDA Pulse Crop Health Initiative projects (Grant Accession No. 0439205 and No. 0439200) and the USDA National Institute of Food and Agriculture Hatch project (Grant Accession No. 7003330).

Acknowledgments: This is contribution No. 23-156-J from the Kansas Agricultural Experimental Station. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture. The USDA is an equal opportunity provider and employer.

Conflicts of Interest: The authors declare that they have no known conflict of interest.

References

1. Peiris, K.H.S.; Wu, X.; Bean, S.R.; Perez-Fajardo, M.; Hayes, C.; Yerka, M.K.; Jagadish, S.V.K.; Ostmeyer, T.; Aramouni, F.M.; Tesso, T.; et al. Near Infrared Spectroscopic Evaluation of Starch Properties of Diverse Sorghum Populations. *Processes* **2021**, *9*, 1942. [\[CrossRef\]](#)
2. Martin, R.; Siliveru, K.; Watt, J.; Blodgett, P.; Alavi, S. Pilot Scale Roller Milling of Chickpeas into a De-Hulled Coarse Meal and Fine Flour. *Processes* **2022**, *10*, 2328. [\[CrossRef\]](#)
3. Pulivarthi, M.K.; Nkurikiye, E.; Watt, J.; Li, Y.; Siliveru, K. Comprehensive Understanding of Roller Milling on the Physicochemical Properties of Red Lentil and Yellow Pea Flours. *Processes* **2021**, *9*, 1836. [\[CrossRef\]](#)
4. Braun, M.; Dunmire, K.; Evans, C.; Stark, C.; Woodworth, J.; Paulk, C. Effects of Grinding Corn with Different Moisture Content on Subsequent Particle Size and Flowability. *Processes* **2021**, *9*, 1372. [\[CrossRef\]](#)
5. Braun, M.; Wecker, H.; Dunmire, K.; Evans, C.; Sodak, M.W.; Kapetanovich, M.; Shepherd, J.; Fisher, R.; Coble, K.; Stark, C.; et al. Evaluation of Hammermill Tip Speed, Air Assist, and Screen Hole Diameter on Ground Corn Characteristics. *Processes* **2021**, *9*, 1768. [\[CrossRef\]](#)
6. McGuire, C.; Siliveru, K.; Ambrose, K.; Alavi, S. Food Powder Flow in Extrusion: Role of Particle Size and Composition. *Processes* **2022**, *10*, 178. [\[CrossRef\]](#)
7. Iwaki, S.; Hayakawa, K.; Fu, B.-X.; Otobe, C. Changes in Hydrophobic Interactions among Gluten Proteins during Dough Formation. *Processes* **2021**, *9*, 1244. [\[CrossRef\]](#)
8. Zhang, Y.; Hu, R.; Tilley, M.; Siliveru, K.; Li, Y. Effect of Pulse Type and Substitution Level on Dough Rheology and Bread Quality of Whole Wheat-Based Composite Flours. *Processes* **2021**, *9*, 1687. [\[CrossRef\]](#)
9. Mariscal-Moreno, R.M.; Chuck-Hernández, C.; Figueroa-Cárdenas, J.D.; Serna-Saldivar, S.O. Physicochemical and Nutritional Evaluation of Bread Incorporated with Ayocote Bean (*Phaseolus coccineus*) and Black Bean (*Phaseolus vulgaris*). *Processes* **2021**, *9*, 1782. [\[CrossRef\]](#)
10. Cheng, Z.; Li, X.; Hu, J.; Fan, X.; Hu, X.; Wu, G.; Xing, Y. Effect of Gaseous Chlorine Dioxide Treatment on the Quality Characteristics of Buckwheat-Based Composite Flour and Storage Stability of Fresh Noodles. *Processes* **2021**, *9*, 1522. [\[CrossRef\]](#)
11. Park, G.; Cho, H.; Kim, K.; Kweon, M. Quality Characteristics and Antioxidant Activity of Fresh Noodles Formulated with Flour-Bran Blends Varied by Particle Size and Blend Ratio of Purple-Colored Wheat Bran. *Processes* **2022**, *10*, 584. [\[CrossRef\]](#)
12. Tang, H.; Xu, C.; Jiang, Y.; Wang, J.; Wang, Z.; Tian, L. Evaluation of Physical Characteristics of Typical Maize Seeds in a Cold Area of North China Based on Principal Component Analysis. *Processes* **2021**, *9*, 1167. [\[CrossRef\]](#)
13. Ye, Q.; Yang, X.; Xie, W.; Yao, J.; Cai, Z. Effect of Heat Resource Effectiveness Change on Rice Potential Yield in Southern China. *Processes* **2021**, *9*, 896. [\[CrossRef\]](#)
14. Heermann, M.L.; Brown, J.; Getty, K.J.K.; Yucel, U. Assessing Functionality of Alternative Sweeteners in Rolled “Sugar” Cookies. *Processes* **2022**, *10*, 868. [\[CrossRef\]](#)
15. Li, S.; Zhao, W.; Min, G.; Li, P.; Zhang, A.; Zhang, J.; Wang, Y.; Liu, Y.; Liu, J. Effects of Different Amylose Contents of Foxtail Millet Flour Varieties on Textural Properties of Chinese Steamed Bread. *Processes* **2021**, *9*, 1131. [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.