

Supporting Information

A Method to Separate Two Main Antioxidants from *Lepidium latifolium* L. Extracts Using Online Medium Pressure Chromatography Tower and Two-Dimensional Inversion/Hydrophobic Interaction Chromatography Based on Online HPLC-DPPH Assay

Xiu Wang^{a,b}, Yupei Zhang^{a,b}, Nan Wu^{a,b}, Jingya Cao^{a,b}, Yanduo Tao^a, Ruitao Yu^{a*}

^aKey Laboratory of Tibetan Medicine Research, Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xining, Qinghai, China

^bUniversity of Chinese Academy of Science, Beijing, China

***Corresponding authors at:** Key Laboratory of Tibetan Medicine Research, Northwest Institute of Plateau Biology, Chinese Academy of Sciences, No. 23 Xinning Road, Xining, China.

Tel: 13897406734

E-mail: yuruitao521@163.com.

Figure S1. ESI mass spectrum of Fr4-1-1




Figure S2. ^1H NMR Spectrum (600 MHz) of Fr4-1-1 (in MeOD- d_4)




Figure S3. ^{13}C NMR Spectrum (600 MHz) of Fr4-1-1 (in $\text{MeOD-}d_4$)




Figure S4. HMBC Spectrum (600 MHz) of Fr4-1-1 (in MeOD-*d*₄)




Figure S5. ESI mass spectrum of Fr4-1-2




Figure S6. ^1H NMR Spectrum (600 MHz) of Fr4-1-2 (in $\text{MeOD-}d_4$)




Figure S7. ^{13}C NMR Spectrum (600 MHz) of Fr4-1-2 (in $\text{MeOD-}d_4$)




Figure S8. HMBC Spectrum (600 MHz) of Fr4-1-2 (in MeOD-*d*₄)

