

Review

Dental Infection and Resistance—Global Health Consequences

Mainul Haque ^{1,*}, Massimo Sartelli ² and Seraj Zohurul Haque ³

¹ Unit of Pharmacology, Faculty of Medicine and Defence Health, Universiti Pertahanan Nasional Malaysia (National Defence University of Malaysia), Kem Sungai Besi, 57000 Kuala Lumpur, Malaysia

² Department of Surgery, Macerata Hospital, via Santa Lucia 2, 62100 Macerata, Italy; massimosartelli@infectionsinsurgery.org

³ Ninewells Hospital & Medical School, Dundee DD1 9SY, Scotland, UK; szzhaque@gmail.com

* Correspondence: runurono@gmail.com; Tel.: +60-109-265-543

Received: 4 December 2018; Accepted: 20 February 2019; Published: 1 March 2019



Abstract: Antibiotics are widely used in dental caries and another dental related issues, both for therapeutic and prophylactic reasons. Unfortunately, in recent years the use of antibiotics has been accompanied by the rapid emergence antimicrobial resistance. Dental caries and periodontal diseases are historically known as the top oral health burden in both developing and developed nations affecting around 20–50% of the population of this planet and the uppermost reason for tooth loss. Dental surgeons and family practitioners frequently prescribed antimicrobials for their patients as outpatient care. Several studies reported that antibiotics are often irrationally- and overprescribed in dental diseases which is the basis of antimicrobial resistance. The aim of this review is to evaluate the use of antibiotics in dental diseases. Almost certainly the promotion of primary oral health care (POHC) in primary health care program especially among the least and middle-income countries (LMIC) may be the answer to ensure and promote rational dental care.

Keywords: dental; oral; maxillofacial; infection; antibiotic; antimicrobial; resistance pattern; epidemiology; common microorganism; nutrition component; biofilm formation; morbidity; mortality; increase healthcare cost

1. Introduction

Penicillin was the first antimicrobial agent discovered in 1928 [1] and the first patient was treated with this newly discovered medicine in 1942 [2]. The therapeutic potential was soon well recognized. World War II promoted extraordinary cooperation between two nations situated either side of the Atlantic, United States and Great Britain, resulting in unbelievably efficient bulk production of penicillin by 1943 [1,3]. Dutch scientists during post World War II developed a new method of producing penicillin which was marketed in 1946, and ultimately ensured mass production of penicillin for the medicine market and reduced the cost of treatment [3,4]. Antibiotics have transformed modern medicine and countless lives have been saved through their use over the years. The invention antibiotic was a turning point for medical science and the quality of human healthcare. Unfortunately, the use of this miracle medicine has been accompanied by the rapid emergence of resistant strains due to the unstoppable spread of antibiotic resistance genes among the microbial community [5,6]. The microbial community develop defense strategies for their existence against the antimicrobial assault, which is a prime instance of microbiological metamorphosis, transformation, and the zenith of evolution. “An immense genetic plasticity of bacterial pathogens that trigger specific responses that result in mutational adaptations, acquisition of genetic material or alteration of gene expression producing resistance to virtually all antibiotics currently available in clinical practice” is considered

as the natural law [7]. In the early 1950s, Japan experienced an epidemic of Shigellosis caused by antibiotic resistance strains of *Shigella*. A few years later Japanese scientists described that resistance to “multiple antibiotics not only developed quickly and simultaneously, but also seemed to transfer from resistant to sensitive strains” [8]. Additionally, before penicillin was marketed for treatment, a bacterial penicillinase (β -lactamases) was recognized by Abraham and Chain, two members of the penicillin invention squad [9]. It is interesting that microbes developed their existence technology, such as β -lactamases, before antibiotics were ever used. A recent database voiced the presence of more than 20,000 possible resistance genes (r genes) of strictly 400 different types, as analyzed from available bacterial genome sequences [10,11]. Microorganism resistances to antimicrobials now extend to all known categories of agents, either natural or synthetic. Antimicrobial resistance (AMR) has not only stalled treatment approaches to infectious diseases but has also influenced healthcare policy and planning [10]. Subsequently, medical specialists are facing the possibility that human beings may return to the pre-antibiotic era of medical care [5] as AMR is threatening not only effective treatment towards bacterial diseases but also other microbes such as parasites, viruses, and fungi [12,13]. There were a lot of reports that dental surgeons often prescribed large amounts of antimicrobials. Often these antibiotics were overprescribed, or prescribed imprudently and irrationally, which in turn promotes microbial resistance. This manuscript will try to discuss issues related to dental infection and resistance and its impact on global health consequences.

2. Materials and Methods

This review has been based on freely accessible literature from Google, Google Scholar, and PubMed, and from the link provided by the Universiti Pertahanan Nasional Malaysia ((UPNM) National Defence University of Malaysia). The terms used were dental, oral, maxillofacial infection, resistance pattern, epidemiology, common microorganism, nutrition component, biofilm formation, morbidity, mortality, and increase healthcare cost. A few manuscripts incorporated required payment to view the full paper but have been provided free of charge by other libraries as part of cooperation with UPNM. This is a narrative review article will give effort to describe and deliberate issues related to dental infection and resistance and its impact on global health significances, from a theoretical and contextual point of view, based on previously published manuscript. Additionally, there has been no attempt to utilize specific a data base and to use a methodological approach to develop a systematic review and meta-analysis.

3. Epidemiology of Dental Infection

Dental caries and periodontal diseases are historically known as the top oral health burden in both developing and developed nations affecting around 20–50% of the population of this planet, and is the uppermost reason for tooth loss [14–16]. Among Indian patients over 30 years of age it has been reported that the significant (almost 80%) cause of loss of teeth is due to periodontal disease [17,18]. Multiple studies reported that in Asia, the Middle East, and across the African sub-Saharan regions, dental caries is a principal public oral health threat [19–21]. Though some studies state that globally dental caries has declined in the population especially in modern countries [21–23], other recent research reports that there is a significant increase in rates of caries [24,25].

Another Indian study said that periodontal disease is widespread in the Eastern Indian state of West Bengal [26]. One Libyan study conducted among a population of 1225 subjects aged 18–34 years, revealed that only 5% population had healthy periodontium [27]. Multiple studies reported that periodontal disease occurrence and severity increases with patients' age [28–31]. Additionally, periodontal diseases occur more often in males and individuals with poor oral hygiene practice [32–34]. Pediatric and adult groups of males were more often sufferers of dental infections and other infections of the oral cavity in comparison with the female population of both rural and urban communities [35–37]. One Dutch study revealed that the incidence of caries decreased in the period of 1990–2009 among an 8–21-year-old studied population in both low and high socioeconomic cluster

and was also statistically significant [38]. Similarly, another British study conducted among 69,318 children aged 5–15 years revealed a 31–51% reduction of caries in the last 40 years (1973–2013). This study also reported a greater reduction among 15-years age group. Additionally, the British study concluded that although there is a significant improvement, caries continues to be a significant liability to the national healthcare system [39]. In Norway, prevalence of caries decreased from 81% to 52.2% in a 15-year period (1985–2000) among 12-year-old children. However, the next 4 years (2001–2004) the prevalence increased to 59.8% with a 3.3% annual growth rate. Nevertheless, the decayed, missing, and filled teeth (DMFT) index kept on steady around 1.6 from 1997 to 2004 [40]. A similar DMFT was also observed in other Scandinavian countries [41]. The DMFT index has been in use by dental surgeons for nearly 80 years and persists as the most reliable tool to assess dental caries for epidemiological study [42]. Another research among 28,522 German first grade children revealed that 25.9% had caries and required treatment. This study examined in further detail 25,020 children and reported that girls possess better oral and dental health when compared to boys regarding decayed missing and filled primary teeth (DMFT). There was a statistically significant ($p < 0.001$) difference between DMFT score among sexes. Boys DMFT score was double than that of girls [43].

4. Common Microorganisms Involved in Dental Infection

Periodontal disease and dental caries are the most frequently reported chronic infective dental diseases and are caused by microorganisms living in the mouth cavity [44,45]. The human oral flora encompasses over 700 microorganisms, and 50% of those microorganisms are uncultivable microbes [45]. The different anatomic areas and its bathing fluid, saliva of the oral cavity, possess oral microbiome comprising the normal oral flora which includes bacteria, archaea, fungi, protozoa, and viruses. These oral microbiomes usually swim in saliva as free-floating microorganisms and form a complex ecological community of biofilm, attaching to different surfaces of the mouth cavity [46,47]. Biofilm is often responsible for several local and systemic diseases [47].

One Japanese study reported that diverse infection of strict anaerobes with facultative anaerobes, especially commensal streptococcal gram-positive bacteria known as viridans streptococci, was identified as the principal pathogens responsible for dentoalveolar infections, periodontitis, and pericoronitis [48]. Another Japanese study identified *Fusobacterium*, *Peptostreptococcus micros*, *Porphyromonas* species, and *Prevotella* species as culprits for dentoalveolar infection [49]. Additionally, 34% of *Prevotella* species were found to produce beta-lactamase [49]. *Peptostreptococcus micros* was also identified by another group of researchers as responsible for progressive periodontitis [50]. Multiple studies revealed polymicrobial flora with frequent involvement of Gram-negative anaerobic pathogens [51–56]. Similar other studies reported that 90% of root canal infections were due to obligatory anaerobic microbes [52,57].

5. Nutrition Component That Promotes Dental Infection

Diet also plays a substantial role in dental caries and enamel erosion [20,58,59]. “Caries is demineralization of the inorganic part of the tooth with the dissolution of the organic substance due to a multifactorial etiology” [60]. Other researchers define “dental caries is a multifactorial disease that results from interactions among a susceptible host, caries-related bacteria, and cariogenic diets” [61]. The demineralization of the dental enamel and dentine by organic acids that form in the dental plaque are because of anaerobic microorganisms metabolizing sugars of the diet [62]. Dietary machinery especially dietary acids contribute to the development of enamel defects (e.g., enamel hypoplasia, fluorosis) [58]. Soft drinks contain both acids and sugars, have the acidogenic and cariogenic property that can lead to dental caries and enamel erosion. Several research studies revealed a positive correlation between drinking soft drinks and dental caries and erosion [63–65]. One study conducted among children of 2–10 years revealed that those children who consumed a high volume of carbonated soft drink also had a significantly higher rate of dental caries than children who had high juice, high milk, and high water in their diet [66]. Additionally, one British

study reported that boys of 14–15 years take much more sugary drinks than that of their female counterpart. Two or more glasses of sugary drink have a statistically significant association with dental caries. The high consumption of sugary drink was found to have a correlation with family income, gender, and mother's education level [67]. Another American study suggests that present-day high consumption of fizzy drink among children, above all the escalation in "soda pop" drinking, were responsible for an intensification of dental caries rates [68]. This study and other research additionally revealed that poor intakes of micronutrients such as riboflavin, copper, vitamin D, and vitamin B₁₂ are associated with an increased rate of caries and gingival diseases [68,69]. Low intake of milk and other dairy products has been associated with carries among children [68]. Other comparable studies also similarly condemned that high intake of non-alcoholic carbonated drinks and low consumption of milk and other dairy products among children and adolescents increases the risk for dental caries and other systemic diseases [63,70,71]. Multiple studies reported that drinking milk ensures the consumption of many macronutrients and micronutrients which in turn promotes and protects health from various diseases [72–74]. Very low-level sugar consumption was observed among the communities of Inuit of Alaska, Ethiopia, Ghana, Nigeria, Sudan, and the islands of Tristan da Cunha and Sant'Elena. These populations often lead traditional lifestyles and rate of carries were found to be very negligible. Researchers observed that as the economic condition of these geographical areas improve, the amount of sugar and other fermentable carbohydrate content increases in the diet, and this paralleled the increases rate of caries [62,75–77]. Additional studies revealed that frequent consumption of fermentable sugars promotes caries formation [78,79], as sugar such as sucrose when fermented to produce lactic acid which lowers pH which ultimately disturbs the environment and damages normal the demineralization and remineralization process, consequentially results in dental enamel destruction through demineralization [78,80].

6. Biofilm Formation among Dental Infection

Researchers with the aid of advanced technology have identified a rich microbial consortium and around 1000 diverse microbial species, which include bacteria, virus, and fungi in the dental biofilm. Biofilms are a structurally and functionally organized community of microorganisms with a three-dimensional configuration and are surrounded in a matrix of extracellular material [81–83]. Among microbes existing in the oral biofilm, many of them are innocent natural inhabitants. Nevertheless, some of them have the inherent quality of damaging the mineralized and soft tissues of teeth and gum [83]. Antonie Philips van Leeuwenhoek, a Dutch scientist in the seventeenth century, first reported regarding observed dental plaque and oral biofilms, and plaque is responsible for caries, and periodontal disease [84]. Multiple studies reported that more than 10¹¹ microorganisms per mg of dental plaque were found [45,85,86]. Several studies reported that the mouth cavity is rich in nutrition, temperature 35/36 °C, pH 6.75–7.25 are the factors which all encourage the growth of many microorganisms and the formation of biofilm [87–89]. Just after tooth brushing and rinsing, the mouth cavity salivary proteins form a thin film over the tooth surfaces. Microorganisms with their microfilaments attach to this thin protein film. If the oral environment alters to become favorable for microorganisms, their population increases in number and communicates by secreting signal molecules and creates a community [90,91]. "The microbes secrete proteins, polysaccharides, nucleic acids, and other substances to the extracellular matrix, additionally containing proteins and nutrients from saliva" [83]. Thereafter, forms the matrix of the biofilm [81].

The principal microbial species found in the plaque of diseased areas are dissimilar from those of healthy areas. This is even though at the healthy locations of the mouth cavity, disease producing microbes are still regularly spotted but in lower numbers [92]. The former suppositions regarding the cause of the dental caries were believed to be due to the increased number of microorganisms in plaque "(Non-specific Plaque Hypothesis)," or that plaque contains particular types of cariogenic bacteria, such as *Streptococcus mutans* and *Streptococcus sobrinus*, within the plaque "(Specific Plaque Hypothesis) [85,93,94]. At present, the fundamental concepts of the earlier two hypotheses were

reunited and widely accepted as the “Ecological Plaque Hypothesis.” The current hypothesis of caries formation describes that the normal oral microbial environment, if stressed, alters towards certain disease-related micro-organisms [61,95–97].

7. Antibiotic Use and Resistance Pattern of Microorganism in Dental Infections

The appearance of penicillin in the market and clinical use has saved many lives throughout the planet, transformed medical science and its’ success in treating infectious disease [3,98,99]. Poor use of antimicrobials promotes the emergence of antibiotic-resistant microbial strains [100,101], increases the possibility of antibiotic-associated adverse reactions [102], and characterizes a waste of healthcare funds [103]. Antimicrobial resistance is a grave public health and patient care issue. Antimicrobial resistant infections are exceedingly problematic in regards to the treatment of infectious diseases. Consequently, prudent use of antimicrobials demands the highest priority to prevent antimicrobial resistance, ensure patient safety and improve healthcare [103]. Multiple studies reported that dental surgeons frequently prescribed inappropriate antibiotics which ultimately promote antimicrobial resistance [103–106].

Antibiotics are widely used in dental caries and another dental related issues, both for therapeutic and prophylactic reasons [107]. Dental surgeons frequently prescribe antibiotics with apprehension that oral cavity normally contains huge number microorganism as normal flora which can cause infections in their patients [108]. It has been reported that antibiotic prophylaxis within 2 hours of dental procedures were beneficial [109], however, antibiotics taken after 4 hours had no benefit achieved [110]. Procedures such as periodontal surgery, scaling and professional teeth cleaning can cause significant bleeding from hard or soft tissues, and such bleeding cannot always be anticipated for prior to the procedure [110]. Another group of scientists believe that “indications for the use of systemic antibiotics in dentistry are limited since most dental and periodontal diseases are best managed by operative intervention and oral hygiene measures” [111]. Currently enormous amounts of antibiotics are prescribed and consumed because of dental infective disorders around the planet [106,112,113]. Regular and frequent use of antibiotics in dental infection often causes long-term public health troubles by leading to the development of resistant microbes including multidrug-resistant pathogens. This may be all the more tragic if there was minimal benefit to patient care by giving the antibiotic in these circumstances [114,115].

One study conducted in India comprising of 68 patients (18–58 years old) examined microbial specimens and reported that “a total of 64 aerobic and 87 anaerobic strains were isolated. The predominant bacteria were *Streptococci viridans* (64%), *Prevotella* (43%), *Peptostreptococcus* (26%), *Porphyromonas* (7%), and *Fusobacterium* (14%). The anaerobic Gram-negative bacilli (40%) were the predominant organisms followed by aerobic gram-positive cocci (34%). Four strains of *Candida albicans* were also identified” [116]. *Streptococci viridans* were highly sensitive to amoxicillin-clavulanate (95%), amoxicillin (90%), and levofloxacin (83%). Similarly, *Prevotella* were highly susceptible to amoxicillin-clavulanate (97%) and less sensitive to erythromycin (62%). Again, *Peptostreptococcus* and *Porphyromonas* are highly sensitive to amoxicillin-clavulanate (100%) and clindamycin (100%) [116]. The identified microorganisms were extremely sensitive to the commonly prescribed antimicrobials, e.g., amoxicillin + clavulanate combination and amoxicillin alone, clindamycin, and levofloxacin, but were more resistant towards erythromycin [116]. Another British study reported that an acute dental abscess was typically polymicrobial, with facultative anaerobes, such as viridans group streptococci and the *Streptococcus anginosus* cluster, with largely strict anaerobes, such as anaerobic cocci, *Prevotella* and *Fusobacterium* species. The highly sophisticated technology, based on non-culture techniques, identified a diverse community of pathogens, e.g., *Treponema* species and anaerobic Gram-positive rods such as *Bulleidia extructa*, *Cryptobacterium curtum*, and *Mogibacterium timidum* [117]. Macrolide antimicrobials were documented as less sensitive towards the ‘viridans group streptococci,’ anaerobic streptococci, *Streptococcus oralis*, and *Prevotella* species [48,118–120]. Another study reported that penicillin shows less reduced susceptibility towards the mitis streptococci group than the anginosus group [121].

The Infection Research Group of Glasgow Dental Hospital and School studied among 155 viridans group streptococci for their “minimum inhibitory concentrations (MICs) to penicillin, amoxicillin, ceftriaxone, erythromycin, clindamycin, rifampicin, vancomycin, and teicoplanin” and revealed that “27% of *S. oralis* were resistant to penicillin, 51% resistant to erythromycin, and 6% resistant to clindamycin. Further, 11% of *S. mitis* were resistant to penicillin, 40% resistant to erythromycin and 3% resistant to clindamycin.” Penicillin-resistant pathogens were also shown to be less sensitive towards other antimicrobials [122]. Researchers of the Barkatullah University, India studied gram-positive cocci (e.g., *Streptococcus mutans*, *S. sobrinus*, *S. oralis*, *S. sanguinis*) and bacilli (*Lactobacillus acidophilus*, *L. rhamnosus*, *L. fermentum*). These pathogens show resistance for each antibiotic in this research as follows: “penicillin V: 72/150 (48%), tetracycline: 99/150 (66%), amoxicillin: 135/150 (90%), cloxacillin: 117/150 (78%), and erythromycin: 90/150 (60%) [123].” Another study from Nepal identified that 91% of invading pathogens were gram-positive and 9% were gram negative. *Streptococcus mutans* were resistant to penicillin (66.15%), tetracycline (60.76%), and cotrimoxazole (20%). *S. aureus* was found to be resistant towards penicillin (91.48%), tetracycline (86.17%) and ampicillin (61.70%). *S. mitis* was resistant to tetracycline (78.12%) and ciprofloxacin (65.62%). *Pseudomonas spp.* were 100% resistant to tetracycline and cotrimoxazole was 90.90% [124]. Janaki medical school researchers reported that from bacterial isolates, 90% were Gram-positive. Among that Gram-positive microorganisms were *S. mutans*, *S. aureus*, *S. mitis*, *S. albus*, and *S. vestibularis*. The remaining 8.48% were Gram-negative microbial isolates. Again among Gram-negative: *Pseudomonas spp.*, *K. pneumoniae*, *P. vulgaris*, and *Enterobacter spp.* Among microbiological isolates were sucrose fermenter and non-sucrose fermenter 93.84% and 6.15% respectively. Documented gram-positive microorganisms were sensitive towards ciprofloxacin, gentamicin, and erythromycin 94.27%, 51.85%, 49.49% respectively. The gram-negative organisms was sensitive to ciprofloxacin, imipenem and gentamicin, and ceftriaxone 100%, 89.28%, and 50% respectively [125]. Another study from Brazil revealed that oral pathogens were highly resistant to ampicillin, amoxicillin+clavulanic acid, cefoxitin, cephalothin, amikacin, chloramphenicol, and nalidixic acid. This study also reported that carbapenems (meropenem and imipenem) were the most active antimicrobial agents and 1.6–2.3% pathogens showed resistance toward these medicine. Low resistance profile was also observed with ciprofloxacin and rifampin [126]. A Mexican study conducted among 60 children with active infections in the primary dentition revealed that Clindamycin in 8 lg/mL and 16 lg/mL exhibited the maximum (85.9%) microbial resistance followed by amoxicillin (43.7%) and amoxicillin-clavulanic acid (12.0%). *Streptococcus oralis* and *Prevotella intermedia*; *Treponema denticola* and *Porphyromonas gingivalis*; *Streptococcus mutans*; *Campylobacter rectus*; and *Streptococcus salivarius* were found 75%, 48.3%, 45.0%, and 40% respectively to be the most predominantly resistant microorganisms identified through polymerase chain reaction [127].

A composite microbial community exists normally in the mouth cavity which contains of diverse bacterial and fungal classes, their accompanying biofilms, and often with produce cytokine which promote relentless inflammation. These various infectious consequences of poor oral health among different age group of patients [92,128]. The Mayo Clinic reported that oral health had potential to enhance several diseases process which include endocarditis, cardiovascular diseases, pregnancy and birth, diabetes mellitus, HIV/AIDS, osteoporosis, Alzheimer’s disease, eating disorders, rheumatoid arthritis, head and neck cancers, and Sjogren’s syndrome [129]. It has been recommended that prescribing practice especially regarding antimicrobials need to be improved among dental surgeons and all health professionals, additionally, dentals surgeons were advised to follow guidelines for prescribing, and awareness regarding antibiotic resistance need build up among common people [114,130].

8. Morbidity, Mortality, and Increase Healthcare Cost in Dental Infections

Oral health is essential component of general health and quality of life. Thereafter, the World Health Organization (WHO) defines “It is a state of being free from the mouth and facial pain, oral and throat cancer, oral infection and sores, periodontal (gum) disease, tooth decay, tooth loss, and other

diseases and disorders that limit an individual's capacity in biting, chewing, smiling, speaking, and psychosocial well-being [131]".

"Detrimental effects of oral infections on general health have been known for almost 3000 years" [132]. Chronic oral and dental disease can be very dangerous, especially among immunocompromised patients [131]. Consequently, different studies have suggested that such chronic infection promotes several life-threatening systemic diseases and increases morbidity and mortality [133–137]. Tooth loss impairs mastication and leads to higher mortality by causing poor diet, nutrition, and eating behavior [138–140]. One Japanese study, utilizing the multivariate-adjusted Cox proportional hazards model, revealed that practicing proper oral care, like that of tooth brushing, regular consultation with dental surgeons, and use of dentures, had an inverse relation with mortality among the elderly population [140,141]. Additionally, periodontal diseases and tooth loss are frequently related to coronary heart disease, stroke, pneumonia, and incident disability [142–146].

Chronic oral infections, particularly periodontitis, are the underlying cause of several fatal systemic diseases like endocarditis [86,147]. Several studies have shown that cardiovascular diseases such as coronary heart disease (CHD), stroke, peripheral vascular disease (PVD), cardiomyopathy, atherosclerosis, and myocardial infarction are linked to chronic infection and inflammation [148–153]. It has been shown that severe chronic microbial dental infections, endocarditis and meningitis are associated with cerebral infarctions among male patients [154,155]. Additionally, research studies revealed that periodontitis has a potential role in regulating many other systemic diseases such as diabetes, respiratory disease, low birth weight, pre-term infants, other pregnancy-related issues, rheumatoid arthritis, and osteoporosis [156,157]. "Several mechanisms have been proposed to explain or support such theories, and oral lesions are indicators of disease progression, and oral cavity can be a window to overall health and body systems [156]." Chronic periodontitis is responsible for producing C-reactive proteins (CRP), interleukin-1b (1L-1b), interleukin-6 (1L-6), and Tumor Necrosis Factor-alpha (TNF- α) and disseminate these inflammatory mediator substances through the human body [158,159]. One Indian study revealed that patients with periodontitis had a higher level of high-sensitive (hs)-CRP than those who were suffering from gingivitis and with healthy gingiva [160]. The Mayo Clinic reported that "a high-sensitivity C-reactive protein (hs-CRP) test, which is more sensitive than a standard test, also can be used to evaluate your risk of developing coronary artery disease, a condition in which the arteries of your heart are narrowed. Coronary artery disease can lead to a heart attack [161]." Furthermore, high level of inflammatory mediator CRP increased the possibility of the development of type 2 diabetes mellitus (T2DM) and suggested that CRP is considered as a strong independent forecaster [162–165]. One recent study of Thailand reported that CRP, IL6, and TNF- α related with the initiation process for the development T2DM. This study additionally revealed that there was a strong co-relationship between CRP and T2DM. Consequently, researchers suggested that CRP is a good biomarker for the diagnosis and assessment of T2DM [166]. Similarly, one meta-analysis comprising of 19 manuscripts involving 39,136 participants and 7924 cases reported that "T2DM risk was strongly associated with elevated levels of inflammatory cytokines (IL-1 β , IL-6, IL-18, CRP), TNF- α , and low levels of adiponectin [167]". Another research studied three important inflammatory mediators (IL-6, CRP, and TNF- α) and revealed that among the elderly population, having low levels of these inflammatory mediators were linked with low death rate and improved quality of life [168]. Multiple studies reported that elevated CRP level indicate chronic periodontal disease, and other oral mucosal diseases including denture related edentulous lesions [169,170].

Chronic periodontal diseases often serve as a reservoir site to produce inflammatory mediators. These mediators have the potential to encourage threat to the fetal-placental unit and increase the risk of adverse pregnancy outcomes [171]. One Iranian study reported that mothers delivering low birth babies (LBW) had statistically significantly higher level of diseased gingiva ($p = 0.042$), and more deep pockets ($p = 0.0006$, Mann–Whitney test) [171]. This study concluded that chronic periodontal infections are one of the possible independent risk factors for LBW [171]. Similarly, one systematic review comprising of 12 prospective studies reported that a statistically significant risk of preterm

delivery for those women suffering from periodontitis (risk ratio (RR):1.70) during pregnancy and a significant risk for LBW (RR: 2.11) [172]. Another similar systematic review and meta-analysis of case-control studies, comprising of 17 case-control studies and involving of a total of 10,148 patients, assessed periodontal disease as a risk factor for preterm birth, LBW babies. This study reported that “the estimated odds ratio was 1.78 (CI 95%: 1.58, 2.01) for preterm birth, 1.82 (CI 95%: 1.51, 1.20) for LBW and 3.00 (CI 95%: 1.93, 4.68) for preterm low birth-weight [173].” Another meta-analysis and systematic review involved 10 case-controlled studies from seven countries (India, Brazil, Iran, Argentina, Jordan, Senegal, and Tanzania) with a total of 2423 patient with a mean age ranging from 13–49 years [174]. This research meta-analysis also found that chronic periodontal infection among pregnant mothers often resulted in the delivery of a child before term and of LBW [174]. One more Indian case-control study concluded that pregnant women having chronic periodontal infections is a significant obstetric risk factor for preterm LBW. These patients of Nellore District, Andhra Pradesh, India frequently delivered preterm LBW child. This study finally commented that both antenatal status and periodontal diseases contribute to delivering preterm LBW child [175]. Age, sex, socioeconomic status, education level, race, genetics issues, CRP also had relation with dental diseases [28].

Dental diseases often remain untreated even in the USA and endure as a substantial public health threat. It has been advocated that preventive dental care improves overall oral health, but not all patients of the USA receive regular dental healthcare [176–180]. Over 10,000 patients visit five hospital systems (a total of seven hospitals) in the Minneapolis—St. Paul metropolitan area in Minnesota, USA for emergency dental care and it costs closely \$5 million per year [175]. These patients were principally treated by minimizing their acute illness such as pain but not addressing underlying pathology of the disease [180]. Another study revealed that in the USA around 35% population or 100 million had no dental coverage [181]. One study reported that pediatric patients receiving treatment for dental diseases in a hospital operating room were the US \$1508 and \$104 for hospital non-operating room [182].

One Indian study reported that dental infections are a major public health threat, having a considerable influence on the quality of life, and work performance both academic and professional [178]. This study additionally reported that an enormous amount of monetary support is required to treat caries-related issues among school children of India. Nonetheless, the Indian budget for public health care is considerably under-resourced for oral health care [183]. Furthermore, in the Indian budget there is no precise, distinct provision for oral health care [183,184]. The healthcare in Malaysia is principally provided under a publicly operated system by the ministry of health which is almost free or has a very nominal charge at point of use [185]. Private health care is also available in big cities [185]. The government of Malaysia believes that “health is a public service to be made available to everyone, with equity of access, both in geographical and cost terms. The Ministry of Health is also the lead agency in the provision of oral healthcare to the nation” [186]. Oral healthcare in Malaysia is provided both by the public and private sectors [186].

On average Brazilian people spend R\$42.19 annually on dental care which comprises dental insurance, appointments, and treatments. Again, R\$5.10 is spent on dental insurance of the total dental care expenditure and of this amount R\$4.70 was for private dental coverage. Nonetheless, only 2.5% of Brazilians spend money on dental insurance [187]. Another study reported that in Brazil dental care is traditionally based on private dental service [188].

Numerous peoples are suffering diverse issues of oral and dental diseases around the planet especially in countries having resource constraints, where government oral health care programs are limited [189]. Thereafter, the encumbrance oral diseases are very high all over the world against a low availability of skilled manpower regarding the management of oral health care [190]. At present multiple studies evidenced that there is urgent need to develop primary oral health care (POHC) programs all over the planet, especially in developing countries to promote effective oral health education and regular dental health check-up programs [189–191].

9. Primary Health Care and Primary Oral Health Care

Health is an ultimate human right which is requisite for the implementing of other issues human rights. Human rights are constitutionally preserved and ensured by almost all countries of this planet for persons and clusters against activities that restrict with their basic self-determinations and human rights. Thereafter, human rights are universal [192,193]. Primary health care ensure both health care and human rights both in developing and developed countries [194,195]. Currently, essential oral health service, regular dental care programs, and oral health education programs with intervention were found lacking throughout the globe especially in LMIC [196–198]. As POHC is an integral part of primary health care [191]. Thereafter, countries need developed POHC program to address this global issue of dental care [191].

10. Conclusions

The mouth cavity is the natural territory for various microbes. These naturally occurred microorganisms often act as a reservoir for resource for pathogenic microbes to produce local oral/dental infections and later systemic infections [199]. Antibiotics are often prescribed and used unnecessarily and excessively, which in turn contributes to the development of resistant microbes [200–203]. Multiple studies reported that at least 30–50% of antimicrobials were prescribed without any scientific reasoning [204,205]. Dental surgeons and family practitioners frequently prescribed antibiotics for their patients (24%) as outpatient care [206]. Several studies reported that antibiotics are often overprescribed by dental surgeons for non-indicated diseases and irrationally in dental diseases which is the basis of antimicrobial resistance [207–209]. Moreover, dental surgeons very rarely documented infection through performing culture-sensitivity tests and the majority of antimicrobial prescribing was the result of “guess work” and described as imprudent use [111,210]. Multiple earlier research studies advocated that dental surgeons should prescribe antibiotics/antimicrobials only to control documented, local systemic infections, but not for mere inflammation [111,211,212]. Additionally, prophylactic use of antimicrobials in dental disease was suggested for only for suspected systemic infective conditions [213,214].

Antimicrobial resistance is a natural phenomenon that occurs as microbes evolve. However, human activities have accelerated the pace at which microorganisms develop and disseminate resistance. Incorrect and inappropriate use of antibiotics are contributing to the development of such resistance. It is necessary that all healthcare workers must recognize dental infections play a central role in preventing the emergence and spread of resistance. The inclusion of POHC as a component of primary health care may be the answer to promote rational dental care.

Author Contributions: Conceptualization: M.H., M.S.; Methodology: M.H., M.S.; Writing—original draft preparation: M.H., M.S. and S.Z.H.; Writing—review and editing: M.H., M.S., and S.Z.H.; Supervision: M.H.; Project administration: M.H.

Funding: This review manuscript receives no financial support.

Acknowledgments: The principal authors are much grateful to the Library of UPNM for their support to provide those manuscripts which are not open access.

Conflicts of Interest: Authors possess no conflict of interest.

References

1. American Chemical Society International Historic Chemical Landmarks. Discovery and Development of Penicillin. Available online: <http://www.acs.org/content/acs/en/education/whatischemistry/landmarks/flemingpenicillin.html> (accessed on 5 October 2018).
2. Rothman, L. This Is What Happened to the First American Treated with Penicillin. 2016. Available online: <http://time.com/4250235/penicillin-1942-history/> (accessed on 5 October 2018).
3. Gaynes, R. The Discovery of Penicillin—New Insights after More Than 75 Years of Clinical Use. *Emerg. Infect. Dis.* **2017**, *23*, 849–853. [CrossRef]

4. Burns, M. The Development of Penicillin in the Netherlands 1940–1950: The Pivotal Role of NV Nederlandsche Gist- en Spiritusfabriek, Delft. Ph.D. Thesis, Department of History, University of Sheffield, Sheffield, UK, 2005. Available online: <http://etheses.whiterose.ac.uk/14682/1/427207.pdf> (accessed on 5 October 2018).
5. Davies, J.; Davies, D. Origins and Evolution of Antibiotic Resistance. *Microbiol. Mol. Biol. Rev.* **2010**, *74*, 417–433. [[CrossRef](#)] [[PubMed](#)]
6. Davies, J.E. Origins, acquisition and dissemination of antibiotic resistance determinants. *Ciba Found Symp.* **1997**, *207*, 15–27. [[PubMed](#)]
7. Munita, J.M.; Arias, C.A. Mechanisms of Antibiotic Resistance. *Microbiol. Spectr.* **2016**, *4*. [[CrossRef](#)]
8. Davies, J. Microbes have the last word. A drastic re-evaluation of antimicrobial treatment is needed to overcome the threat of antibiotic-resistant bacteria. *EMBO Rep.* **2007**, *8*, 616–621. [[CrossRef](#)] [[PubMed](#)]
9. Abraham, E.P.; Chain, E. An Enzyme from Bacteria able to Destroy Penicillin. 1940. *Rev. Infect. Dis.* **1988**, *10*, 677–678. [[PubMed](#)]
10. D’Costa, V.M.; Griffiths, E.; Wright, G.D. Expanding the soil antibiotic resistome: Exploring environmental diversity. *Curr. Opin. Microbiol.* **2007**, *10*, 481–489. [[CrossRef](#)] [[PubMed](#)]
11. Liu, B.; Pop, M. ARDB—Antibiotic Resistance Genes Database. *Nucleic Acids Res.* **2009**, *37*, D443–D447. [[CrossRef](#)] [[PubMed](#)]
12. World Health Organization. *Antimicrobial Resistance: Global Report on Surveillance*; World Health Organization: Geneva, Switzerland, 2014. Available online: http://apps.who.int/iris/bitstream/handle/10665/112642/9789241564748_eng.pdf (accessed on 7 October 2018).
13. World Health Organization. *Antimicrobial Resistance*; World Health Organization: Geneva, Switzerland, 2018. Available online: <http://www.who.int/news-room/fact-sheets/detail/antimicrobial-resistance> (accessed on 7 October 2018).
14. Nazir, M.A. Prevalence of periodontal disease, its association with systemic diseases and prevention. *Int. J. Health Sci.* **2017**, *11*, 72–80.
15. Sanz, M.; D’Aiuto, F.; Deanfield, J.; Fernandez-Avilés, F. European workshop in periodontal health and cardiovascular disease—scientific evidence on the association between periodontal and cardiovascular diseases: A review of the literature. *Eur. Heart J. Suppl.* **2010**, *12*, B3–B12. [[CrossRef](#)]
16. Petersen, P.E.; Ogawa, H. The global burden of periodontal disease: Towards integration with chronic disease prevention and control. *Periodontol 2000* **2012**, *60*, 15–39. [[CrossRef](#)] [[PubMed](#)]
17. Shaju, J.P.; Zade, R.M.; Das, M. Prevalence of periodontitis in the Indian population: A literature review. *J. Indian Soc. Periodontol.* **2011**, *15*, 29–34. [[CrossRef](#)] [[PubMed](#)]
18. Peter, S. *Essentials of Preventive and Community Dentistry*; Arya MEDI Publishing House: New Delhi, India, 2007.
19. Petersen, P.E.; Bourgeois, D.; Ogawa, H.; Estupinan-Day, S.; Ndiaye, C. The Global Burden of Oral Diseases and Risks to Oral Health. *Bull. World Health Organ.* **2005**, *83*, 661–669. [[PubMed](#)]
20. Peterson, P.E. The World Oral Health Report 2003: Continuous Improvement of Oral Health in the 21st Century—The Approach of the WHO Global Oral Health Programme. *Community Dent. Oral Epidemiol.* **2003**, *31*, 3–23. [[CrossRef](#)]
21. Congiu, G.; Campus, G.; Lugliè, P.F. Early Childhood Caries (ECC) Prevalence and Background Factors: A Review. *Oral Health Prev. Dent.* **2014**, *12*, 71–76. [[PubMed](#)]
22. Zhang, S.; Chau, A.M.; Lo, E.C.; Chu, C.H. Dental Caries and Erosion Status of 12-Year-Old Children in Hong Kong. *BMC Public Health* **2014**, *14*, 7. [[CrossRef](#)] [[PubMed](#)]
23. Guaré, R.O.; Ciamponi, A.L.; BR Santos, M.T.; Gorjão, R.; Diniz, M.B. Caries Experience and Salivary Parameters among Overweight Children and Adolescents. *Dent. J.* **2013**, *1*, 31–40. [[CrossRef](#)]
24. Lagerweij, M.D.; van Loveren, C. Declining Caries Trends: Are We Satisfied? *Curr. Oral Health Rep.* **2015**, *2*, 212–217. [[CrossRef](#)] [[PubMed](#)]
25. Hong, L.; Ahmed, A.; McCunniff, M.; Overman, P.; Mathew, M. Obesity and Dental Caries in Children Aged 2–6 Years in the United States: National Health and Nutrition Examination Survey 1999–2002. *J. Public Health Dent.* **2008**, *68*, 227–233. [[CrossRef](#)] [[PubMed](#)]
26. Kundu, D.; Mehta, R.; Rozra, S. Periodontal status of a given population of West Bengal: An epidemiological study. *J. Indian Soc. Periodontol.* **2011**, *15*, 126–129. [[PubMed](#)]
27. Peeran, S.W.; Singh, A.J.A.R.; Alagamuthu, G.; Naveen Kumar, P.G. Periodontal status and its risk factors among young adults of the Sebha city (Libya). *Dent. Res. J.* **2013**, *10*, 533–538.

28. Bokhari, S.A.H.; Suhail, A.M.; Malik, A.R.; Imran, M.F. Periodontal disease status and associated risk factors in patients attending a Dental Teaching Hospital in Rawalpindi, Pakistan. *J. Indian Soc. Periodontol.* **2015**, *19*, 678–682. [[CrossRef](#)] [[PubMed](#)]
29. AlJehani, Y.A. Risk Factors of Periodontal Disease: Review of the Literature. *Int. J. Dent.* **2014**, *2014*, 182513. [[CrossRef](#)] [[PubMed](#)]
30. Tadjoeidin, F.M.; Fitri, A.H.; Kuswandani, S.O.; Sulijaya, B.; Soeroso, Y. The correlation between age and periodontal diseases. *J. Int. Dent. Med. Res.* **2017**, *10*, 327–332.
31. Frencken, J.E.; Sharma, P.; Stenhouse, L.; Green, D.; Laverty, D.; Dietrich, T. Global epidemiology of dental caries and severe periodontitis—A comprehensive review. *J. Clin. Periodontol.* **2017**, *4*, S94–S105. [[CrossRef](#)] [[PubMed](#)]
32. Schulze, A.; Busse, M. Gender Differences in Periodontal Status and Oral Hygiene of Non-Diabetic and Type 2 Diabetic Patients. *Open Dent. J.* **2016**, *10*, 287–297. [[CrossRef](#)] [[PubMed](#)]
33. Petersen, P.E.; Ogawa, H. Strengthening the prevention of periodontal disease: The WHO approach. *J. Periodontol.* **2005**, *76*, 2187–2193. [[CrossRef](#)] [[PubMed](#)]
34. Quadri, M.F.A.; Shubayr, M.A.; Hattan, A.H.; Wafi, S.A.; Jafer, A.H. Oral Hygiene Practices among Saudi Arabian Children and Its Relation to Their Dental Caries Status. *Int. J. Dent.* **2018**, *2018*, 3234970. [[CrossRef](#)] [[PubMed](#)]
35. Flynn, T.R.; Shanti, R.M.; Hayes, C. Severe odontogenic infections, part 2: Prospective outcomes study. *J. Oral Maxillofac. Surg.* **2006**, *64*, 1104–1113. [[CrossRef](#)] [[PubMed](#)]
36. Azodo, C.C.; Chukwumah, N.M.; Ezeja, E.B. Dentoalveolar abscess among children attending a dental clinic in Nigeria. *Odontostomatol. Trop.* **2012**, *35*, 41–46. [[PubMed](#)]
37. Sanadhya, S.; Aapaliya, P.; Jain, S.; Sharma, N.; Choudhary, G.; Dobaria, N. Assessment and comparison of clinical dental status and its impact on oral health-related quality of life among rural and urban adults of Udaipur, India: A cross-sectional study. *J. Basic Clin. Pharm.* **2015**, *6*, 50–58. [[PubMed](#)]
38. Schuller, A.A.; van Dommelen, P.; Poorterman, J.H. Trends in oral health in young people in the Netherlands over the past 20 years: A study in a changing context. *Community Dent. Oral Epidemiol.* **2014**, *42*, 178–184. [[CrossRef](#)] [[PubMed](#)]
39. Murray, J.J.; Vernazza, C.R.; Holmes, R.D. Forty years of national surveys: An overview of children’s dental health from 1973–2013. *Br. Dent. J.* **2015**, *219*, 281–285. [[CrossRef](#)] [[PubMed](#)]
40. Haugejorden, O.; Magne Birkeland, J. Ecological time-trend analysis of caries experience at 12 years of age and caries incidence from age 12 to 18 years: Norway 1985–2004. *Acta Odontol. Scand.* **2006**, *64*, 368–375. [[CrossRef](#)] [[PubMed](#)]
41. WHO Oral Health Country/Area Profile Programme. EURO Incl. DMFT for 12-Year-Olds. Available online: <http://www.whocollab.od.mah.se/euro.html> (accessed on 16 October 2018).
42. Broadbent, J.M.; Thomson, W.M. For debate: Problems with the DMF index pertinent to dental caries data analysis. *Community Dent. Oral Epidemiol.* **2005**, *33*, 400–409. [[CrossRef](#)] [[PubMed](#)]
43. Weusmann, J.; Mahmoodi, B.; Azaripour, A.; Kordsmeyer, K.; Walter, C.; Willershausen, B. Epidemiological investigation of caries prevalence in first-grade school children in Rhineland-Palatinate, Germany. *Head Face Med.* **2015**, *11*, 33. [[CrossRef](#)] [[PubMed](#)]
44. Asikainen, S.; Alaluusua, S. Bacteriology of dental infections. *Eur. Heart J.* **1993**, *14*, 43–50. [[PubMed](#)]
45. Aas, J.A.; Paster, B.J.; Stokes, L.N.; Olsen, I.; Dewhirst, F.E. Defining the Normal Bacterial Flora of the Oral Cavity. *J. Clin. Microbiol.* **2005**, *43*, 5721–5732. [[CrossRef](#)] [[PubMed](#)]
46. Samaranayake, L.; Matsubara, V.H. Normal Oral Flora and the Oral Ecosystem. *Dent. Clin. N. Am.* **2017**, *61*, 199–215. [[CrossRef](#)] [[PubMed](#)]
47. Zhang, Y.; Wang, X.; Li, H.; Ni, C.; Du, Z.; Yan, F. Human oral microbiota and its modulation for oral health. *Biomed. Pharmacother.* **2018**, *99*, 883–893. [[CrossRef](#)] [[PubMed](#)]
48. Kuriyama, T.; Karasawa, T.; Nakagawa, K.; Saiki, Y.; Yamamoto, E.; Nakamura, S. Bacteriologic features and antimicrobial susceptibility in isolates from orofacial odontogenic infections. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol. Endodontol.* **2000**, *90*, 600–608. [[CrossRef](#)] [[PubMed](#)]
49. Kuriyama, T.; Williams, D.W.; Yanagisawa, M.; Iwahara, K.; Shimizu, C.; Nakagawa, K.; Yamamoto, E.; Karasawa, T. Antimicrobial susceptibility of 800 anaerobic isolates from patients with dentoalveolar infection to 13 oral antibiotics. *Oral Microbiol. Immunol.* **2007**, *22*, 285–288. [[CrossRef](#)] [[PubMed](#)]

50. Rams, T.E.; Feik, D.; Listgarten, M.A.; Slots, J. Peptostreptococcus micros in human periodontitis. *Oral Microbiol. Immunol.* **1992**, *7*, 1–6. [[CrossRef](#)] [[PubMed](#)]
51. Rôças, I.N.; Siqueira, J.F., Jr.; Aboim, M.C.; Rosado, A.S. Denaturing gradient gel electrophoresis analysis of bacterial communities associated with failed endodontic treatment. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol. Endodontol.* **2004**, *98*, 741–749. [[CrossRef](#)]
52. Kayaoglu, G.; Ørstavik, D. Virulence factors of enterococcus faecalis: Relationship to endodontic disease. *Crit. Rev. Oral Biol. Med.* **2004**, *15*, 308–320. [[CrossRef](#)] [[PubMed](#)]
53. Rôças, I.N.; Siqueira, J.F., Jr.; Santos, K.R. Association of enterococcus faecalis with different forms of periradicular diseases. *J. Endodontol.* **2004**, *30*, 315–320. [[CrossRef](#)] [[PubMed](#)]
54. Schleifer, K.H.; Kilpper-balz, R. Transfer of streptococcus faecalis and streptococcus faecium to the genus enterococcus nom. rev. as enterococcus faecalis comb. nov. and enterococcus faecium comb. nov. *Int. J. Syst. Bacteriol.* **1984**, *34*, 31–34. [[CrossRef](#)]
55. Ryan, K.; Ray, C.G. *Sherris Medical Microbiology: An Introduction to Infectious Diseases*; McGraw-Hill Medical: New York, NY, USA, 2004; pp. 294–295.
56. Siqueira, J.F., Jr.; Rôças, I.N. Uncultivated phylotypes and newly named species associated with primary and persistent endodontic infections. *J. Clin. Microbiol.* **2005**, *43*, 3314–3319. [[CrossRef](#)] [[PubMed](#)]
57. Schäfer, E.; Bossmann, K. Antimicrobial efficacy of chloroxylenol and chlorhexidine in the treatment of infected root canals. *Am. J. Dent.* **2001**, *14*, 233–237. [[PubMed](#)]
58. Moynihan, P.; Petersen, P.E. Diet, nutrition and the prevention of dental diseases. *Public Health Nutr.* **2004**, *7*, 201–226. [[CrossRef](#)] [[PubMed](#)]
59. Burgess, J. Diet and Oral Health. *Drugs and Diseases. Dentistry.* Medscape. 2015. Available online: <https://emedicine.medscape.com/article/2066208-overview> (accessed on 24 October 2018).
60. Scardina, G.A.; Messina, P. Good oral health and diet. *J. Biomed. Biotechnol.* **2012**, *2012*, 720692. [[CrossRef](#)] [[PubMed](#)]
61. Xu, H.; Tian, J.; Hao, W.; Zhang, Q.; Zhou, Q.; Shi, W.; Qin, M.; He, X.; Chen, F. Oral Microbiome Shifts from Caries-Free to Caries-Affected Status in 3-Year-Old Chinese Children: A Longitudinal Study. *Front. Microbiol.* **2018**, *9*, 2009. [[CrossRef](#)] [[PubMed](#)]
62. Bang, G.; Kristoffersen, T. Dental caries and diet in an Alaskan Eskimo population. *Scand. J. Dent. Res.* **1972**, *80*, 440–444. [[CrossRef](#)] [[PubMed](#)]
63. Cheng, R.; Yang, H.; Shao, M.Y.; Hu, T.; Zhou, X.D. Dental erosion and severe tooth decay related to soft drinks: A case report and literature review. *J. Zhejiang Univ. Sci. B* **2009**, *10*, 395–399. [[CrossRef](#)] [[PubMed](#)]
64. Kannan, A.; Ahmed, M.A.A.; Duraisamy, P.; Manipal, S.; Adusumillil, P. Dental hard tissue erosion rates and soft drinks—A gender-based analysis in Chennai city, India. *Saudi J. Dent. Res.* **2014**, *5*, 21–27. [[CrossRef](#)]
65. Wang, Y.L.; Chang, C.C.; Chi, C.W.; Chang, H.H.; Chiang, Y.C.; Chuang, Y.C.; Chang, H.H.; Huang, G.F.; Liao, Y.S.; Lin, C.P. Erosive potential of soft drinks on human enamel: An in vitro study. *J. Formos. Med. Assoc.* **2014**, *113*, 850–856. [[CrossRef](#)] [[PubMed](#)]
66. Sohn, W.; Burt, B.A.; Sowers, M.R. Carbonated soft drinks and dental caries in the primary dentition. *J. Dent. Res.* **2006**, *85*, 262–266. [[CrossRef](#)] [[PubMed](#)]
67. Skinner, J.; Byun, R.; Blinkhorn, A.; Johnson, G. Sugary drink consumption and dental caries in New South Wales teenagers. *Aust. Dent. J.* **2015**, *60*, 169–175. [[CrossRef](#)] [[PubMed](#)]
68. Marshall, T.A.; Levy, S.M.; Broffitt, B.; Warren, J.J.; Eichenberger-Gilmore, J.M.; Burns, T.L.; Stumbo, P.J. Dental caries and beverage consumption in young children. *Pediatrics* **2003**, *112*, E184–E191. [[CrossRef](#)] [[PubMed](#)]
69. Hugar, S.M.; Dhariwal, N.S.; Majeed, A.; Badakar, C.; Gokhale, N.; Mistry, L. Assessment of Vitamin B12 and Its Correlation with Dental Caries and Gingival Diseases in 10- to 14-year-old Children: A Cross-sectional Study. *Int. J. Clin. Pediatr. Dent.* **2017**, *10*, 142–146. [[CrossRef](#)] [[PubMed](#)]
70. Shenkin, J.D.; Heller, K.E.; Warren, J.J.; Marshall, T.A. Soft drink consumption and caries risk in children and adolescents. *Gen. Dent.* **2003**, *51*, 30–36. [[PubMed](#)]
71. Campmans-Kuijpers, M.J.; Singh-Povel, C.; Steijns, J.; Beulens, J.W. The association of dairy intake of children and adolescents with different food and nutrient intakes in the Netherlands. *BMC Pediatr.* **2016**, *16*, 2. [[CrossRef](#)] [[PubMed](#)]

72. Weinberg, L.G.; Berner, L.A.; Groves, J.E. Nutrient contributions of dairy foods in the United States, Continuing Survey of Food Intakes by Individuals, 1994–1996, 1998. *J. Am. Diet Assoc.* **2004**, *104*, 895–902. [[CrossRef](#)] [[PubMed](#)]
73. Ranganathan, R.; Nicklas, T.A.; Yang, S.J.; Berenson, G.S. The nutritional impact of dairy product consumption on dietary intakes of adults (1995–1996): The Bogalusa Heart Study. *J. Am. Diet. Assoc.* **2005**, *105*, 1391–1400. [[CrossRef](#)] [[PubMed](#)]
74. Muehlhoff, E.; Bennett, A.; McMahon, D. Milk, and Dairy Products in Human Nutrition. The Food and Agriculture Organization (FAO) of the United Nations. 2013. Available online: <http://www.fao.org/docrep/018/i3396e/i3396e.pdf> (accessed on 25 October 2018).
75. Gupta, P.; Gupta, N.; Pawar, A.P.; Birajdar, S.S.; Natt, A.S.; Singh, H.P. Role of sugar and sugar substitutes in dental caries: A review. *ISRN Dent.* **2013**, *2013*, 519421. [[CrossRef](#)] [[PubMed](#)]
76. Olsson, B. Dental health situation in privileged children in Addis Ababa, Ethiopia. *Community Dent. Oral Epidemiol.* **1979**, *7*, 37–41. [[CrossRef](#)] [[PubMed](#)]
77. Scheinin, A.; Mäkinen, K.K. Turku sugar studies. An overview. *Acta Odontol. Scand.* **1976**, *34*, 405–408. [[CrossRef](#)] [[PubMed](#)]
78. Sheiham, A.; James, W.P. A reappraisal of the quantitative relationship between sugar intake and dental caries: The need for new criteria for developing goals for sugar intake. *BMC Public Health.* **2014**, *14*, 863. [[CrossRef](#)] [[PubMed](#)]
79. Bradshaw, D.J.; Lynch, R.J. Diet and the microbial etiology of dental caries: New paradigms. *Int. Dent. J.* **2013**, *63*, 64–72. [[CrossRef](#)] [[PubMed](#)]
80. Touger-Decker, R.; van Loveren, C. Sugars and dental caries. *Am. J. Clin. Nutr.* **2003**, *78*, 881s–892s. [[CrossRef](#)] [[PubMed](#)]
81. Saini, R.; Saini, S.; Sharma, S. Biofilm: A dental microbial infection. *J. Nat. Sci. Biol. Med.* **2011**, *2*, 71–75. [[CrossRef](#)] [[PubMed](#)]
82. Chandki, R.; Banthia, P.; Banthia, R. Biofilms: A microbial home. *J. Indian Soc. Periodontol.* **2011**, *15*, 111–114. [[PubMed](#)]
83. Øilo, M.; Bakken, V. Biofilm and Dental Biomaterials. *Materials* **2015**, *8*, 2887–2900. [[CrossRef](#)]
84. van Leeuwenhoek, A. An abstract of a letter from antonie van leeuwenhoek, Sep. 12, 1683. about animals in the scurf of the teeth. *Philos. Trans. R. Soc. Lond.* **1684**, *14*, 568–574.
85. Souto, R.; de Andrade, A.F.B.; Uzeda, M.; Colombo, A.P.V. Prevalence of “non-oral” pathogenic bacteria in subgingival biofilm of subjects with chronic periodontitis. *Braz. J. Microbiol.* **2006**, *37*, 208–215. [[CrossRef](#)]
86. Li, X.; Kolltveit, K.M.; Tronstad, L.; Olsen, I. Systemic diseases caused by oral infection. *Clin. Microbiol. Rev.* **2000**, *13*, 547–558. [[CrossRef](#)] [[PubMed](#)]
87. Marsh, P.D. Are dental diseases examples of ecological catastrophes? *Microbiology* **2003**, *149*, 279–294. [[CrossRef](#)] [[PubMed](#)]
88. de Almeida Pdel, V.; Grégio, A.M.; Machado, M.A.; de Lima, A.A.; Azevedo, L.R. Saliva composition and functions: A comprehensive review. *J. Contemp. Dent. Pract.* **2008**, *9*, 72–80. [[PubMed](#)]
89. Jenkinson, H.F.; Lappin-Scott, H.M. Biofilms adhere to stay. *Trends Microbiol.* **2001**, *9*, 9–10. [[CrossRef](#)]
90. Avila, M.; Ojcius, D.M.; Yilmaz, O. The oral microbiota: Living with a permanent guest. *DNA Cell Biol.* **2009**, *28*, 405–411. [[CrossRef](#)] [[PubMed](#)]
91. Sbordone, L.; Bortolaia, C. Oral microbial biofilms and plaque-related diseases: Microbial communities and their role in the shift from oral health to disease. *Clin. Oral Investig.* **2003**, *7*, 181–188. [[CrossRef](#)] [[PubMed](#)]
92. Marsh, P.D. Dental plaque as a biofilm and a microbial community—Implications for health and disease. *BMC Oral Health* **2006**, *6*, S14. [[CrossRef](#)] [[PubMed](#)]
93. Theilade, E. The non-specific theory in microbial etiology of inflammatory periodontal diseases. *J. Clin. Periodontol.* **1986**, *13*, 905–911. [[CrossRef](#)] [[PubMed](#)]
94. Loesche, W.J. Role of Streptococcus mutans in human dental decay. *Microbiol. Rev.* **1986**, *50*, 353–380. [[PubMed](#)]
95. Marsh, P.D. Microbial ecology of dental plaque and its significance in health and disease. *Adv. Dent. Res.* **1994**, *8*, 263–271. [[CrossRef](#)] [[PubMed](#)]
96. Yang, F.; Zeng, X.; Ning, K.; Liu, K.L.; Lo, C.C.; Wang, W.; Chen, J.; Wang, D.; Huang, R.; Chang, X.; et al. Saliva microbiomes distinguish caries-active from healthy human populations. *ISME J.* **2012**, *6*, 1–10. [[CrossRef](#)] [[PubMed](#)]

97. Rosier, B.T.; De Jager, M.; Zaura, E.; Krom, B.P. Historical and contemporary hypotheses on the development of oral diseases: Are we there yet? *Front. Cell Infect. Microbiol.* **2014**, *4*, 92. [[CrossRef](#)] [[PubMed](#)]
98. Fleming, A. The story of penicillin. *Bull. Georgetown Univ. Med. Cent.* **1955**, *8*, 128–132. [[PubMed](#)]
99. Adedeji, W.A. The Treasure Called Antibiotics. *Ann. Ib. Postgrad Med.* **2016**, *14*, 56–57. [[PubMed](#)]
100. Costelloe, C.; Metcalfe, C.; Lovering, A.; Mant, D.; Hay, A.D. Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: Systematic review and meta-analysis. *BMJ* **2010**, *340*, c2096. [[CrossRef](#)] [[PubMed](#)]
101. Bryce, A.; Hay, A.D.; Lane, I.F.; Thornton, H.V.; Wootton, M.; Costelloe, C. Global prevalence of antibiotic resistance in pediatric urinary tract infections caused by *Escherichia coli* and association with routine use of antibiotics in primary care: Systematic review and meta-analysis. *BMJ* **2016**, *352*, i939. [[CrossRef](#)] [[PubMed](#)]
102. Shehab, N.; Patel, P.R.; Srinivasan, A.; Budnitz, D.S. Emergency department visits for antibiotic-associated adverse events. *Clin. Infect. Dis.* **2008**, *47*, 735–743. [[CrossRef](#)] [[PubMed](#)]
103. Cope, A.L.; Chestnutt, I.G. Inappropriate Prescribing of Antibiotics in Primary Dental Care: Reasons and Resolutions. *Prim. Dent. J.* **2014**, *3*, 33–37. [[CrossRef](#)] [[PubMed](#)]
104. Maslamani, M.; Sedeqi, F. Antibiotic and Analgesic Prescription Patterns among Dentists or Management of Dental Pain and Infection during Endodontic Treatment. *Med. Princ. Pract.* **2017**, *27*, 66–72. [[CrossRef](#)] [[PubMed](#)]
105. Teoh, L.; Stewart, K.; Marino, R.J.; McCullough, M.J. Part 1. Current prescribing trends of antibiotics by dentists in Australia from 2013 to 2016. *Aust. Dent. J.* **2018**, *63*, 329–337. [[CrossRef](#)] [[PubMed](#)]
106. Marra, F.; George, D.; Chong, M.; Sutherland, S.; Patrick, D.M. Antibiotic prescribing by dentists has increased: Why? *J. Am. Dent. Assoc.* **2016**, *147*, 320–327. [[CrossRef](#)] [[PubMed](#)]
107. Roda, R.P.; Bagán, J.V.; Bielsa, J.M.S.; Pastor, E.C. Antibiotic use in dental practice. A review. *Med. Oral Patol. Oral Cir. Bucal* **2007**, *12*, E186–92.
108. Peedikayil, F.C. Antibiotics: Use and misuse in pediatric dentistry. *J. Indian Soc. Pedod. Prev. Dent.* **2011**, *29*, 282–287. [[CrossRef](#)] [[PubMed](#)]
109. Berney, P.; Francioli, P. Successful prophylaxis of experimental streptococcal endocarditis with single-dose amoxicillin administered after bacterial challenge. *J. Infect. Dis.* **1990**, *161*, 281–285. [[CrossRef](#)] [[PubMed](#)]
110. Dajani, A.S.; Taubert, K.A.; Wilson, W.; Bolger, A.F.; Bayer, A.; Ferrieri, P.; Gewitz, M.H.; Shulman, S.T.; Nouri, S.; Newburger, J.W.; et al. Prevention of bacterial endocarditis: Recommendations by the American Heart Association. *J. Am. Dent. Assoc.* **1997**, *128*, 1142–1151. [[CrossRef](#)] [[PubMed](#)]
111. Dar-Odeh, N.S.; Abu-Hammad, O.A.; Al-Omiri, M.K.; Khraisat, A.S.; Shehabi, A.A. Antibiotic prescribing practices by dentists: A review. *Ther. Clin. Risk Manag.* **2010**, *6*, 301–306. [[CrossRef](#)] [[PubMed](#)]
112. Oberoi, S.S.; Dhingra, C.; Sharma, G.; Sardana, D. Antibiotics in dental practice: How justified are we. *Int. Dent. J.* **2015**, *65*, 4–10. [[CrossRef](#)] [[PubMed](#)]
113. Roberts, R.M.; Bartoces, M.; Thompson, S.E.; Hicks, L.A. Antibiotic prescribing by general dentists in the United States, 2013. *J. Am. Dent. Assoc.* **2017**, *148*, 172–178. [[CrossRef](#)] [[PubMed](#)]
114. Lee, C.R.; Cho, I.H.; Jeong, B.C.; Lee, S.H. Strategies to minimize antibiotic resistance. *Int. J. Environ. Res. Public Health* **2013**, *10*, 4274–4305. [[CrossRef](#)] [[PubMed](#)]
115. Lee, C.R.; Lee, J.H.; Kang, L.W.; Jeong, B.C.; Lee, S.H. Educational effectiveness, target, and content for prudent antibiotic use. *Biomed. Res. Int.* **2015**, *2015*, 214021. [[CrossRef](#)] [[PubMed](#)]
116. Chunduri, N.S.; Madasu, K.; Goteki, V.R.; Karpe, T.; Reddy, H. Evaluation of bacterial spectrum of orofacial infections and their antibiotic susceptibility. *Ann. Maxillofac. Surg.* **2012**, *2*, 46–50. [[CrossRef](#)] [[PubMed](#)]
117. Robertson, D.; Smith, A.J. The microbiology of the acute dental abscess. *J. Med. Microbiol.* **2009**, *58*, 155–162. [[CrossRef](#)] [[PubMed](#)]
118. Teng, L.J.; Hsueh, P.R.; Chen, Y.C.; Ho, S.W.; Luh, K.T. Antimicrobial susceptibility of viridans group streptococci in Taiwan with an emphasis on the high rates of resistance to penicillin and macrolides in *Streptococcus oralis*. *J. Antimicrob. Chemother.* **1998**, *41*, 621–627. [[CrossRef](#)] [[PubMed](#)]
119. Kuriyama, T.; Karasawa, T.; Nakagawa, K.; Yamamoto, E.; Nakamura, S. Incidence of beta-lactamase production and antimicrobial susceptibility of anaerobic gram-negative rods isolated from pus specimens of orofacial odontogenic infections. *Oral Microbiol. Immunol.* **2001**, *16*, 10–15. [[CrossRef](#)] [[PubMed](#)]
120. Kuriyama, T.; Karasawa, T.; Nakagawa, K.; Yamamoto, E.; Nakamura, S. Bacteriology and antimicrobial susceptibility of gram-positive cocci isolated from pus specimens of orofacial odontogenic infections. *Oral Microbiol. Immunol.* **2002**, *17*, 132–135. [[CrossRef](#)] [[PubMed](#)]

121. Smith, A.J.; Jackson, M.S. Susceptibility of viridans group streptococci isolated from dento-alveolar infections to eight antimicrobial agents. *J. Antimicrob. Chemother.* **2003**, *52*, 1045–1046. [[CrossRef](#)] [[PubMed](#)]
122. Smith, A.; Jackson, M.S.; Kennedy, H. Antimicrobial susceptibility of viridans group streptococcal blood isolates to eight antimicrobial agents. *Scand. J. Infect. Dis.* **2004**, *36*, 259–263. [[CrossRef](#)] [[PubMed](#)]
123. Dwivedi, D.; Kushwah, T.; Kushwah, M.; Singh, V. Antibiotic susceptibility pattern against pathogenic bacteria causing Dental Caries. *South Asian J. Exp. Biol.* **2011**, *1*, 31–35.
124. Yadav, K.; Prakash, S.; Yadav, N.P.; Sah, R.S. Multi-Drug Resistance of Bacterial Isolates among Dental Caries Patients. *Janaki Med. Coll. J. Med. Sci.* **2015**, *3*, 37–44. [[CrossRef](#)]
125. Yadav, K.; Prakash, S. Antibiogram profiles against polymicrobial pathogens among dental caries patients at Janaki Medical College teaching hospital, Nepal. *Int. J. Appl. Dent. Sci.* **2015**, *1*, 156–162.
126. Gaetti-Jardim, E.C.; Marqueti, A.C.; Faverani, L.P.; Gaetti-Jardim Júnior, E. Antimicrobial resistance of aerobes and facultative anaerobes isolated from the oral cavity. *J. Appl. Oral Sci.* **2010**, *18*, 551–559. [[CrossRef](#)] [[PubMed](#)]
127. Loyola-Rodriguez, J.P.; Garcia-Cortes, J.O.; Martinez-Martinez, R.E.; Patiño-Marin, N.; Martínez-Castañón, G.A.; Zavala-Alonso, N.V.; Amano, A. Molecular identification and antibiotic resistant bacteria isolated from primary dentition infections. *Aust. Dent. J.* **2014**, *59*, 497–503. [[CrossRef](#)] [[PubMed](#)]
128. Sanz, M.; Beighton, D.; Curtis, M.A.; Cury, J.A.; Dige, I.; Dommisch, H.; Ellwood, R.; Giacaman, R.A.; Herrera, D.; Herzberg, M.C.; et al. Role of microbial biofilms in the maintenance of oral health and in the development of dental caries and periodontal diseases. Consensus report of group 1 of the Joint EFP/ORCA workshop on the boundaries between caries and periodontal disease. *J. Clin. Periodontol.* **2017**, *44*, S5–S11. [[CrossRef](#)] [[PubMed](#)]
129. Mayo Clinic Staff. Oral health: A Window to Your Overall Health. Healthy Lifestyle Adult Health. Mayo Clinic. Available online: <https://www.mayoclinic.org/healthy-lifestyle/adult-health/in-depth/dental/art-20047475> (accessed on 13 January 2019).
130. Dar-Odeh, N.; Fadel, H.T.; Abu-Hammad, S.; Abdeljawad, R.; Abu-Hammad, O.A. Antibiotic Prescribing for Oro-Facial Infections in the Pediatric Outpatient: A Review. *Antibiotics* **2018**, *7*, 38. [[CrossRef](#)] [[PubMed](#)]
131. World Health Organization. Oral Health. The 11th IDAR World Congress on Preventive Dentistry. Available online: http://www.who.int/oral_health/events/congress-preventive-dentistry-oct2017-outcomes/en/ (accessed on 9 November 2018).
132. Meurman, J.H.; Hämäläinen, P. Oral health and morbidity—Implications of oral infections on the elderly. *Gerodontology* **2006**, *23*, 3–16. [[CrossRef](#)] [[PubMed](#)]
133. Meurman, J.H. Dental infections and general health. *Quintessence Int.* **1997**, *28*, 807–811. [[PubMed](#)]
134. Slots, J. Update on general health risk of periodontal disease. *Int. Dent. J.* **2003**, *53*, 200–207. [[CrossRef](#)] [[PubMed](#)]
135. Meurman, J.H.; Qvarnström, M.A. Chronic oral infections—a threat to health? *Duodecim* **1995**, *111*, 1348–1354. [[PubMed](#)]
136. Kraut, R.A.; Hicks, J.L. Bacterial endocarditis of dental origin: Report of case. *J. Oral Surg.* **1976**, *34*, 1031–1034. [[PubMed](#)]
137. Martos, R.; Márton, I. Correlations between dental-oral infections and cardiovascular disease. *Fogorvosi Szemle* **2008**, *101*, 101–105. [[PubMed](#)]
138. Adolph, M.; Darnaud, C.; Thomas, F.; Pannier, B.; Danchin, N.; Batty, G.D.; Boucharda, P. Oral health in relation to all-cause mortality: The IPC cohort study. *Sci. Rep.* **2017**, *7*, 44604. [[CrossRef](#)] [[PubMed](#)]
139. Kim, J.K.; Baker, L.A.; Davarian, S.; Crimmins, E. Oral health problems and mortality. *J. Dent. Sci.* **2013**, *8*. [[CrossRef](#)] [[PubMed](#)]
140. Hayasaka, K.; Tomata, Y.; Aida, J.; Watanabe, T.; Kakizaki, M.; Tsuji, I. Tooth loss and mortality in elderly Japanese adults: Effect of oral care. *J. Am. Geriatr. Soc.* **2013**, *61*, 815–820. [[CrossRef](#)] [[PubMed](#)]
141. Chambrone, L.; Chambrone, D.; Lima, L.A.; Chambrone, L.A. Predictors of tooth loss during long-term periodontal maintenance: A systematic review of observational studies. *J. Clin. Periodontol.* **2010**, *37*, 675–684. [[CrossRef](#)] [[PubMed](#)]
142. Janket, S.J.; Baird, A.E.; Chuang, S.K.; Jones, J.A. Meta-analysis of periodontal disease and risk of coronary heart disease and stroke. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol. Endodontol.* **2003**, *95*, 559–569. [[CrossRef](#)] [[PubMed](#)]

143. Azarpazhooh, A.; Leake, J.L. Systematic review of the association between respiratory diseases and oral health. *J. Periodontol.* **2006**, *77*, 1465–1482. [[CrossRef](#)] [[PubMed](#)]
144. Wolff, J.L.; Boulton, C.; Boyd, C.; Anderson, G. Newly reported chronic conditions and onset of functional dependency. *J. Am. Geriatr. Soc.* **2005**, *53*, 851–855. [[CrossRef](#)] [[PubMed](#)]
145. Aida, J.; Kondo, K.; Hirai, H.; Nakade, M.; Yamamoto, T.; Hanibuchi, T.; Osaka, K.; Sheiham, A.; Tsakos, G.; Watt, R.G. Association between dental status and incident disability in an older Japanese population. *J. Am. Geriatr. Soc.* **2012**, *60*, 338–343. [[CrossRef](#)] [[PubMed](#)]
146. Tsakos, G.; Watt, R.G.; Rouxel, P.L.; de Oliveira, C.; Demakakos, P. Tooth loss associated with physical and cognitive decline in older adults. *J. Am. Geriatr. Soc.* **2015**, *63*, 91–99. [[CrossRef](#)] [[PubMed](#)]
147. Chen, P.C.; Tung, Y.C.; Wu, P.W.; Wu, L.S.; Lin, Y.S.; Chang, C.J.; Kung, S.; Chu, P.H. Dental Procedures and the Risk of Infective Endocarditis. *Medicine (Baltimore)* **2015**, *94*, e1826. [[CrossRef](#)] [[PubMed](#)]
148. Esdaile, J.M.; Abrahamowicz, M.; Grodzicky, T.; Li, Y.; Panaritis, C.; du Berger, R.; Côte, R.; Grover, S.A.; Fortin, P.R.; Clarke, A.E.; et al. Traditional Framingham risk factors fail to fully account for accelerated atherosclerosis in systemic lupus erythematosus. *Arthritis Rheum.* **2001**, *44*, 2331–2337. [[CrossRef](#)]
149. Otto, C.M. Heartbeat: Chronic inflammatory disorders and cardiovascular disease. *Heart* **2016**, *102*, 1935–1936. [[CrossRef](#)] [[PubMed](#)]
150. Mason, J.C.; Libby, P. Cardiovascular disease in patients with chronic inflammation: Mechanisms underlying premature cardiovascular events in rheumatologic conditions. *Eur. Heart J.* **2014**, *36*, 482–489. [[CrossRef](#)] [[PubMed](#)]
151. Dunne, M. Infectious Agents and Cardiovascular Disease. In *The Infectious Etiology of Chronic Diseases: Defining the Relationship, Enhancing the Research, and Mitigating the Effects: Workshop Summary*; Institute of Medicine (US) Forum on Microbial Threats, Knobler, S.L., O'Connor, S., Lemon, S.M., Eds.; National Academies Press: Washington, DC, USA, 2004. Available online: <https://www.ncbi.nlm.nih.gov/books/NBK83710/> (accessed on 11 November 2018).
152. Syrjänen, J. Vascular diseases and oral infections. *J. Clin. Periodontol.* **1990**, *17*, 497–500. [[CrossRef](#)] [[PubMed](#)]
153. Valtonen, V.V. Infection as a risk factor for infarction and atherosclerosis. *Ann. Med.* **1991**, *23*, 539–543. [[CrossRef](#)] [[PubMed](#)]
154. Syrjänen, J.; Peltola, J.; Valtonen, V.; Iivanainen, M.; Kaste, M.; Huttunen, J.K. Dental infections in association with cerebral infarction in young and middle-aged men. *J. Intern. Med.* **1989**, *225*, 179–184. [[CrossRef](#)] [[PubMed](#)]
155. Syrjänen, J. Infection as a risk factor for cerebral infarction. *Eur. Heart J.* **1993**, *14*, 17–19. [[PubMed](#)]
156. Chaudhry, S.; Jaiswal, R.; Sachdeva, S. Dental considerations in cardiovascular patients: A practical perspective. *Indian Heart J.* **2016**, *68*, 572–575. [[CrossRef](#)] [[PubMed](#)]
157. Rautemaa, R.; Lauhio, A.; Cullinan, M.P.; Seymour, G.J. Oral infections and systemic disease—An emerging problem in medicine. *Clin. Microbiol. Infect.* **2007**, *13*, 1041–1047. [[CrossRef](#)] [[PubMed](#)]
158. Saini, R.; Saini, S.; Saini, S.R. Periodontal diseases: A risk factor to cardiovascular disease. *Ann. Card. Anaesth.* **2010**, *13*, 159–161. [[CrossRef](#)] [[PubMed](#)]
159. Gani, D.K.; Lakshmi, D.; Krishnan, R.; Emmadi, P. Evaluation of C-reactive protein and interleukin-6 in the peripheral blood of patients with chronic periodontitis. *J. Indian Soc. Periodontol.* **2009**, *13*, 69–74. [[CrossRef](#)] [[PubMed](#)]
160. Bansal, T.; Dhruvakumar, D.; Pandey, A. Comparative evaluation of C-reactive protein in peripheral blood of patients with healthy gingiva, gingivitis, and chronic periodontitis: A clinical and particle-enhanced turbidimetric immuno-analysis. *J. Indian Soc. Periodontol.* **2014**, *18*, 739–743. [[CrossRef](#)] [[PubMed](#)]
161. Mayo Clinic Staff. C-Reactive Protein Test. Overview. 2017. Available online: <https://www.mayoclinic.org/tests-procedures/c-reactive-protein-test/about/pac-20385228> (accessed on 11 November 2018).
162. Belfki, H.; Ben Ali, S.; Bougatef, S.; Ben Ahmed, D.; Haddad, N.; Jmal, A.; Abdennebi, M.; Ben Romdhane, H. Association between C-reactive protein and type 2 diabetes in a Tunisian population. *Inflammation* **2012**, *35*, 684–689. [[CrossRef](#)] [[PubMed](#)]
163. Dongway, A.C.; Faggad, A.S.; Zaki, H.Y.; Abdalla, B.E. C-reactive protein is associated with low-density lipoprotein cholesterol and obesity in type 2 diabetic Sudanese. *Diabetes Metab. Syndr. Obes.* **2015**, *8*, 427–435. [[PubMed](#)]
164. Hu, F.B.; Meigs, J.B.; Li, T.Y.; Rifai, N.; Manson, J.E. Inflammatory markers and risk of developing type 2 diabetes in women. *Diabetes* **2004**, *53*, 693–700. [[CrossRef](#)] [[PubMed](#)]

165. Doi, Y.; Kiyohara, Y.; Kubo, M.; Ninomiya, T.; Wakugawa, Y.; Yonemoto, K.; Iwase, M.; Iida, M. Elevated C-reactive protein is a predictor of the development of diabetes in a general Japanese population: The Hisayama Study. *Diabetes Care* **2005**, *28*, 2497–2500. [[CrossRef](#)] [[PubMed](#)]
166. Phosat, C.; Panprathip, P.; Chumpathat, N.; Prangthip, P.; Chantratita, N.; Soonthornworasiri, N.; Puduang, S.; Kwanbunjan, K. Elevated C-reactive protein, interleukin 6, tumor necrosis factor alpha and glycemic load associated with type 2 diabetes mellitus in rural Thais: A cross-sectional study. *BMC Endocr. Disord.* **2017**, *17*, 44. [[CrossRef](#)] [[PubMed](#)]
167. Liu, C.; Feng, X.; Li, Q.; Wang, Y.; Li, Q.; Hua, M. Adiponectin, TNF- α and inflammatory cytokines and risk of type 2 diabetes: A systematic review and meta-analysis. *Cytokine* **2016**, *86*, 100–109. [[CrossRef](#)] [[PubMed](#)]
168. Giovannini, S.; Onder, G.; Liperoti, R.; Russo, A.; Carter, C.; Capoluongo, E.; Pahor, M.; Bernabei, R.; Landi, F. Interleukin-6, C-reactive protein, and tumor necrosis factor-alpha as predictors of mortality in frail, community-living elderly individuals. *J. Am. Geriatr. Soc.* **2011**, *59*, 1679–1685. [[CrossRef](#)] [[PubMed](#)]
169. Bansal, T.; Pandey, A.; Deepa, D.; Asthana, A.K. C-Reactive Protein (CRP) and its Association with Periodontal Disease: A Brief Review. *J. Clin. Diagn. Res.* **2014**, *8*, ZE21–ZE24. [[PubMed](#)]
170. Noh, M.K.; Jung, M.; Kim, S.H.; Lee, S.R.; Park, K.H.; Kim, D.H.; Kim, H.H.; Park, Y.G. Assessment of IL-6, IL-8 and TNF- α levels in the gingival tissue of patients with periodontitis. *Exp. Ther. Med.* **2013**, *6*, 847–851. [[CrossRef](#)] [[PubMed](#)]
171. Haerian-Ardakani, A.; Eslami, Z.; Rashidi-Meibodi, F.; Haerian, A.; Dallalnejad, P.; Shekari, M.; Taghavi, A.M.; Akbari, S. Relationship between maternal periodontal disease and low birth weight babies. *Iran J. Reprod Med.* **2013**, *11*, 625–630. [[PubMed](#)]
172. Chambrone, L.; Guglielmetti, M.R.; Pannuti, C.M.; Chambrone, L.A. Evidence grade associating periodontitis to preterm birth and/or low birth weight: I. A systematic review of prospective cohort studies. *J. Clin. Periodontol.* **2011**, *38*, 795–808. [[CrossRef](#)] [[PubMed](#)]
173. Corbella, S.; Taschieri, S.; Francetti, L.; De Siena, F.; Del Fabbro, M. Periodontal disease as a risk factor for adverse pregnancy outcomes: A systematic review and meta-analysis of case-control studies. *Odontology* **2012**, *100*, 232–240. [[CrossRef](#)] [[PubMed](#)]
174. Teshome, A.; Yitayeh, A. Relationship between periodontal disease and preterm low birth weight: Systematic review. *Pan Afr. Med. J.* **2016**, *24*, 215. [[CrossRef](#)] [[PubMed](#)]
175. Mannem, S.; Chava, V.K. The relationship between maternal periodontitis and preterm low birth weight: A case-control study. *Contemp. Clin. Dent.* **2011**, *2*, 88–93. [[CrossRef](#)] [[PubMed](#)]
176. Benjamin, R.M. Oral health: The silent epidemic. *Public Health Rep.* **2010**, *125*, 158–159. [[CrossRef](#)] [[PubMed](#)]
177. Patrick, D.L.; Lee, R.S.; Nucci, M.; Grembowski, D.; Jolles, C.Z.; Milgrom, P. Reducing oral health disparities: A focus on social and cultural determinants. *BMC Oral Health* **2006**, *6*, S4. [[CrossRef](#)] [[PubMed](#)]
178. Kenney, G.M.; McFeeters, J.R.; Justin, Y.Y. Preventive dental care and unmet dental needs among low-income children. *Am. J. Public Health* **2005**, *95*, 1360–1366. [[CrossRef](#)] [[PubMed](#)]
179. Davis, E.E.; Deinard, A.S.; Maïga, E.W. Doctor, my tooth hurts: The costs of incomplete dental care in the emergency room. *J. Public Health Dent.* **2010**, *70*, 205–210. [[CrossRef](#)] [[PubMed](#)]
180. Donoff, B.; McDonough, J.E.; Riedy, C.A. Integrating oral and general health care. *N. Engl. J. Med.* **2014**, *371*, 2247–2249. [[CrossRef](#)] [[PubMed](#)]
181. Manski, R.J.; Brown, E. *Dental Use, Expenses, Private Dental Coverage, and Changes, 1996 and 2004*; MEPS Chartbook No.17. Medical Expenditure Panel Survey; Agency for Healthcare Research and Quality U.S. Department of Health & Human Services: Rockville, MD, USA, 2007. Available online: https://meps.ahrq.gov/data_files/publications/cb17/cb17.pdf (accessed on 14 November 2018).
182. Griffin, S.O.; Gooch, B.F.; Beltrán, E.; Sutherland, J.N.; Barsley, R. Dental services, costs, and factors associated with hospitalization for Medicaid-eligible children, Louisiana 1996–1997. *J. Public Health Dent.* **2000**, *60*, 21–27. [[CrossRef](#)] [[PubMed](#)]
183. Gambhir, R.S.; Gupta, T. Need for Oral Health Policy in India. *Ann. Med. Health Sci. Res.* **2016**, *6*, 50–55. [[CrossRef](#)] [[PubMed](#)]
184. Parkash, H.; Shah, N. *National Oral Health Care Programme: Implementation Strategies*; National Oral Health Care Programme: New Delhi, India, 2001.
185. Thomas, S.; Beh, L.; Nordin, R.B. Health care delivery in Malaysia: Changes, challenges, and champions. *J. Public Health Afr.* **2011**, *2*, E23. [[CrossRef](#)] [[PubMed](#)]

186. Oral Health Division, Ministry of Health Malaysia. Oral Healthc. Malays. 2005. Available online: https://www.mah.se/upload/FAKULTETER/OD/Avdelningar/who/WPRO/Malaysia/data/oral_healthcare_in_malaysia_05.pdf (accessed on 14 November 2018).
187. Cascaes, A.M.; Camargo, M.B.J.; Castilhos, E.D.; Silva, L.E.R.; Barros, A.J.D. Private dental insurance expenditure in Brazil. *Rev. Saude Publica* **2018**, *52*, 24. [[CrossRef](#)] [[PubMed](#)]
188. Narvai, P.C. Collective oral health: Ways from sanitary dentistry to buccality. *Rev. Saude Publica* **2006**, *40*, 141–147. [[CrossRef](#)] [[PubMed](#)]
189. Petersen, P.E. Strengthening of oral health systems: Oral health through primary health care. *Med. Princ. Pract.* **2014**, *23*, 3–9. [[CrossRef](#)] [[PubMed](#)]
190. Mumghamba, E.G. Integrating a primary oral health care approach in the dental curriculum: A Tanzanian experience. *Med. Princ. Pract.* **2013**, *23*, 69–77. [[CrossRef](#)] [[PubMed](#)]
191. Honkala, E. Primary oral health care. *Med. Princ. Pract.* **2014**, *23*, 17–23. [[CrossRef](#)] [[PubMed](#)]
192. Lassen, E.M.; Mayrhofer, M.; Vedel Kessing, P.; Sano, H.O.; García San José, D.; Frank Jørgensen, R. Factors which Enable or Hinder the Protection of Human Rights. The Danish Institute of Human Rights. Frame Fostering Human Rights among European (External and Internal) Policies. Large-Scale Fp7 Collaborative Project. GA no. 320000, 2017. Available online: https://www.humanrights.dk/sites/humanrights.dk/files/media/dokumenter/udgivelser/research/frame/frame_-_factors_which_enable_or_hinder_the_protection_of_human_rights.pdf (accessed on 14 January 2019).
193. Meier, B.M. Human Rights in the World Health Organization: Views of the Director-General Candidates. *Health Hum. Rights* **2017**, *19*, 293–298. [[PubMed](#)]
194. Okoro, C. Primary Health Care in London: Onwards from Alma Ata. *Lond. J. Prim. Care* **2008**, *1*, 61–65. [[CrossRef](#)]
195. De Maeseneer, J.; Willems, S.; Sutter, A.D.; Van de Geuchte, I.; Billings, M. Primary Health Care as a Strategy for Achieving Equitable Care: A Literature Review Commissioned by the Health Systems Knowledge Network. 2007. Available online: https://www.who.int/social_determinants/resources/csdh_media/primary_health_care_2007_en.pdf (accessed on 14 January 2019).
196. Kandelman, D.; Arpin, S.; Baez, R.J.; Baehni, P.C.; Petersen, P.E. Oral health care systems in developing and developed countries. *Periodontol. 2000* **2012**, *60*, 98–109. [[CrossRef](#)] [[PubMed](#)]
197. Petersen, P.E. Challenges to improvement of oral health in the 21st century—the approach of the WHO Global Oral Health Programme. *Int. Dent. J.* **2004**, *54*, 329–343. [[CrossRef](#)] [[PubMed](#)]
198. Petersen, P.E.; Yamamoto, T. Improving the oral health of older people: The approach of the WHO Global Oral Health Programme. *Community Dent. Oral Epidemiol.* **2005**, *33*, 81–92. [[CrossRef](#)] [[PubMed](#)]
199. Palmer, R.J. Composition and development of oral bacterial communities. *Periodontol. 2000* **2014**, *64*, 20–39. [[CrossRef](#)] [[PubMed](#)]
200. Franco, B.E.; Altigracia Martínez, M.; Sánchez Rodríguez, M.A.; Wertheimer, A.I. The determinants of the antibiotic resistance process. *Infect. Drug Resist.* **2009**, *2*, 1–11. [[PubMed](#)]
201. Ventola, C.L. The antibiotic resistance crisis: Part 1, causes and threats. *Pharm. Ther.* **2015**, *40*, 277–283.
202. Fleming-Dutra, K.E.; Hersh, A.L.; Shapiro, D.J.; Bartoces, M.; Enns, E.A.; File, T.M., Jr.; Finkelstein, J.A.; Gerber, J.S.; Hyun, D.Y.; Linder, J.A.; et al. Prevalence of Inappropriate Antibiotic Prescriptions among US Ambulatory Care Visits, 2010–2011. *JAMA* **2016**, *315*, 1864–1873. [[CrossRef](#)] [[PubMed](#)]
203. Centers for Disease Control and Prevention (CDC). Office-related antibiotic prescribing for persons aged ≤ 14 years—United States, 1993–1994 to 2007–2008. *MMWR Morb. Mortal. Wkly. Rep.* **2011**, *60*, 1153–1156.
204. Shapiro, D.J.; Hicks, L.A.; Pavia, A.T.; Hersh, A.L. Antibiotic prescribing for adults in ambulatory care in the USA, 2007–2009. *J. Antimicrob. Chemother.* **2014**, *69*, 234–240. [[CrossRef](#)] [[PubMed](#)]
205. Gonzales, R.; Malone, D.C.; Maselli, J.H.; Sande, M.A. Excessive antibiotic use for acute respiratory infections in the United States. *Clin. Infect. Dis.* **2001**, *33*, 757–762. [[CrossRef](#)] [[PubMed](#)]
206. Hicks, L.A.; Bartoces, M.G.; Roberts, R.M.; Suda, K.J.; Hunkler, R.J.; Taylor, T.H., Jr.; Schrag, S.J. US outpatient antibiotic prescribing variation according to geography, patient population, and provider specialty in 2011. *Clin. Infect. Dis.* **2015**, *60*, 1308–1316. [[CrossRef](#)] [[PubMed](#)]
207. Wasan, H.; Gupta, P.; Mathur, A.; Mutneja, E.; Mathur, V.P.; Gupta, Y.K. Influence of Qualification and Practice Settings of Dental Practitioners on Antimicrobial Prescribing in Delhi and National Capital Region, India. *J. Nat. Sci. Biol. Med.* **2017**, *8*, 229–234. [[PubMed](#)]

208. Löffler, C.; Böhmer, F. The effect of interventions aiming to optimize the prescription of antibiotics in dental care-A systematic review. *PLoS ONE* **2017**, *12*, e0188061. [[CrossRef](#)] [[PubMed](#)]
209. Segura-Egea, J.J.; Martín-González, J.; Jiménez-Sánchez, M.D.C.; Crespo-Gallardo, I.; Saúco-Márquez, J.J.; Velasco-Ortega, E. Worldwide pattern of antibiotic prescription in endodontic infections. *Int. Dent. J.* **2017**, *67*, 197–205. [[CrossRef](#)] [[PubMed](#)]
210. Garg, A.K.; Agrawal, N.; Tewari, R.K.; Kumar, A.; Chandra, A. Antibiotic prescription pattern among Indian oral healthcare providers: A cross-sectional survey. *J. Antimicrob. Chemother.* **2014**, *69*, 526–528. [[CrossRef](#)] [[PubMed](#)]
211. Cope, A.; Francis, N.; Wood, F.; Mann, M.K.; Chestnutt, I.G. Systemic antibiotics for symptomatic apical periodontitis and acute apical abscess in adults. *Cochrane Database Syst. Rev.* **2014**, *26*, CD010136. [[CrossRef](#)] [[PubMed](#)]
212. Iqbal, A. The Attitudes of Dentists Towards the Prescription of Antibiotics During Endodontic Treatment in North of Saudi Arabia. *J. Clin. Diagn. Res.* **2015**, *9*, ZC82. [[CrossRef](#)] [[PubMed](#)]
213. Ramu, C.; Padmanabhan, T.V. Indications of antibiotic prophylaxis in dental practice- review. *Asian Pac. J. Trop. Biomed.* **2012**, *2*, 749–754. [[CrossRef](#)]
214. Habib, G.; Lancellotti, P.; Antunes, M.J.; Bongiorni, M.G.; Casalta, J.P.; Del Zotti, F.; Dulgheru, R.; El Khoury, G.; Erba, P.A.; Iung, B.; et al. 2015 ESC Guidelines for the management of infective endocarditis: The Task Force for the Management of Infective Endocarditis of the European Society of Cardiology (ESC). Endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the European Association of Nuclear Medicine (EANM). *Eur. Heart J.* **2015**, *36*, 3075–3128. [[PubMed](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).