

Supplementary Table S1. Details about distribution of the study population regarding the species, age, gender and body condition score of animals, and the year of sampling.

	N total	Age		Gender			Body Condition Score					Year of sampling			
		Young	Adult	Female	Male	Unknown	0/5	1/5	2/5	3/5	4/5	2018	2019	2020	2021
Barn owl (<i>Tyto alba</i>)	90	72/90 80%	18/90 20%	20/90 22.2%	20/90 22.2%	50/90 55.6%	0/90 0%	0/90 0%	19/90 21.1%	71/90 78.9%	0/90 0%	11/90 12.2%	28/90 31.1%	28/90 31.1%	23/90 25.6%
Eurasian eagle-owl (<i>Bubo bubo</i>)	19	3/19 15.8%	16/19 84.2%	7/19 36.8%	5/19 26.4%	7/19 36.8%	4/19 21.1%	2/19 10.5%	10/19 52.6%	3/19 15.8%	0/19 0%	9/19 47.4%	4/19 21.1%	3/19 15.8%	3/19 15.8%
Tawny owl (<i>Strix aluco</i>)	10	1/10 10%	9/10 90%	1/10 10%	1/10 10%	8/10 80%	1/10 10%	0/10 0%	2/10 20%	7/10 70%	0/10 0%	5/10 50%	1/10 10%	2/10 10%	2/10 10%
European owl (<i>Athene noctua</i>)	10	5/10 50%	5/10 50%	2/10 20%	1/10 10%	7/10 70%	0/10 0%	1/10 10%	1/10 10%	7/10 70%	1/10 10%	8/10 80%	0/10 0%	0/10 0%	2/10 20%
Eurasian Scops-owl (<i>Otus scops</i>)	4	3/4 75%	1/4 25%	0/4 0%	0/4 0%	4/4 100%	0/4 0%	0/4 0%	0/4 0%	4/4 100%	0/4 0%	3/4 75%	1/4 25%	0/4 0%	0/4 0%
Northern long-eared owl (<i>Asio otus</i>)	1	0/1 0%	1/1 100%	0/1 0%	1/1 100%	0/1 0%	0/1 0%	0/1 0%	1/1 100%	0/1 0%	0/1 0%	0/1 0%	0/1 0%	1/1 100%	0/1 0%
TOTAL	134	84/134 62.7%	50/134 37.3%	30/134 22.4%	28/134 20.9%	76/134 56.7%	5/134 3.7%	3/134 2.2%	33/134 24.6%	92/134 68.7%	1/134 0.7%	36/134 26.9%	34/134 25.4%	34/134 25.4%	30/134 22.3%

N = number of raptors; % = percentage