

Supplementary material

Table S1. List of oligonucleotides

<i>CgACT1 F</i>	TTCAACGTTCCAGCCTTCT
<i>CgACT1 R</i>	GTAACACCGTCACCAAGAGT
<i>CgYAP1 F</i>	GTGGACTTGGAGAACTTGAC
<i>CgYAP1 R</i>	ACTTCCGCAACTCTTCAC
<i>CgMSN4 F</i>	CATTAATCCCGTATTGCTGTCC
<i>CgMSN4 R</i>	CTTCGTTGTCGTAATCATTGCC
<i>CgCTA1 F</i>	ACAAACTAACTGAAGGACCCA
<i>CgCTA1 R</i>	GACATAGTGCCATTCACCT
<i>CgSOD1 F</i>	CCACTATCACCTACGAGATTGC
<i>CgSOD1 R</i>	CACCGACGTGTCTGTTCTC
<i>CgSOD2 F</i>	CACTACTCCAAGCACCACC
<i>CgSOD2 R</i>	TTCGTGTACCCACCTCCA
<i>CgSKN7 F</i>	ATGAGCCCTTCATTGCCT
<i>CgSKN7 R</i>	TGGCGATTCAAGTTGTCCCT
<i>CgAFT1 F</i>	TACCCGCAGGAAATAGAGATCG
<i>CgAFT1 R</i>	TTCATGTTCCCTAGCCAAAGCC
<i>CgAFT2 F</i>	GCCATGAACCAAAGGACCTACG
<i>CgAFT2 R</i>	GTAGAGAAAAGTACACCGCACCC
<i>CgYAP5 F</i>	GTTGATCTGGATGGTCTGAAGG
<i>CgYAP5 R</i>	TCGGAGTAATTGAAGGTGAGC
<i>CgCCC1 F</i>	GTTCAGCAATGTGGACCCGC
<i>CgCCC1 R</i>	GATCACAGCTTCGCGTCAC
<i>CgFTR1 F</i>	AGCCGGTTGTTCTCAAGAGG
<i>CgFTR1 R</i>	CACCACTTCAGAACATCACC
<i>CgISU1 F</i>	CGTCCGTTTACCAAGGATGTACC
<i>CgISU1 R</i>	GTTCCGCAGAGTCTTGTCCA

Table S2. Intracellular iron content in *C. glabrata* strains

The amount of Fe in the samples was measured using an atomic absorption spectrometer in flame emission mode (Perkin Elmer 1100). Cells were incubated in the absence or presence of 1mM FeCl₃. * P <0,001

µg Fe / 10¹⁰ cells		
FeCl₃	wt	<i>Cgerg6A</i>
-	7.3 ± 0.0	6.1 ± 0.1
+	576.5 ± 2.5*	1305 ± 25*