



Supplementary Material

Table S1. Individual phenolic content in peel extract of small (S), medium (M), and large (L) avocado fruit.

Phenolic compound	Fruit Size Group		
	S	M	L
	µg/g of dry matter		
(-) Epicatechin	24.1 ± 2.2 a	18.5 ± 0.1 b	4.8 ± 0.5 c
Hesperidin	11.9 ± 1.9 a	6.9 ± 0.6 a	9.3 ± 0.9 a
(+) Catechin hydrate	5.3 ± 1.3 a	5.1 ± 0.1 a	1.6 ± 0.3 b
Kaempferol	5.1 ± 0.2 a	5.8 ± 1.0 a	5.6 ± 0.2 a
(±) Narangenin	4.1 ± 0.2 a	5.6 ± 1.3 a	1.5 ± 0.1 b
Naringin	3.6 ± 0.7 a	2.9 ± 0.6 a.b	1.2 ± 0.0 b
Caffeic acid	2.4 ± 0.2 a	2.3 ± 0.2 a	0.9 ± 0.0 b
Rutin hydrate	2.2 ± 0.4 a	2.5 ± 0.3 a	1.7 ± 0.0 a
Coumaric acid	1.9 ± 0.2 a	1.6 ± 0.2 a	0.4 ± 0.1 b
Quercetin	0.6 ± 0.2 a.b	0.2 ± 0.3 a	1.1 ± 0.0 b
Ferulic acid	0.7 ± 0.2 a	0.4 ± 0.1 a	0.0 ± 0.0

Mean ± standard deviation (n=2). Different letters in the same row represent significant differences according to the Tukey's test ($P < 0.05$).



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).