



## Correction

# Correction: Herfort et al. Influence of Sheep's Wool Vegetation Mats on the Plant Growth of Perennials. *Horticulturae* 2023, 9, 384

Susanne Herfort <sup>1,\*</sup>, Kerstin Pflanz <sup>1</sup>, Marina-Sandra Larsen <sup>1</sup>, Thomas Mertschun <sup>2</sup> and Heiner Grüneberg <sup>3,\*</sup>

<sup>1</sup> Institute of Agricultural and Urban Ecological Projects, Humboldt-Universität zu Berlin (IASP), 10115 Berlin, Germany

<sup>2</sup> Uabg Society for Environmental Analysis Soil and Water Protection mbH, 12459 Berlin, Germany

<sup>3</sup> Faculty of Life Sciences, Albrecht Daniel Thaer-Institute of Agricultural and Horticultural Sciences, Urban Ecophysiology of Plants, 14195 Berlin, Germany

\* Correspondence: susanne.herfort@iasp.hu-berlin.de (S.H.); hgrueneberg@hu-berlin.de (H.G.)

There was an error in the original publication [1]. A correction has been made to Abstract, 3.1. Nitrogen Content of the Vegetation Mats, paragraph 1, Tables 3 and 7.

**Abstract:** The vegetation mats with sheep's wool (V1–V3) contained 192.6, 154.0, and 283.5 g nitrogen (N)/m<sup>2</sup> and the coir mats (V4) contained 7.5 g N/m<sup>2</sup>.

**3.1. Nitrogen Content of the Vegetation Mats,** paragraph 1: The analysis of total nitrogen showed 10.4% for the sheep's wool fibres and 0.3% for the coconut fibres. The subsequently calculated nitrogen content in the four vegetation mat types averaged between 7.5 g of nitrogen (variant 100c) and 283.5 g of nitrogen (sandwiched variant) per m<sup>2</sup> vegetation mat (Table 3).

**Table 3.** Calculated nitrogen content of the fibres (air dry) of the used vegetation mats.

Mat Variant	N-Content of Sheep's Wool Fibres in the Vegetation Mat (g/m <sup>2</sup> )	N-Content of Coconut Fibre in the Vegetation Mat (g/m <sup>2</sup> )	Total N-Content of Fibres in the Vegetation Mat (g/m <sup>2</sup> )
V1: 50sw/50c	187.2	5.4	192.6
V2: 30sw/70c	145.6	8.4	154.0
V3: Sandwich	280.8	2.7	283.5
V4: 100c	0	7.5	7.5

**Table 7.** Pearson correlation between nitrogen content and plant height.

	1st Date	3rd Date	5th Date	7th Date
<i>Achillea clypeolata</i> 'Moonshine'	-	-	-	0.617 **
Sig. (2-tailed)	-	-	-	<0.001
<i>Aster dumosus</i> 'Prof. Anton Kippenberg'	0.542 **	0.808 **	0.878 **	0.834 **
Sig. (2-tailed)	<0.001	<0.001	<0.001	<0.001
<i>Aster dumosus</i> 'Silberball'	0.572 **	0.817 **	0.792 **	0.807 **
Sig. (2-tailed)	<0.001	<0.001	<0.001	<0.001
<i>Centranthus ruber</i> 'Coccineus'	-	-	0.372 **	-
Sig. (2-tailed)	-	-	0.009	-
<i>Salvia nemorosa</i> 'Rosakönigin'	0.437 **	0.536 **	0.561 **	0.446 **
Sig. (2-tailed)	0.002	<0.001	<0.001	0.002

\*\* The correlation is significant at a level of 0.01 (2-tailed).

The authors apologize for any inconvenience caused and state that the scientific conclusions are unaffected. The original publication has also been updated.



**Citation:** Herfort, S.; Pflanz, K.; Larsen, M.-S.; Mertschun, T.; Grüneberg, H. Correction: Herfort et al. Influence of Sheep's Wool Vegetation Mats on the Plant Growth of Perennials. *Horticulturae* 2023, 9, 384. *Horticulturae* 2023, 9, 1166. <https://doi.org/10.3390/horticulturae9111166>

Received: 5 October 2023

Accepted: 13 October 2023

Published: 25 October 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## Reference

1. Herfort, S.; Pflanz, K.; Larsen, M.-S.; Mertschun, T.; Grüneberg, H. Influence of Sheep's Wool Vegetation Mats on the Plant Growth of Perennials. *Horticulturae* **2023**, *9*, 384. [[CrossRef](#)]

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.