



Supplementary Materials of Detecting Enteric Pathogens in Low-Risk Drinking Water in Dhaka, Bangladesh: An Assessment of the WHO Water Safety Categories

Table S1. Target genes, primer sequences, amplicon size for detection of the signature and pathogenic genes of bacteria.

Genes	Primer Name	Primer Sequences (5' → 3')	Amplicon Size (bp)	References
Diarrheagenic <i>E. coli</i> genes				
<i>eltB</i>	LT-F	tctctatgtgcatacgaggc	322	[1]
	LT-R	ccatactgattggccaaat		
<i>estA</i>	ST-F	gctaaaccagta ^a ggtcttcaaaa	147	[1]
	ST-F	cccggtac ^a gcaggattacaaca		
<i>vt1</i>	VT1-F	gaagagtccgtgggattacg	130	[2]
	VT1-R	agcgatgcagcttataaa		
<i>vt2</i>	VT2-F	accgttttcagattt ^a cacata	298	[1]
	VT2-R	tacacaggaggagttcagacagt		
<i>eaeA</i>	eae-F	cacacgaataaaactgactaaaatg	376	[1]
	eae-R	aaaaaacgtgacccgcacctaaat		
<i>bfpA</i>	bfpA-F	ttcttggtgcttgcgtctttt	367	[1]
	bfpA-R	ttttgttgttatctttgtaa		
<i>ipaH</i>	ipaH-F	gctggaaaaactcagfgcct	424	[3]
	ipaH-R	ccagtcgttattcattct		
pCVD	EA-F	ctggcgaagactgtatcat	630	[4]
	EA-R	caatgtatagaaatccgtt		
<i>V. cholerae</i> species specific gene				
<i>ompW</i>	ompW-F	caccaagaaggtaactttatgt	304	[5]
	ompW-R	ggtttgtcaatttagcttccacc		
<i>Salmonella</i> spp.				
<i>invA</i>	S-invA-F	cggtggtttaagcgtactctt	796	[6]
	S-invA-R	cgaatatgcgtccacaaggta		
<i>Salmonella enteritidis</i>				
<i>IE-1</i>	SE-IE1-F	agtgcatactttaatgac	316	[7]
	SE-IE1-R	actatgtcgatacggcgg		
<i>Salmonella typhimurium</i>				
<i>Flic-C</i>	ST-flic C-F	cccgcttacagggtggactac	432	[8]
	ST-flic C-R	agcgggtttcgggtgtt		
<i>Campylobacter</i> spp.				
16S rRNA	C412F	ggatgacactttcgagc	816	[9]
	C1228R	cattgttagcacgtgtgtc		
<i>Campylobacter coli</i>				
<i>cueE</i>	CC18F	ggtagatgttctaaagcgg	502	[9]
	CC519R	ataaaagactatcgccgtg		
<i>Campylobacter jejuni</i>				
cj0414	CJ-F	caaataaaggtagaggtagaatgt	161	[10]
	CJ-R	ccataaggactagctagctgat		

References

1. Svenungsson, B.; Lagergren, Å.; Ekwall, E.; Evengård, B.; Hedlund, K.O.; Kärnell, A.; Löfdahl, S.; Svensson, L.; Weintraub, A. Enteropathogens in adult patients with diarrhea and healthy control subjects: A 1-year prospective study in a Swedish clinic for infectious diseases. *Clin. Infect. Dis.* **2000**, *30*, 770–778.
2. Pollard, D.; Johnson, W.; Lior, H.; Tyler, S.; Rozee, K. Rapid and specific detection of verotoxin genes in Escherichia coli by the polymerase chain reaction. *J. Clin. Microbiol.* **1990**, *28*, 540–545.
3. Tornieporth, N.G.; John, J.; Salgado, K.; de Jesus, P.; Latham, E.; Melo, M.; Gunzburg, S.T.; Riley, L.W. Differentiation of pathogenic Escherichia coli strains in Brazilian children by PCR. *J. Clin. Microbiol.* **1995**, *33*, 1371–1374.
4. Schmidt, H.; Knop, C.; Franke, S.; Aleksic, S.; Heesemann, J.; Karch, H. Development of PCR for screening of enteroaggregative Escherichia coli. *J. Clin. Microbiol.* **1995**, *33*, 701–705.
5. Nandi, B.; Nandy, R.K.; Mukhopadhyay, S.; Nair, G.B.; Shimada, T.; Ghose, A.C. Rapid method for species-specific identification of Vibrio cholerae using primers targeted to the gene of outer membrane protein OmpW. *J. Clin. Microbiol.* **2000**, *38*, 4145–4151.
6. Fratamico, P.M. Comparison of culture, polymerase chain reaction (PCR), TaqMan Salmonella, and Transia Card Salmonella assays for detection of Salmonella spp. in naturally-contaminated ground chicken, ground turkey, and ground beef. *Mol. Cell. Probes* **2003**, *17*, 215–221.
7. Wang, S.J.; Yeh, D.B. Designing of polymerase chain reaction primers for the detection of Salmonella enteritidis in foods and faecal samples. *Lett. Appl. Microbiol.* **2002**, *34*, 422–427.
8. Paião, F.; Arisitides, L.; Murate, L.; Vilas-Bôas, G.; Vilas-Boas, L.; Shimokomaki, M. Detection of Salmonella spp., Salmonella Enteritidis and Typhimurium in naturally infected broiler chickens by a multiplex PCR-based assay. *Braz. J. Microbiol.* **2013**, *44*, 37–42.
9. Linton, D.; Lawson, A.; Owen, R.; Stanley, J. PCR detection, identification to species level, and fingerprinting of Campylobacter jejuni and Campylobacter coli direct from diarrheic samples. *J. Clin. Microbiol.* **1997**, *35*, 2568–2572.
10. Wang, R.F.; Slavik, M.F.; Cao, W.W. A Rapid PCR Method for Direct Detection of Low Numbers of Campylobacter Jejuni. *J. Rapid Methods Autom. Microbiol.* **1992**, *1*, 101–108.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.