

Article

The Zeldovich Number: A Universal Dimensionless Measure for the Electromagnetic Field

Iwo Bialynicki-Birula ^{1,*} and Zofia Bialynicka-Birula ²¹ Center for Theoretical Physics, Polish Academy of Sciences, Aleja Lotników 32/46, 02-668 Warsaw, Poland² Institute of Physics, Polish Academy of Sciences, Aleja Lotników 32/46, 02-668 Warsaw, Poland; zbbirula@gmail.com

* Correspondence: birula@cft.edu.pl

Abstract: In this paper we extend the Zeldovich formula, which was originally derived for the free electromagnetic field and was interpreted as the number of photons. We show that our extended formula gives a universal dimensionless measure of the overall strength of electromagnetic fields: free fields and fields produced by various sources, in both the classical and the quantum theory. In particular, we find that this number—called here the Zeldovich number—for macroscopic systems is very large, of the order of 10^{20} . For the hydrogen atom in the ground state, the Zeldovich number is equal to 0.025 and for the xenon atom it is around 50.

Keywords: Zeldovich number; photon number; measure of the electromagnetic field

1. Introduction

The formula measuring the number of photons was derived by Yakov B. Zeldovich for the free electromagnetic field. In the original derivation [1] of this formula, Zeldovich assumed that photons can be identified with monochromatic oscillations of the electromagnetic field. However, we will show, that this formula is universal; it can be used without any restrictions for all electromagnetic fields. In the quantum theory of the electromagnetic field the Zeldovich formula plays multiple roles. It can be used to measure the probability of various field configurations and to define the norm of quantum photon states.

In this study we extend the use of the Zeldovich formula from free fields to field configurations generated by various sources. In particular, we will study the problem of the photons attached to the hydrogen atom which has been treated in nonrelativistic case by Francesco Persico and his collaborators [2–5]. The extension to the relativistic theory was treated in [6]. All these attempts have not given a definite answer to the question: How many photons are attached to an atom? We obtain an answer to this question employing the notion of the Zeldovich number.

Since in the general case the name “photon number” is not always justified, we shall use instead the term “Zeldovich number” and we denote it by N_Z . The number N_Z is a *dimensionless* quantity which measures the overall strength of the electromagnetic field. This number is a useful characterization of the field and its sources.

In Section 2, we extend the derivation of the formula for N_Z to a general electromagnetic field. In Section 3, we calculate N_Z in classical electrodynamics. In Sections 4 and 5, we extend the calculations to atomic physics, nonrelativistic and relativistic. This is done by associating the wave functions of electrons with electromagnetic fields.

2. The Zeldovich Number

The starting point of our calculations is the formula for the total energy of the electromagnetic field,



Citation: Bialynicki-Birula, I.; Bialynicka-Birula, Z. The Zeldovich Number: A Universal Dimensionless Measure for the Electromagnetic Field. *Physics* **2023**, *5*, 862–872. <https://doi.org/10.3390/physics5030055>

Received: 6 June 2023

Accepted: 11 July 2023

Published: 15 August 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

$$\mathcal{E}(t) = \frac{1}{2} \int d^3r [\mathbf{E}(\mathbf{r}, t) \cdot \mathbf{D}(\mathbf{r}, t) + \mathbf{B}(\mathbf{r}, t) \cdot \mathbf{H}(\mathbf{r}, t)]. \tag{1}$$

We do not assume, as has been done by Zeldovich, that the field vectors satisfy the free Maxwell equations. It is sufficient that the field vectors are sufficiently regular to have the Fourier transforms.

It is convenient to use the field vectors, \mathcal{D} and \mathcal{H} measured in purely geometrical units, i.e., in $1/m^2$. The \mathcal{D} and \mathcal{H} vectors are related to the standard physical vectors \mathbf{D} and \mathbf{H} by the formulas,

$$\mathcal{D} = \frac{\mathbf{D}}{e}, \quad \mathcal{H} = \frac{\mathbf{H}}{ec}. \tag{2}$$

The expression for the field energy expressed in terms of the new vectors is,

$$\mathcal{E}(t) = \frac{e^2}{2\epsilon_0} \int d^3r [\mathcal{D}(\mathbf{r}, t) \cdot \mathcal{D}(\mathbf{r}, t) + \mathcal{H}(\mathbf{r}, t) \cdot \mathcal{H}(\mathbf{r}, t)]. \tag{3}$$

Next, we rewrite this formula in terms of the Fourier transforms,

$$\mathcal{E}(t) = \frac{e^2}{2\epsilon_0} \int d^3k [\tilde{\mathcal{D}}^*(\mathbf{k}, t) \cdot \tilde{\mathcal{D}}(\mathbf{k}, t) + \tilde{\mathcal{H}}^*(\mathbf{k}, t) \cdot \tilde{\mathcal{H}}(\mathbf{k}, t)]. \tag{4}$$

The original formula for \mathcal{N}_Z will be obtained in two steps. In the first step we divide the integrand in Equation (4) by $\hbar ck$ to obtain a dimensionless quantity,

$$\mathcal{N}_Z[\mathcal{D}, \mathcal{H}] = 2\pi\alpha \int \frac{d^3k}{k} [\tilde{\mathcal{D}}^*(\mathbf{k}, t) \cdot \tilde{\mathcal{D}}(\mathbf{k}, t) + \tilde{\mathcal{H}}^*(\mathbf{k}, t) \cdot \tilde{\mathcal{H}}(\mathbf{k}, t)], \tag{5}$$

where α is the fine structure constant. This is the representation in terms of Fourier transforms. In the next step, to obtain the original form of \mathcal{N}_Z we convert Equation (5) back to the \mathbf{r} -space with the use of the relations,

$$\tilde{\mathcal{D}}(\mathbf{k}, t) = \int \frac{d^3r}{(2\pi)^{3/2}} e^{-ik \cdot r} \mathcal{D}(\mathbf{r}, t), \tag{6}$$

$$\tilde{\mathcal{H}}(\mathbf{k}, t) = \int \frac{d^3r}{(2\pi)^{3/2}} e^{-ik \cdot r} \mathcal{H}(\mathbf{r}, t), \tag{7}$$

and the formula,

$$\int \frac{d^3k}{k} e^{ik \cdot (r-r')} = \frac{1}{2\pi^2 |r-r'|^2}. \tag{8}$$

The resulting expression has the form obtained by Zeldovich (apart from our different scaling of the electromagnetic field),

$$\mathcal{N}_Z[\mathcal{D}, \mathcal{H}] = \frac{\alpha}{\pi} \int d^3r \int d^3r' \left(\mathcal{D}(\mathbf{r}, t) \frac{1}{|r-r'|^2} \cdot \mathcal{D}(\mathbf{r}', t) + \mathcal{H}(\mathbf{r}, t) \frac{1}{|r-r'|^2} \cdot \mathcal{H}(\mathbf{r}', t) \right). \tag{9}$$

The derivation of this expression by Zeldovich was based on the expansion of the solutions of Maxwell equations in free space into monochromatic propagating waves. We arrived at the same formula without making any assumptions concerning the dynamics of the electromagnetic field. Certainly, the interpretation of \mathcal{N}_Z in this general case as the photon number is highly problematic, but \mathcal{N}_Z is very well defined.

In the case of free fields, \mathcal{N}_Z is the total number of photons and it has some remarkable properties. It is a constant of motion and despite its nonrelativistic appearance it is invariant

not only under all Lorentz transformations, but also under the conformal transformations [7]. Since conformal transformations include the transformations to accelerated frames of reference, the invariance of \mathcal{N}_Z may help to better understand the Unruh effect [8]. As a mathematical object, \mathcal{N}_Z plays the role of a norm for the photon wave function [7,9,10]. The scalar product obtained from this norm by polarization identity serves as a perfect measure of fidelity for photon states [11]. It also appears as the exponent in the Wigner functional of the electromagnetic field [12–14] (there are some misplaced factors of 2 in this reference that were corrected in [13]). In this way it determines the relative probabilities of various field configurations. Due to its connection to the Wigner functional, \mathcal{N}_Z can be generalized to characterize also thermal states of electromagnetic fields [13,14].

3. The Zeldovich Number for Macroscopic Fields

In this Section, we calculate \mathcal{N}_Z for field configurations created by the following classical sources: (i) the two oppositely-charged metallic spheres, and (ii) the current flowing in a circular loop.

In the case of the two oppositely-charged metallic spheres, the charged density is,

$$\rho(\mathbf{r}) = \frac{Q}{4\pi a^2} [\delta(a - |\mathbf{r} + \mathbf{d}/2|) - \delta(a - |\mathbf{r} - \mathbf{d}/2|)], \tag{10}$$

where a is the sphere radius, $|\mathbf{d}|$ is the distance between the spheres, and Q is the total charge. We assume that the charge is distributed uniformly on the surface. We choose the z direction along the vector \mathbf{d} and use the spherical coordinates. To calculate \mathcal{N}_Z we will need the Fourier transform the $\tilde{\rho}(\mathbf{k})$,

$$\begin{aligned} \tilde{\rho}(\mathbf{k}) &= \int \frac{d^3r}{(2\pi)^{3/2}} \rho(\mathbf{r}) e^{-i\mathbf{k}\cdot\mathbf{r}} \\ &= \frac{2iQ}{(2\pi)^{3/2}} \sin(dk/2 \cos \theta) \frac{\sin(ak)}{ak}, \end{aligned} \tag{11}$$

where we took into account that a shift by $\pm \mathbf{d}/2$ in the position space results in the multiplication by $e^{\pm i\mathbf{k}\cdot\mathbf{d}/2}$ in the Fourier space. For electrostatic fields, the Fourier transform $\tilde{\mathcal{D}}$ of the displacement vector is,

$$e\tilde{\mathcal{D}}(\mathbf{k}) = -i \frac{\mathbf{k}\tilde{\rho}(\mathbf{k})}{k^2}. \tag{12}$$

The substitution of this expression into Equation (5) gives,

$$\mathcal{N}_Z = 2\pi\alpha \left(\frac{Q}{e}\right)^2 \int_0^\infty \frac{dk}{k} \int_0^\pi d\theta \sin \theta \int_0^{2\pi} d\phi \tilde{\rho}^*(\mathbf{k}) \tilde{\rho}(\mathbf{k}). \tag{13}$$

The integrals in this formula can be analytically calculated and the final result, as expected, depends only on the dimensionless ratio, $b = d/a$,

$$\mathcal{N}_Z = \frac{\alpha}{12\pi b} \left(\frac{Q}{e}\right)^2 [(b+2)^3 \ln(b+2) + (b-2)^3 \ln(|b-2|) - 2b(4 + 12 \ln 2 + b^2 \ln b)]. \tag{14}$$

When b is very large, we obtain,

$$\mathcal{N}_Z \approx \frac{2\alpha}{\pi} \left(\frac{Q}{e}\right)^2 \ln b. \tag{15}$$

The unbounded logarithmic growth of \mathcal{N}_Z is the manifestation of the infrared catastrophe, well known in quantum electrodynamics. In the case of a large separation the Coulomb fields of each sphere is practically not shielded and \mathcal{N}_Z for an unshielded charge is infinite.

In the case of the current I flowing in a closed loop, the current density has the following form in cylindrical coordinates [15],

$$\mathbf{j}(\rho, z, \phi) = I\delta(a - \rho)\delta(z)\{-\sin \phi, \cos \phi, 0\}. \tag{16}$$

To calculate \mathcal{N}_Z , one needs the Fourier transform of the current,

$$\begin{aligned} \tilde{\mathbf{j}}(\mathbf{k}) &= \int \frac{d^3r}{(2\pi)^{3/2}} e^{-i\mathbf{k}\cdot\mathbf{r}} \mathbf{j}(\rho, z, \phi) \\ &= \frac{aI}{(2\pi)^{3/2}} \int_0^{2\pi} d\phi e^{-ia(k_x \cos \phi + k_y \sin \phi)} \{-\sin \phi, \cos \phi, 0\} \\ &= \frac{iaI}{\sqrt{2\pi}} \frac{J_1(ak_\perp)}{k_\perp} \{k_y, -k_x, 0\}, \end{aligned} \tag{17}$$

where $k_\perp = \sqrt{k_x^2 + k_y^2}$ and J_1 is the Bessel function. For static fields, the Fourier transform of the magnetic field vector is,

$$ec\tilde{\mathbf{H}}(\mathbf{k}) = -i\frac{\mathbf{k} \times \tilde{\mathbf{j}}(\mathbf{k})}{k^2}. \tag{18}$$

The substitution of this expression into Equation (5) gives,

$$\begin{aligned} \mathcal{N}_Z &= \alpha \left(\frac{aI}{ec}\right)^2 \int_0^\infty dk_\perp \int_{-\infty}^\infty dk_z \int_0^{2\pi} d\phi \frac{k_\perp J_1(ak_\perp)^2}{(k_\perp^2 + k_z^2)^{3/2}} \\ &= 4\pi\alpha \left(\frac{aI}{ec}\right)^2. \end{aligned} \tag{19}$$

Note that the increase of the ring radius and the proportional decrease of the current leaves the Zeldovich number unchanged.

For macroscopic systems \mathcal{N}_Z is very large. In the electrostatic case even for tiny charges of one microcoulomb on both spheres and for $b = 10$, one obtains \mathcal{N}_Z equal to 1.6×10^{20} . In turn, for the current of one ampere flowing in a loop with the radius of 1 m, one obtains \mathcal{N}_Z equal to 4×10^{19} . These very large values are due to the mismatch between the value of elementary charge, which appears in the definition of \mathcal{N}_Z , and those values in macroscopic fields.

4. The Zeldovich Number for the Hydrogen Atom

In our calculations of \mathcal{N}_Z associated with the hydrogen atom we will use the ground state electron wave function satisfying the Dirac equation. The wave functions satisfying the nonrelativistic Schrödinger equation would be simpler, but the relativistic theory allows for a uniform treatment of the electric and magnetic fields.

The ground state is doubly degenerate (disregarding tiny corrections due to the hyperfine interactions). The two states differ in the sign of the projection of the total angular momentum on a chosen direction. We choose, as is customary, the z direction. The state with the positive sign has the following normalized Dirac wave function [16],

$$\begin{aligned} \psi(x, y, z, t) &= \sqrt{\frac{\gamma + 1}{8\pi\Gamma(2\gamma + 1)}} \left(\frac{2\alpha}{\lambda}\right)^{3/2} e^{-iEt/\hbar} \\ &\times \left(\frac{2\alpha r}{\lambda}\right)^{\gamma-1} e^{-\alpha r/\lambda} \left\{1, 0, \frac{i\alpha}{\gamma + 1} \frac{z}{r}, \frac{i\alpha}{\gamma + 1} \frac{x + iy}{r}\right\}, \end{aligned} \tag{20}$$

where $\gamma = \sqrt{1 - \alpha^2}$ and $\lambda = \hbar/mc$ is the reduced electron Compton wave length.

The probability density $\rho_e = \psi^*\psi$ and the probability current density $\mathbf{j}_e = \psi^*\boldsymbol{\alpha}\psi$ for the electron are,

$$\rho_e(r) = \frac{e^{-2r/b}(2r/b)^{2\gamma+1}}{4\pi r^3\Gamma(2\gamma+1)}, \tag{21}$$

$$\mathbf{j}_e(r) = \{-y, x, 0\} \frac{\alpha\rho_e(r)}{r}, \tag{22}$$

where $b = \lambda/\alpha$ is the Bohr radius. These sources produce the electromagnetic field \mathcal{D} and \mathcal{H} generated by the electron. Since we are using the rescaled electric and magnetic field, the sources (21) and (22) must be also rescaled. The substitution of the electronic \mathcal{D} into (5) produces the infrared divergence. This divergence has a clear physical interpretation analogous to the one encountered for two charged spheres. The infinite result is just due to the unshielded electron charge. Atoms are neutral and the introduction of the compensating charge of the nucleus will remove the infrared divergence.

The exact formula for the charge distribution in the nucleus is not important. The tiny size of the nucleus as compared to the size of the electronic cloud makes the difference between various models of the nucleus negligible. In our calculations of the electromagnetic field associated with the hydrogen atom we have assumed that the proton charge and the proton magnetic moment are distributed uniformly within a sphere with a sharp boundary (cf. [17]) with the radius a taken from experiment. Thus, the proton charge density and the current density will be described by the Heaviside step function,

$$\rho_p(r) = \frac{3}{4\pi a^3}\Theta(a-r), \tag{23}$$

$$\mathbf{j}_p(r) = \frac{\{y, -x, 0\}}{r} \frac{3\mu}{\pi a^4}\Theta(a-r), \tag{24}$$

where μ measures the strength of the proton magnetic moment μ_p ,

$$\mu = \mu_p/(ec) = 5.8 \times 10^{-16}\text{m}. \tag{25}$$

The densities (23) and (24) satisfy the conditions that the electric and magnetic fields outside the proton are correct,

$$\mathcal{D}_{\text{out}}(\mathbf{r}) = \frac{\mathbf{r}}{4\pi r^3}, \tag{26}$$

$$\mathcal{H}_{\text{out}}(\mathbf{r}) = \frac{\mu}{4\pi r^3} \left(\frac{3\mathbf{r}(\mathbf{r} \cdot \mathbf{n}_z)}{r^2} - \mathbf{n}_z \right), \tag{27}$$

where \mathbf{n}_z is the unit vector along the z axis. We assumed that the proton is at rest. Hence the only contribution to its current comes from the proton magnetic moment μ .

There are three scale parameters in our problem separated by a few orders of magnitude: the proton radius $a = 8.5 \times 10^{-16}\text{m}$, the Compton wave length of the electron $\lambda = 3.86 \times 10^{-13}\text{m}$, and the atomic scale parameter, given by the Bohr radius $b = 5.29 \times 10^{-11}\text{m}$.

The Zeldovich number strongly depends on the proton radius, which is fixed by the experiment. Given the smallness of the proton radius, as compared to λ and b , the eventual changes of the charge distribution inside the proton have almost no influence.

The field vectors $\mathcal{D}(\mathbf{r})$ and $\mathcal{H}(\mathbf{r})$ are the solutions of the Maxwell equations,

$$\nabla \cdot \mathcal{D}(\mathbf{r}) = \rho(r), \quad \nabla \times \mathcal{H}(\mathbf{r}) = \mathbf{j}(r), \tag{28}$$

where

$$\rho(r) = \rho_p(r) - \rho_e(r), \quad \mathbf{j}(r) = \mathbf{j}_p(r) - \mathbf{j}_e(r). \tag{29}$$

Note that the elementary charge e does not appear explicitly in Equations (28) since we are using the field vectors in the geometrical units defined in Equation (2). The terms with time derivatives do not appear in these equations because the sources do not depend on time.

The solutions of the Maxwell equations are most easily obtained with the use of two scalar functions $\phi(r)$ and $\alpha(r)$,

$$\mathcal{D}(\mathbf{r}) = -\nabla\phi(r), \tag{30}$$

$$\mathcal{H}(\mathbf{r}) = \nabla \times \{-y, x, 0\}\alpha(r). \tag{31}$$

After substituting these formulas into the Maxwell Equation (28), one obtains two ordinary differential equations for $\phi(r)$ and $\alpha(r)$,

$$-\phi''(r) - 2/r\phi'(r) = \rho(r), \tag{32}$$

$$-\alpha''(r) - 4/r\alpha'(r) = \chi(r), \tag{33}$$

where $\rho(r)$ and $\chi(r)$ are,

$$\rho(r) = \frac{3}{4\pi a^3}\Theta(a-r) + \rho_e(r), \tag{34}$$

$$\chi(r) = \frac{3\mu}{\pi a^4 r}\Theta(a-r) - \frac{\alpha\rho_e(r)}{r}. \tag{35}$$

The solutions of the Equations (32) and (33) can be expressed as double integrals of the source terms,

$$\phi(r) = \int_r^\infty \frac{dv}{v^2} \int_0^v du u^2 \rho(u), \tag{36}$$

$$\alpha(r) = \int_r^\infty \frac{dv}{v^4} \int_0^v du u^4 \chi(u). \tag{37}$$

These integrals can be evaluated in closed form and the results are:

$$\phi(r) = \frac{1}{4\pi r} \left[\left(\frac{3a^2r - r^3}{2a^3}\Theta(a-r) + \Theta(r-a) \right) - \left(1 - \frac{\Gamma(1+2\gamma, 2r/b) - (2r/b)\Gamma(2\gamma, 2r/b)}{\Gamma(1+2\gamma)} \right) \right], \tag{38}$$

$$\frac{d}{dr}\phi(r) = -\frac{1}{4\pi r^2} \left[\left(\frac{r^3}{a^3}\Theta(a-r) + \Theta(r-a) \right) - \left(1 - \frac{\Gamma(2\gamma+1, 2r/b)}{\Gamma(2\gamma+1)} \right) \right], \tag{39}$$

$$\alpha(r) = \frac{1}{4\pi r^3} \left[\mu \left(\frac{4ar^3 - 3r^4}{a^4}\Theta(a-r) + \Theta(r-a) \right) - \lambda \frac{\Gamma(2\gamma+2, 2r/b) - (2r/b)^3\Gamma(2\gamma-1, 2r/b) - \Gamma(2\gamma-1)}{6\Gamma(2\gamma+1)} \right], \tag{40}$$

$$\frac{d}{dr}\alpha(r) = -\frac{1}{4\pi r^4} \left[3\mu \left(\frac{r^4}{a^4}\Theta(a-r) + \Theta(r-a) \right) - \lambda \frac{\Gamma(2\gamma+2) - \Gamma(2\gamma+2, 2r/b)}{2\Gamma(2\gamma+1)} \right], \tag{41}$$

where $\Gamma(z, a)$ is the incomplete gamma function [18]. The field vectors are constructed from these scalar functions according to the formulas which follow from Equations (30) and (31),

$$\mathcal{D}(\mathbf{r}) = -\frac{\mathbf{r}}{r} \frac{d}{dr}\phi(r), \quad \mathcal{H}(\mathbf{r}) = -\frac{r\mathbf{z}}{r} \frac{d}{dr}\alpha(r) + \mathbf{n}_z \left(r \frac{d}{dr}\alpha(r) + 2\alpha(r) \right), \tag{42}$$

so that,

$$\mathcal{D}(\mathbf{r}) = \frac{\mathbf{r}}{4\pi r^3} \left[\left(\frac{r^3}{a^3} \Theta(a-r) + \Theta(r-a) \right) - \left(1 - \frac{\Gamma(2\gamma+1, 2r/b)}{\Gamma(2\gamma+1)} \right) \right], \tag{43}$$

$$\begin{aligned} \mathcal{H}(\mathbf{r}) &= \frac{z\mathbf{r}}{4\pi r^5} \left[3\mu \left(\frac{r^3}{a^3} \Theta(a-r) + \Theta(r-a) \right) - \lambda \frac{\Gamma(2\gamma+2) - \Gamma(2\gamma+2, 2r/b)}{2\Gamma(2\gamma+1)} \right] \\ &- \frac{n_z}{4\pi r^3} \left[\mu \left(\frac{9r^4 - 8ar^3}{a^4} \Theta(a-r) + \Theta(r-a) \right) - \lambda \frac{\Gamma(2\gamma+2) - \Gamma(2\gamma+2, 2r/b) - 2(2r/b)^3 \Gamma(2\gamma-1, 2r/b)}{6\Gamma(2\gamma+1)} \right]. \end{aligned} \tag{44}$$

Owing to the huge difference in the sizes of the proton and the electron clouds, one cannot show the complete behavior of the field vectors on a single plot. In Figure 1 we show the behavior of \mathcal{D} on the atomic scale and in the inset we show the behavior of \mathcal{D} near $r = 0$ on the scale of the proton radius a . The electric displacement field has the radial form and the magnetic field plotted in Figure 2 exhibits a typical field configuration of a magnet.




Figure 1. The value of the modulus of the electric displacement field $|\mathcal{D}|$ plotted on the atomic scale for the hydrogen atom in the ground state. In order to interpret this plot as the total amount of charge enclosed within the sphere of radius r (Gauss law) is multiplied $|\mathcal{D}|$ by $4\pi r^2$. The details of the behavior of $|\mathcal{D}|$ close to the center are shown in the inset on the scale of the proton radius. Starting from the origin, the enclosed charge increases as $(r/a)^3$ and reaches the value of 1 at the proton radius a .




Figure 2. The magnetic field surrounding the hydrogen atom in the ground state. The plot of the field configuration in the $y = 0$ plane gives the complete information, owing to the rotational symmetry around the z axis. The coordinates x and z are measured in λ .

In order to calculate the Zeldovich number we could use in principle the fields in the position representation and the original Zeldovich Formula (9). However, the use of the Fourier transforms greatly simplifies the calculations. These transforms are obtained from

the same purely algebraic Equations (12) and (18) as in the classical theory. The solutions of these equations are,

$$\tilde{\mathcal{D}}(\mathbf{k}) = -i\mathbf{k} \frac{\tilde{\rho}(k)}{k^2}, \tag{45}$$

$$\tilde{\mathcal{H}}(\mathbf{k}) = i((\mathbf{k} \cdot \mathbf{n}_z)\mathbf{k} - k^2\mathbf{n}_z) \frac{1}{k^3} \frac{d\tilde{\chi}(k)}{dk}. \tag{46}$$

The Fourier transform of spherically symmetric functions become one-dimensional integrals,

$$\tilde{f}(k) = \int \frac{d^3r}{(2\pi)^{3/2}} e^{-i\mathbf{k} \cdot \mathbf{r}} f(r) = \sqrt{\frac{2}{\pi}} \int_0^\infty dr r \frac{\sin(kr)}{k} f(r). \tag{47}$$

This integral can be evaluated in a closed form for both functions $\rho(r)$ and $\chi(r)$,

$$\tilde{\rho}(k) = \frac{3 \sin(ak) - 3ak \cos(ak)}{(2\pi)^{3/2} a^3 k^3} - \frac{\sin[2\gamma \arctan(bk/2)]}{(2\pi)^{3/2} \gamma bk(1 + k^2 b^2/4)^\gamma}, \tag{48}$$

$$\tilde{\chi}(k) = 6\mu \frac{2ak \sin(ak) + (2 - a^2 k^2) \cos(ak) - 2}{(2\pi)^{3/2} a^6 k^4} - \alpha \frac{2 \sin[(2\gamma - 1) \arctan(bk/2)]}{(2\pi)^{3/2} \gamma (2\gamma - 1) b^2 k (1 + b^2 k^2/4)^{\gamma-1/2}}, \tag{49}$$

$$\begin{aligned} \frac{d\tilde{\chi}(k)}{dk} &= 6\mu \frac{ak(a^2 k^2 - 8) \sin(ak) + 4(a^2 k^2 - 2) \cos(ak) + 8}{(2\pi)^{3/2} a^6 k^5} \\ &- \alpha \frac{(2\gamma - 1)bk \cos[2\gamma \arctan(bk/2)] - 2(2 + \gamma b^2 k^2) \sin[2\gamma \arctan(bk/2)]}{(2\pi)^{3/2} 2\gamma (2\gamma - 1) b^2 k^2/4 (1 + b^2 k^2/4)^{\gamma+1/2}}. \end{aligned} \tag{50}$$

The substitution of the expression (45) for $\tilde{\mathcal{D}}$ into Equation (5) gives the formula for the contribution $\mathcal{N}_Z[\mathcal{D}]$, which is associated with the electric field, to the total value,

$$\mathcal{N}_Z[\mathcal{D}] = 2\pi\alpha \int \frac{d^3k}{k^3} \tilde{\rho}(k)^2. \tag{51}$$

The integration over k cannot be done analytically, and one has to resort to numerical integration. The formula for \mathcal{N}_Z with $\tilde{\rho}(k)$ given by Equation (48) contains the dimensional parameters a and b . However, \mathcal{N}_Z is dimensionless so that it can only depend on a dimensionless ratio. Let us make use of this property and change the dimensional integration variable k to the dimensionless variable $\kappa = bk$ and introduce the dimensionless ratio as $s = a/b = 1.6 \times 10^{-5}$. The resulting integral is,

$$\begin{aligned} \mathcal{N}_Z[\mathcal{D}] &= \frac{\alpha}{\pi} \int_0^\infty \frac{d\kappa}{\kappa} \\ &\times \left[\frac{3 \sin(s\kappa) - 3s\kappa \cos(s\kappa)}{s^3 \kappa^3} - \frac{\sin(2\gamma \arctan(\kappa/2))}{\gamma \kappa (1 + \kappa^2/4)^\gamma} \right]^2 \end{aligned} \tag{52}$$

and the “electric Zeldovich number” associated with the hydrogen atom in the ground state is $\mathcal{N}_Z[\mathcal{D}] = 0.025$. By the way, a similar number 0.02 was obtained with the use of very crude arguments in ref. [13].

The “magnetic Zeldovich number” for the hydrogen atom is given by the integral,

$$\mathcal{N}_Z[\mathcal{H}] = 2\pi\alpha \int \frac{d^3k}{k^7} (k^2 - (\mathbf{k} \cdot \mathbf{n}_z)^2)^2 \left(\frac{d\tilde{\chi}(k)}{dk} \right)^2. \tag{53}$$

After the integration over the angles, one obtains,

$$\mathcal{N}_Z[\mathcal{H}] = \frac{16\pi^2\alpha}{3} \int_0^\infty \frac{dk}{k} \left(\frac{d\tilde{\chi}(k)}{dk} \right)^2. \tag{54}$$

Using Equation (49) and replacing $kb = \kappa$ in Equation (50), one obtains,

$$\mathcal{N}_Z[\mathcal{H}] = \frac{2\alpha}{3\pi} \int_0^\infty \frac{d\kappa}{\kappa} \left[6d \frac{s\kappa(s^2\kappa^2 - 8) \sin(s\kappa) + 4(s^2\kappa^2 - 2) \cos(s\kappa) + 8}{s^5\kappa^5} - \alpha \frac{\kappa \cos \sigma - \sin \sigma}{2\gamma(2\gamma - 1)\kappa^2(1 + \kappa^2/4)^\gamma} \right]^2, \tag{55}$$

where $d = \mu/a = 0.68$ and $\sigma = 2\gamma \arctan(\kappa/2)$. Numerical integration produces a tiny number $\mathcal{N}_Z[\mathcal{H}] = 6 \times 10^{-5}$ which is negligible in comparison to the number $\mathcal{N}_Z[\mathcal{D}]$ for the electric field. The reason for this huge difference is a rather slow motion of electrons in comparison to the speed of light. This results in the appearance of the fine structure constant in Equation (22).

We also calculate the energy carried by the electromagnetic field associated with the hydrogen atom. As expected the contribution of the magnetic field is negligible. The energy carried by the electric field comes almost entirely from the Coulomb field of the proton because this field is very strong at small distances. The associated energy is quite substantial; it equals $2m_e c^2$, twice the rest energy of the electron. Of course, the Coulomb energy of the proton cannot be counted as a separate contribution because it is already included in the observed proton rest energy. The total field energy associated with the electronic wave function is very tiny. This energy is also dominated by the electric part and it is equal to $1.67 \times 10^{-5} m_e c^2$.

5. The Zeldovich Number for Heavier Atoms

As an example, let us choose the atoms of noble gases because their closed shells produce spherically symmetric charge distribution which greatly simplifies the calculations. We restrict ourselves here to the calculation of the “electric Zeldovich number” since, as was seen in the case of the hydrogen atom, the “magnetic Zeldovich number” is much smaller. This number grows rapidly with the increase of the atomic number Z . The calculation of the exact value of the average photon number requires the knowledge of the total wave function of mutually interacting electrons, but to obtain an order of magnitude estimate one can neglect these interactions. We also use the electron wave functions in the nonrelativistic approximation,

$$\psi_{nlm}(\varrho, \theta, \phi) = \sqrt{\frac{(n-l-1)!}{2n(n+l)!}} \left(\frac{2Z}{nb} \right)^{3/2} \varrho^l e^{-\varrho/2} L_{n-l-1}^{2l+1}(\varrho) Y_l^m(\theta, \phi), \tag{56}$$

where $\varrho = \frac{2Zr}{nb}$. Note that we use a different font to distinguish the rescaled radial variable ϱ from the probability density ρ . In most textbooks on quantum mechanics the lower index of the associated Laguerre polynomial $L_{n-l-1}^{2l+1}(\varrho)$ has a different meaning; instead of $n-l-1$ it is equal to $n+l$. We follow here the notation of *Mathematica* [19]. The probability density corresponding to the wave function (56) is,

$$\rho_{nlm}(\varrho, \theta, \phi) = \frac{(n-l-1)!}{2n(n+l)!} \left(\frac{2Z}{nb} \right)^3 \varrho^{2l} e^{-\varrho} \left[L_{n-l-1}^{2l+1}(\varrho) \right]^2 Y_l^m(\theta, \phi) Y_l^m(\theta, -\phi), \tag{57}$$

The only dependence on the quantum number m is through the spherical harmonics. Therefore, for given n and l one can sum up $2(2l+1)$ contributions from different values of m and obtain the total contribution ρ_{nl} , from the fully filled (n, l) -shell. The additional

factor 2 is due to the two possible electron spin orientations. The density $\rho_{nl}(\varrho)$ of the electrons for the (n, l) shell is,

$$\rho_{nl}(\varrho) = \frac{2(2l + 1)}{4\pi} \frac{(n - l - 1)!}{2n(n + l)!} \left(\frac{2Z}{nb}\right)^3 \varrho^{2l} e^{-\varrho} \left[L_{n-l-1}^{2l+1}(\varrho)\right]^2. \tag{58}$$

The total charge density must also include the contribution $\rho_{nucl}(\varrho)$ from the nucleus. We assume, as was done for the proton in Equation (23), that the charge is distributed evenly within the sphere whose radius depends on the atomic mass number A . Then,

$$\rho_{nucl}(\varrho) = \frac{3Z}{4\pi a(A)^3} \Theta(a(A) - r). \tag{59}$$

We use the commonly accepted (see, e.g., [20]) formula for $a(A)$,

$$a(A) = 1.2 A^{1/3} \times 10^{-15} \text{ m}. \tag{60}$$

In order to determine the Fourier transform of the electric field $\hat{D}(\mathbf{k})$ one needs the Fourier transform of the electron charge density for all shells and the nucleus charge density. The calculations are simple but tedious. In order to obtain the results for all stable noble gas atoms we calculate, according to Equation (47), the Fourier transforms of the functions $\rho_{nl}(r)$ for all closed shells. All the integrals are evaluated analytically and they are equal to the ratios of the polynomials in k^2 . Finally, for each atom we add up the contributions from the relevant shells and evaluate the final value (51) of the Zeldovich number.

The results are shown in Figure 3. As expected, the dependence on Z is almost quadratic (dashed line). The deviation is due to the increase of the nucleus radius with the increasing atomic mass number A . This effect diminishes the strength of the electric field at the center of the atom.




Figure 3. The Zeldovich number for the atoms of noble gases. The dashed line shows the expected approximate quadratic dependence.

6. Conclusions

The results presented in this work have probably no immediate applications. They offer, however, a fresh point of view by introducing a new universal dimensionless measure of the overall strength of the electromagnetic field, the same in the classical and in the quantum domain. For a free quantized electromagnetic field, this measure gives the number of photons. The Zeldovich number can also be viewed as a measure of the strength of the sources of the electromagnetic field. There is one property of N_Z which is worth stressing: it is an intensive and not an extensive property of the system. It does not depend on the size of the system but only on the *dimensionless ratios* of various parameters to the overall system size.

Author Contributions: The authors contributed equally to this manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Zeldovich, Y.B. Number of quanta as an invariant of the classical electromagnetic field. *Sov. Phys. Doklady* **1966**, *10*, 771–772.
2. Passante, R.; Compagno, G.; Persico, F. Cloud of virtual photons in the ground state of the hydrogen atom. *Phys. Rev. A* **1985**, *31*, 2827–2841. [[CrossRef](#)] [[PubMed](#)]
3. Persico, F.; Compagno, G.; Passante, R. Ground state photons dressing atoms and molecules. In *Quantum Optics IV*; Harvey, J.D., Walls, D.F., Eds.; Springer: Berlin/Heidelberg, Germany, 1986; pp. 172–180. [[CrossRef](#)]
4. Compagno, G.; Passante, R.; Persico, F.; Salamone, G.M. Cloud of virtual photons surrounding a nonrelativistic electron. *Acta Phys. Pol. A* **1994**, *85*, 667–676. [[CrossRef](#)]
5. Compagno, G.; Palma, G.M.; Passante, R.; Persico, F. Atoms dressed and partially dressed by the zero-point fluctuations of the electromagnetic field. *J. Phys. B* **1995**, *8*, 1105–1158. [[CrossRef](#)]
6. Radozycki, T. The electromagnetic virtual cloud of the ground-state hydrogen atom—A quantum field theory approach. *J. Phys. A* **1990**, *23*, 4911–4923. [[CrossRef](#)]
7. Gross, L. Norm invariance of mass-zero equations under the conformal group. *J. Math. Phys.* **1964**, *5*, 687–695. [[CrossRef](#)]
8. Crispino, L.C.B.; Higuchi, A.; Matsas, G.E.A. The Unruh effect and its applications. *Rev. Mod. Phys.* **2008**, *80*, 787–838. [[CrossRef](#)]
9. Białynicki-Birula, I. Photon wave function. *Prog. Opt.* **1996**, *36*, 245–294. [[CrossRef](#)]
10. Białynicki-Birula, I.; Białynicka-Birula, Z. The role of the Riemann-Silberstein vector in classical and quantum theories of electromagnetism. *J. Phys. A Math. Theor.* **2013**, *46*, 053001. [[CrossRef](#)]
11. Białynicki-Birula, I.; Białynicka-Birula, Z. Three measures of fidelity for photon states. *Phys. Rev. A* **2020**, *102*, 042201. [[CrossRef](#)]
12. Białynicki-Birula, I. The Wigner functional of the electromagnetic field. *Opt. Commun.* **2000**, *179*, 237–246. [[CrossRef](#)]
13. Białynicki-Birula, I. The structure of the vacuum and the photon number. In *Decoherence and Entropy in Complex Systems*; Elze, H.T., Ed.; Springer: Berlin/Heidelberg, Germany, 2004; pp. 287–295. [[CrossRef](#)]
14. Białynicki-Birula, I. Relativistic Wigner functions. *EPJ Web Conf.* **2014**, *78*, 01001. [[CrossRef](#)]
15. Jackson, J.D. *Classical Electrodynamics*; John Wiley & Sons, Inc.: Hoboken, NJ, USA, 1999; p. 181. Available online: <https://www.scribd.com/doc/188342140/J-D-Jackson-Classical-Electrodynamics> (accessed on 10 July 2023).
16. Bjorken, J.D.; Drell, S.D. *Relativistic Quantum Mechanics*; McGraw-Hill Book Company: New York, NY, USA, 1964; p. 55. Available online: <https://emineter.files.wordpress.com/2018/10/james-d-bjorken-sidney-d-drell-relativistic-quantum-mechanics-1964.pdf> (accessed on 10 July 2023).
17. Marchildon, L. *Quantum Mechanics: From Basic Principles to Numerical Methods and Applications*; Springer: Berlin/Heidelberg, Germany, 2002; p. 211. [[CrossRef](#)]
18. Olver, F.W.J.; Lozier, D.W.; Boisvert, R.F.; Clark, C.W. (Eds.) *NIST Handbook of Mathematical Functions*; NIST: Gaithersburg, MD, USA; Cambridge University Press: New York, NY, USA, 2010; p. 174. Available online: <https://univ.jeanpaulcalvi.com/Posters/ConfAuchWeb/abramovitz2.pdf> (accessed on 10 July 2023).
19. Wolfram, S. *Mathematica*, Version 13.1; Wolfram Research, Inc.: Champaign, IL, USA, 2022. Available online: <https://www.wolfram.com/mathematica/> (accessed on 10 July 2023).
20. Basdevant, J.L.; Rich, J.; Spiro, M. *Fundamentals in Nuclear Physics: From Nuclear Structure to Cosmology*; Springer: New York, NY, USA, 2005; p. 151. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.