

Supporting information

Exploration of the Divergent Outcomes for the Nenitzescu Reaction of Piperazinone Enaminoesters

Rebecca Hermans ¹, Max Van Hoof ¹, Luc Van Meervelt ² and Wim Dehaen ^{1,*}



¹ Sustainable Chemistry for Metals and Molecules, Department of Chemistry, KU Leuven, Celestijnenlaan 200F, B-3001 Leuven, Belgium

² Biochemistry, Molecular and Structural Biology, Department of Chemistry, KU Leuven, Celestijnenlaan 200F, B-3001 Leuven, Belgium



* Correspondence: wim.dehaen@kuleuven.be

NMR Spectra



Ethyl (Z)-2-(3-oxopiperazin-2-ylidene)acetate (15)




Methyl (Z)-2-(3-oxopiperazin-2-ylidene)acetate (22)




Ethyl (Z)-2-(5-methyl-3-oxopiperazin-2-ylidene)acetate (23)





Ethyl (Z)-2-(3-oxo-3,4-dihydroquinoxalin-2(1*H*)-ylidene)acetate (24)







Ethyl (Z)-2-((5*S*,6*S*)-3-oxo-5,6-diphenylpiperazin-2-ylidene)acetate (25)





Ethyl 5-hydroxy-7-methyl-2-(2-oxoimidazolidin-1-yl)benzofuran-3-carboxylate (19)








**Ethyl-2-(5-methyl-3,6-dioxocyclohexa-1,4-dien-1-yl)-2-(3-oxopiperazin-2-ylidene)acetate
(20a)**




**Ethyl-2-(4-methyl-3,6-dioxocyclohexa-1,4-dien-1-yl)-2-(3-oxopiperazin-2-ylidene)acetate
(20b)**





Ethyl 8-hydroxy-1-oxo-1,2,3,4-tetrahydropyrazino[1,2-a]indole-10-carboxylate (27)










Ethyl 8-hydroxy-6-methyl-1-oxo-1,2,3,4-tetrahydropyrazino[1,2-a]indole-10-carboxylate (18b) and ethyl 8-hydroxy-7-methyl-1-oxo-1,2,3,4-tetrahydropyrazino[1,2-a]indole-10-carboxylate (18a)





(Z)-3-(5-Hydroxy-7-methyl-2-oxobenzofuran-3(2H)-ylidene)piperazin-2-one (21a)





NOESY

