



Proceeding Paper Cytotoxicity Studies of Eugenol Amino Alcohols Derivatives [†]

Cláudia Teixeira ^{1,2}, Nuno F. S. Pinto ¹, David M. Pereira ^{2,*}, Renato B. Pereira ², Maria José G. Fernandes ¹, Elisabete M. S. Castanheira ³, António Gil Fortes ¹ and Maria Sameiro T. Gonçalves ¹

- ¹ Centre of Chemistry (CQ/UM), University of Minho, Campus of Gualtar, 4710-057 Braga, Portugal; pg40748@alunos.uminho.pt (C.T.); nuno_pinto1993@hotmail.com (N.F.S.P.); mjfernandes@quimica.uminho.pt (M.J.G.F.); gilf@quimica.uminho.pt (A.G.F.); msameiro@quimica.uminho.pt (M.S.T.G.)
- ² REQUIMTE/LAQV, Laboratory of Pharmacognosy, Department of Chemistry, Faculty of Pharmacy, University of Porto, R. Jorge Viterbo Ferreira, 228, 4050-313 Porto, Portugal; ren.pereira@gmail.com
- ³ Centre of Physics of Minho and Porto Universities (CF-UM-UP), University of Minho, Campus of Gualtar, 4710-057 Braga, Portugal; ecoutinho@fisica.uminho.pt
- * Correspondence: dpereira@ff.up.pt
- + Presented at the 25th International Electronic Conference on Synthetic Organic Chemistry, 15–30 November 2021; Available online: https://ecsoc-25.sciforum.net/.

Abstract: Eugenol is a phenylpropanoid displaying a wide range of biological activities. In this study, the cytotoxic activity of various β -amino alcohols derivatives from eugenol was evaluated in AGS (gastric cancer) and A549 (lung cancer) cell lines. The results show that some structural modifications resulted in enhanced cytotoxic activity towards cancer cells. In addition, the activation of caspase-3 and hence apoptosis induced by these molecules, was also explored. Considering the obtained results, some structure/activity relationships can be drawn, which may guide future structural improvements for anticancer agents.

Keywords: eugenol; β-amino alcohols; essential oils; cytotoxicity; caspase-3; anticancer



Citation: Teixeira, C.; Pinto, N.F.S.; Pereira, D.M.; Pereira, R.B.; Fernandes, M.J.G.; Castanheira, E.M.S.; Fortes, A.G.; Gonçalves, M.S.T. Cytotoxicity Studies of Eugenol Amino Alcohols Derivatives. *Chem. Proc.* 2022, *8*, 105. https:// doi.org/10.3390/ecsoc-25-11689

Academic Editor: Julio A. Seijas

Published: 14 November 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/).

1. Introduction

Humans have relied on plants for treating a wide range of diseases since pre-historical times [1,2]. Drug discovery and design continues to be inspired by natural products, with 23.5% of 1881 new drugs, approved between 1981 and 2019, being unaltered natural products, botanical drugs, or natural derivatives [3].

Essential oils, natural matrices comprising several secondary metabolites of low molecular weight, gained interest due to their biological activities and application across several industries. Eugenol, the major constituent of clove oil, displays numerous biological properties such as anti-inflammatory, antioxidant, anti-tumorigenic, anti-microbial and cytotoxic activity [4,5]. Eugenol also has an interesting structure to be used as a starting molecule to obtain several synthetic derivatives. For these reasons several studies on eugenol or its synthetic derivatives have been conducted [6].

Cancer is a multistep process that involves dynamic changes in the genome and allows cells to invade several tissues, increasing cell proliferation and escape from programmed cell death processes, such as apoptosis [7,8]. Apoptosis is a type of cell death characterized by specific morphological and biochemical changes and requires the activation of caspases, a group of cysteine-aspartic acid proteases. It can be triggered by two pathways: the intrinsic/mitochondrial pathway and the extrinsic/death receptor pathway [9–12]. The ability to escape from apoptosis is a hallmark of cancer and, for this reason, new therapies that reactivate apoptotic mechanisms hold great promise to counteract cancer [8,9]. It is to be noted that eugenol has already been shown to induce apoptosis in different cancer cell lines [13].

Considering all these facts, cytotoxicity against AGS (human gastric adenocarcinoma) and A549 (human lung adenocarcinoma) cell lines of was screened with several β -amino alcohols from eugenol. The most potent molecules were selected and evaluated for their capacity to induce apoptosis.

2. Results and Discussion

2.1. Synthesis of Eugenol β-Amino Alcohols Derivatives 3–9

Eugenol **1** was reacted with *m*-chloroperbenzoic acid in dichloromethane to give the epoxide **2** [14], which was further reacted with several aromatic and aliphatic amines in ethanol/water as solvent by using a known procedure [15], to yield the corresponding β -amino alcohol derivatives **3–9** [16] (Figure 1).



Figure 1. Structures of eugenol 1, eugenol epoxide 2 and the corresponding β -amino alcohols 3–9.

2.2. Toxicity of β -Amino Alcohols Eugenol Derivatives **3–9**

The effect of β -amino alcohols 3–9 as well as precursor molecules 1 and 2 were studied in AGS and A549 cell lines at 24 h (100 μ M) (Figure 2).



Figure 2. Effect of eugenol and its derivatives (100 μ M) in (a) AGS and (b) A549 cell lines. The results correspond to the mean of each well of three independent experiments performed in triplicate. *** *p* < 0.001.

When evaluating the impact of all the molecules obtained, it was clear that for both cell lines, a few derivatives displayed higher cytotoxicity than eugenol **1** and its epoxide **2**, which was devoid of activity. Among all derivatives, compound **5** was the most potent for AGS cells, leading to 28% reduction in cell viability and for A549 cells, compound **4** reduced cell viability by 31%.

Considering these results, we were interested in understanding if a process of programmed cell death, such as apoptosis, was taking place. After the results from the viability assay, the most potent molecule for each cell line was tested for possible triggering of cell death by apoptosis. For this purpose, the activation of the effector caspase-3 was studied (Figure 3).



Figure 3. Effect of the compounds **5** and **4** at 100 μ M on the activity of caspase-3, on (**a**) AGS cell line and (**b**) A549 cell line. The results correspond to the mean of each well of four independent experiments performed in duplicate. The values were normalized for DNA content. *** *p* < 0.001.

This experiment revealed that for the AGS cell line, compound **5** at 100 μ M elicited an increase in caspase-3 activity of about 12-fold. Compound **4** was also active, while in much lower magnitude.

In addition, compound **5** had a stronger effect in caspase-3 activation allowing us to conclude that cell viability reduction is caused by activation of apoptosis.

3. Experimental

3.1. Cell Culture and Viability Assessment

AGS (human gastric adenocarcinoma) and A549 (human lung adenocarcinoma) cells are broadly used in pharmacological studies and create a versatile pair, due to the different ways in which the cells to respond to chemotherapy treatments, with AGS being more responsive than A549.

Cells were maintained in DMEM plus GlutaMAXTM with 10% FBS and 1% penicillin/streptomycin, at 37 °C with 5% CO₂ in a humidified incubator. For the assessment of viability, a resazurin-based method was used. AGS and A549 cells were seeded at a density of 15,000 and 10,000 cells/well, respectively, incubated for 24 h, and then exposed to the compounds under study for another 24 h period. After this period, a commercial solution of resazurin was added (1:10) and incubated for 30 min. Finally, the fluorescence was read at 560 nm.

3.2. Caspase Activity

AGS and A549 cells were seeded at the same density described for viability assessment and exposed to the compounds under study for the same period of time. After the incubation period, 50 μ L of supernatant was removed from each well, followed by the addition of 50 μ L of caspase-3 substrate (1:200), and the plate was incubated for 45 min. Lastly, fluorexscent was read at 535–620 nm.

3.3. DNA Quantification

Cells were cultured and exposed to the molecules under study, as described above. After the incubation period, the culture medium was replaced by 50 μ L of ultra-pure water and then the plate was incubated for 30 min in a humidified incubator at 37 °C with 5% CO₂ and subsequently frozen at -80 °C. DNA quantification was performed by using *QubitTM* dsDNA HS/Protein assay kit, the *QubitTM* 4 Fluorometer reader was used.

4. Conclusions

The cytotoxic activity of various β -amino alcohols derivatives from eugenol evaluated in AGS and A549 cell lines revealed that compounds **4** and **5** are the most promising at 100 μ M. These molecules enhanced the cytotoxic activity in relation to eugenol. In addition, compound **5** was able to increase caspase-3 activity, which is involved in the apoptosis cell death process. Therefore, compound **5** could be used as an anticancer agent and as a starting molecule for the design of new and more potent anticancer molecules.

Author Contributions: Conceptualization, D.M.P., A.G.F. and M.S.T.G.; methodology, D.M.P., A.G.F. and M.S.T.G.; formal analysis, D.M.P., R.B.P., A.G.F. and M.S.T.G.; investigation, C.T. and N.F.S.P.; writing—original draft preparation, C.T. and M.S.T.G. writing, reviewing and editing, D.M.P., M.S.T.G., M.J.G.F., R.B.P. and E.M.S.C.; supervision, D.M.P., R.B.P. and A.G.F.; project administration, M.S.T.G. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by FCT under project PTDC/ASP-AGR/30154/2017 (PO-CI-01-0145-FEDER-030154) of COMPETE 2020, co-financed by FEDER and EU. FCT-Portugal and FEDER-COMPETE/QREN-EU also gave financial support to the research centres CQ/UM (UIDB/00686/2020), CF-UM-UP (UIDB/04650/2020) and REQUIMTE (UIDB/50006/2020). The NMR spectrometer Bruker Avance III 400 (part of the National NMR Network) was financed by and FEDER.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Cragg, G.M.; Newman, D.J. Nature: A vital source of leads for anticancer drug development. *Phytochem. Rev.* **2009**, *8*, 313–331. [CrossRef]
- Sen, T.; Samanta, S.K. Medicinal plants, human health and biodiversity: A broad review. *Adv. Biochem. Eng. Biotechnol.* 2015, 147, 59–110. [PubMed]
- 3. Newman, D.J.; Cragg, G.M. Natural products as sources of new drugs over the nearly four decades from 01/1981 to 09/2019. *J. Nat. Prod.* 2020, *83*, 770–803. [CrossRef] [PubMed]
- Blowman, K.; Magalhaes, M.; Lemos, M.F.L.; Cabral, C.; Pires, I.M. Anticancer properties of essential oils and other natural products. *Evid. Based Complement. Altern. Med.* 2018, 2018, 3149362. [CrossRef] [PubMed]
- Nehme, R.; Andrés, S.; Pereira, R.B.; Jemaa, M.B.; Bouhallab, S.; Ceciliani, F.; López, S.; Rahali, F.Z.; Ksouri, R.; Pereira, D.M.; et al. Essential oils in livestock: From health to food quality. *Antioxidants* 2021, 10, 330. [CrossRef] [PubMed]
- Kaufman, T.S. The multiple faces of eugenol. A versatile starting material and building block for organic and bio-organic synthesis and a convenient precursor toward bio-based fine chemicals. J. Braz. Chem. Soc. 2015, 26, 1055–1086. [CrossRef]
- 7. Hanahan, D.; Weinberg, R.A. Hallmarks of cancer: The next generation. Cell 2011, 144, 646–674. [CrossRef] [PubMed]
- 8. Hanahan, D.; Weinberg, R.A. The hallmarks of cancer. Cell 2000, 100, 57–70. [CrossRef]
- Pistritto, G.; Trisciuoglio, D.; Ceci, C.; Garufi, A.; D'Orazi, G. Apoptosis as anticancer mechanism: Function and dysfunction of its modulators and targeted therapeutic strategies. *Aging* 2016, *8*, 603–619. [CrossRef] [PubMed]
- 10. Elmore, S. Apoptosis: A review of programmed cell death. Toxicol. Pathol. 2007, 35, 495–516. [CrossRef] [PubMed]
- 11. Ouyang, L.; Shi, Z.; Zhao, S.; Wang, F.T.; Zhou, T.T.; Liu, B.; Bao, J.K. Programmed cell death pathways in cancer: A review of apoptosis, autophagy and programmed necrosis. *Cell Prolif.* **2012**, *45*, 487–498. [CrossRef] [PubMed]
- 12. D'Arcy, M.S. Cell death: A review of the major forms of apoptosis, necrosis and autophagy. *Cell Biol. Int.* **2019**, *43*, 582–592. [CrossRef] [PubMed]
- Jaganathan, S.K.; Supriyanto, E. Antiproliferative and molecular mechanism of eugenol-induced apoptosis in cancer cells. Molecules 2012, 17, 6290–6304. [CrossRef] [PubMed]
- 14. Silva, F.F.M.; Monte, F.J.Q.; Lemos, T.L.G.; Nascimento, P.G.G.; Costa, A.K.M.; Paiva, L.M.M. Eugenol derivatives: Synthesis, characterization, and evaluation of antibacterial and antioxidant activities. *Chem. Cent. J.* **2018**, *12*, 1–9. [CrossRef] [PubMed]
- 15. Azizi, N.; Saidi, M.R. Highly chemoselective addition of amines to epoxides in water. *Org. Lett.* **2015**, *7*, 3649–3651. [CrossRef] [PubMed]
- Pinto, N.F.S.; Fernandes, M.J.G.; Pereira, R.B.; Vieira, T.F.; Rodrigues, A.R.O.; Pereira, D.M.; Sousa, S.F.; Castanheira, E.M.S.; Fortes, A.G.; Gonçalves, M.S.T. Amino alcohols from eugenol as potential semisynthetic insecticides: Chemical, biological and computational insights. *Molecules* 2021, 26, 6616.