

Proceeding Paper

# Azide-Alkyne Cycloaddition Catalyzed by a Glucose/Benedict Reagent System<sup>†</sup>

Jesús G. García Torres<sup>1</sup>, Andrés F. Becerra-Buitrago<sup>2</sup>, Luis Carlos García-Sánchez<sup>2</sup>, Marco A. García-Eleno<sup>1,3</sup>, M. V. Basavanag Unnamatla<sup>1,3</sup>  and Erick Cuevas-Yañez<sup>1,3,\*</sup> 

<sup>1</sup> Centro Conjunto de Investigación en Química Sustentable UAEM-UNAM, Carretera Toluca-Atlaconulco Km. 14.5, Toluca 50200, Estado de México, Mexico; [jesus.alfa.aquilae@gmail.com](mailto:jesus.alfa.aquilae@gmail.com) (J.G.G.T.); [magciae@uaemex.mx](mailto:magciae@uaemex.mx) (M.A.G.-E.); [mvbasavanagu@uaemex.mx](mailto:mvbasavanagu@uaemex.mx) (M.V.B.U.)

<sup>2</sup> Facultad de Ciencias y Educación, Universidad Distrital Francisco José de Caldas, Carrera 7 No. 40B-53, Bogotá D.C. 11021-110231588, Colombia; [andresbecerrab@gmail.com](mailto:andresbecerrab@gmail.com) (A.F.B.-B.); [lcgarcia@udistrital.edu.co](mailto:lcgarcia@udistrital.edu.co) (L.C.G.-S.)

<sup>3</sup> Facultad de Química, Universidad Autónoma del Estado de México, Paseo Colón esq. Paseo Tollocan, Toluca 50120, Estado de México, Mexico

\* Correspondence: [ecuevasy@uaemex.mx](mailto:ecuevasy@uaemex.mx); Tel.: +52-722-276-6610 (ext. 7734)

† Presented at the 25th International Electronic Conference on Synthetic Organic Chemistry, 1–30 November 2021; Available online: <https://ecsoc-25.sciforum.net/>.

**Abstract:** Benedict's Reagent is commonly used in identifying reducing sugars through a redox process where a Cu(I) species is generated. Despite the simplicity of this reaction, this has barely been investigated as a copper (I) source for catalytic processes. In this report, diverse organic azides and alkynes were reacted in the presence of catalytic amounts of a Glucose–Benedict Reagent system, obtaining the corresponding 1,2,3-triazoles through a simple and environmentally friendly synthetic procedure.

**Keywords:** click chemistry; Benedict reagent; glucose; alkyne; azide; 1,2,3-triazole



**Citation:** Torres, J.G.G.;

Becerra-Buitrago, A.F.;

García-Sánchez, L.C.; García-Eleno,

M.A.; Unnamatla, M.V.B.;

Cuevas-Yañez, E. Azide-Alkyne

Cycloaddition Catalyzed by a  
Glucose/Benedict Reagent System.

*Chem. Proc.* **2022**, *8*, 64. <https://doi.org/10.3390/ecsoc-25-11634>

Academic Editor: Julio A. Seijas

Published: 12 November 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The notion of Click Chemistry was initially conceived by Sharpless 20 years ago [1], and shortly thereafter this group, together with the Meldal group [2], developed the copper-catalyzed azide-alkyne cycloaddition (CuAAC), the first click reaction. This was the departure point for an amazingly rapid growth of the CuAAC reaction, including its applications in diverse fields such as materials science, catalysis, or medicinal chemistry [3–6].


Since its discovery, the copper catalyst has represented a fundamental part of the CuAAC reaction. Therefore, a considerable number of copper-based catalytic systems have been devised to catalyze this reaction. A widely used copper catalytic system is based on the bio-reduction of a copper(II) salt. Seminal reports describe the use of sodium ascorbate as a reducing agent to provide a copper(I) catalyst from copper(II) sulfate which efficiently catalyzes the CuAAC reaction using a mixture of tBuOH–H<sub>2</sub>O as solvent [7].

From these original conditions which served as a model, many adaptations were made to prepare alternative copper catalytic systems. For instance, the well-known Fehling reagent was reduced by hydrazine to give in situ a copper(I) system which catalyzed CuAAC reaction, yielding the corresponding 1,2,3-triazoles [8]. In this regard, our group observed that reducing sugars can also be used as reducing agents in combination with the Fehling reagent to generate a Cu(I) catalyst for the synthesis of both 1,2,3-triazoles and bi-1,2,3-triazoles through the Click Chemistry approach [9,10]. On the other hand, an analogue Benedict reagent was reduced by ascorbic acid, affording a Cu<sub>2</sub>O catalyst that was used as a catalytic system in the preparation of a series of 1,2,3-triazoles [11].

This background prompted us to propose that a combination of Benedict reagent and a reducing sugar would provide an efficient copper(I) source to catalyze the CuAAC reaction. In this report, we disclose our most recent findings in this area.

## 2. Results and Discussion

Preliminary studies were carried out to find the optimal reaction conditions, and glucose was selected as the reducing sugar in these studies. An initial mixture of Benedict reagent, glucose, 1-Azido-4-bromobenzene **1** and 1-chloro-4-prop-2-ynoxybenzene **2** according to Scheme 1 and Table 1 remained unreactive at room temperature, presumably due to Cu(I) species not being formed under these conditions, which was evident by the blue-colored solution in the reaction mixture indicating the presence of Cu(II) ion.




**Scheme 1.** Synthesis of 1,2,3-triazole **3** catalyzed by Glucose/Benedict Reagent System.

**Table 1.** Synthesis of triazole **3** catalyzed by Glucose/Benedict Reagent System.

Entry	Catalyst Ratio (% mmol)	Temperature (°C)	Reaction Time (h)	%Yield
1	2.5	Room Temperature (R.T.)	24	0
2	5	R.T.	24	0
3	10	R.T.	24	0
4	2.5	reflux	24	30
5	5	reflux	24	44
6	10	reflux	24	53
7	2.5	50	24	67
8	5	50	24	67
9	10	50	24	68
10	2.5	50 °C to R.T.	12	32
11	5	50 °C to R.T.	12	49
12	10	50 °C to R.T.	12	51
13	2.5	50 °C to R.T.	24	70
14	5	50 °C to R.T.	24	73
15	10	50 °C to R.T.	24	74
16	2.5	50 °C to R.T.	48	71
17	5	50 °C to R.T.	48	74
18	10	50 °C to R.T.	48	75

This fact drove us to perform the following experiments at higher temperatures. Reactions heated to reflux temperature gave triazole **3** together with traces of other side products identified as bis-acetylenes. In order to avoid these kinds of compounds, an adaptation was accomplished. A premixed glucose-Benedict reagent solution was heated to 50 °C for 5 min and cooled to room temperature, and the resulting red Cu<sub>2</sub>O suspension was treated successively with azide **1** and alkyne **2**, obtaining the corresponding triazole **3** with a yield of up to 73%.

The success of this process promoted an in-depth study to determine the scope of this reaction. Thus, an array of organic azides and alkynes were reacted in the presence of catalytic amounts of glucose/Benedict reagent system, giving the triazoles **3-11** in yields ranging from 40 to 73% (Scheme 2).



**Scheme 2.** Synthesized 1,2,3-triazoles catalyzed by Glucose/Benedict Reagent System.

An outstanding feature of this procedure is related to the high selectivity displayed, due to only 1,2,3-triazole being obtained, whereas the bi-1,2,3-triazole formation was not detected, unlike in the use of the Fehling-glucose catalytic system which promotes the generation of these compounds [9,10]. An explanation for this behavior might be found in the strong dependence on the alkalinity of the solution often observed in these processes. Previous reports emphasize that an excess of NaOH contributes to the formation of bi-1,2,3-triazole [12]. In this case, the pH is controlled by the concentration of less basic  $\text{Na}_2\text{CO}_3$  and the Cu (I) ion is stabilized by the citrate ligand. These conditions provide a convenient source of Cu (I) ion but also avoid the triazolide oxidative coupling to bi-1,2,3-triazole. Hence, the glucose/Benedict reagent system exhibits a catalytic activity that is useful for the synthesis of 1,2,3-triazoles.

As far as we know, these are the first examples of the use of the glucose/Benedict reagent system as a copper source for the CuAAC reaction, which represents an alternative catalytic system for the synthesis of 1,2,3-triazoles through a simple and environmentally friendly methodology.

### 3. Experimental Section

The starting materials were purchased from Aldrich Chemical Co. and were used without further purification. Benedict's reagent was prepared by mixing sodium citrate (17.3 g),  $\text{Na}_2\text{CO}_3$  (5.0 g) and  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  (17.3 g) in distilled  $\text{H}_2\text{O}$  (50 mL). The solvents were distilled before use. Silica plates of 0.20 mm thickness were used for thin layer chromatography. Melting points were determined with a Krüss Optronic melting point apparatus, and they were uncorrected.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded using a Bruker Avance 300-MHz; the chemical shifts ( $\delta$ ) are given in ppm relative to TMS as an internal standard (0.00). For analytical purposes, the mass spectra were recorded on a Shimadzu GCMS-QP2010 Plus in the EI mode, 70 eV, and 200 °C via direct inlet probe.

Only the molecular and parent ions ( $m/z$ ) are reported. IR spectra were recorded on a Bruker Tensor 27.

### 3.1. General Procedure for the Synthesis of 1,2,3-Triazoles Catalyzed by Glucose/Benedict Reagent System

Glucose (0.045 g, 0.25 mmol) was added to a solution of Benedict reagent (0.75 mL, 0.005 mmol) in MeOH (10 mL) and H<sub>2</sub>O (4 mL). The resulting mixture was heated at 50 °C for 5 min and cooled to room temperature. The mixture was treated successively with a solution of azide (1 mmol) in MeOH (2 mL) and alkyne (1 mmol) in MeOH (2 mL). The reaction mixture was stirred for 24 h at room temperature. A 2% EDTA.2Na aqueous solution (10 mL) was added, and the stirring was continued for additional 12 h. Charcoal (0.5 g) was added, the mixture was filtered through celite, and the solvent was removed under reduced pressure. The final product was purified by crystallization.

#### 3.1.1. 1-(4-Bromophenyl)-4-(4-chlorophenoxymethyl)-1,2,3-triazole 3

1-Azido-4-bromobenzene **1** and 1-chloro-4-prop-2-ynoxybenzene **2** afforded 1-(4-Bromophenyl)-4-(4-chlorophenoxymethyl)-1,2,3-triazole **3** as a white solid, m.p. 112 °C (73%). IR (ATR)  $\nu_{\text{max}}$  3140, 3102, 2925, 2859, 1485, 823, 799 cm<sup>-1</sup>. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  8.03 (s, 1H), 7.68 (m, 4H), 7.27 (d, 2H), 6.96 (d, 2H), 5.26 (s, 2H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  156.7, 144.9, 135.9, 133.0, 129.5, 126.4, 122.63, 122.0, 120.8, 116.1, 62.16. MS [EI+]  $m/z$  (%): 363 [M]<sup>+</sup> (5), 141 [C<sub>7</sub>H<sub>6</sub>Cl]<sup>+</sup> (100).

#### 3.1.2. 4-(4-Chlorophenoxymethyl)-1-phenyl-1,2,3-triazole 4

Phenyl azide and 1-chloro-4-prop-2-ynoxybenzene **2** afforded 4-(4-Chlorophenoxymethyl)-1-phenyl-1,2,3-triazole **4**, m.p. 97 °C (55%). IR (ATR)  $\nu_{\text{max}}$  3050, 1500, 1250, 825 cm<sup>-1</sup>. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  8.05 (s, 1H), 7.63 (m, 2H), 7.49 (m, 2H), 7.26 (m, 2H), 6.97 (m, 2H), 5.28 (s, 2H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  155.6, 142.1, 133.5, 129.7, 128.2, 126.8, 123.8, 119.4, 115.0, 74.64. MS [EI+]  $m/z$  (%): 285 [M]<sup>+</sup> (5), 77 [C<sub>6</sub>H<sub>5</sub>]<sup>+</sup> (100).

#### 3.1.3. (1-Benzyl-1,2,3-triazol-4-ylmethoxy)-benzaldehyde 5

Benzyl azide and 4-Prop-2-ynoxybenzaldehyde afforded (1-Benzyl-1,2,3-triazol-4-ylmethoxy)-benzaldehyde **5** as a white solid, m.p. 79 °C (43%). IR (ATR)  $\nu_{\text{max}}$  1660, 1600 cm<sup>-1</sup>. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  9.88 (s, 1H), 7.54 (s, 1H), 7.82–7.83 (dd, 2H,  $J = 3$  Hz,  $J = 9$  Hz), 7.36–7.38 (m, 3H), 7.26–7.29 (m, 2H), 7.07–7.09 (dd, 2H,  $J = 3$  Hz,  $J = 9$  Hz), 5.54 (s, 2H), 5.26 (s, 2H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  190.7, 163.3, 143.6, 134.3, 130.3, 129.2, 128.9, 128.1, 122.7, 115.0, 62.2, 54.3. MS [EI+]  $m/z$  (%): 293 [M]<sup>+</sup> (5), 91 [C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>]<sup>+</sup> (100).

#### 3.1.4. 1-Benzyl-4-phenyl-1,2,3-triazole 6

Phenylacetylene and benzyl azide afforded 1-Benzyl-4-phenyl-1,2,3-triazole **6** as a white solid, m.p. 131 °C (45%). IR (ATR)  $\nu_{\text{max}}$  3250, 2850, 1650, 1600 cm<sup>-1</sup>. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.82 (m, 2H), 7.68 (s, 1H), 7.41 (m, 4H), 7.33 (m, 1H), 5.59 (s, 2H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  148.2, 134.6, 130.5, 129.1, 128.8, (2 X CH), 128.7, 127.9, 125.6, 119.5, 54.2. MS (EI+)  $m/z$  (%): 235[M]<sup>+</sup> (21), 206 [M – HN<sub>2</sub>]<sup>+</sup> (74), 116 [M – C<sub>6</sub>H<sub>5</sub>N<sub>3</sub>]<sup>+</sup> (100).

#### 3.1.5. 1,4-Diphenyl-1,2,3-triazole 7

Phenylacetylene and phenyl azide afforded 1,4-diphenyl-1,2,3-triazole **7** as a white solid, m.p. 97 °C (40%). IR (ATR)  $\nu_{\text{max}}$  3050, 1600 cm<sup>-1</sup>. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.25–7.92 (m, 10H), 8.20 (s, 1H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  120.5, 121.5, 125.8, 128.2, 128.4, 128.5, 128.7, 128.9, 128.9, 129.7, 130.7, 130.8, 134.9, 147.6. MS [EI+]  $m/z$  (%): 222 [M + 1]<sup>+</sup> (5), 193 [M – N<sub>2</sub>]<sup>+</sup> (95), 165 (100).

#### 3.1.6. 4-Phenyl-1-(3-phenylpropyl)-1,2,3-triazole 8

Phenylacetylene and 3-Azidopropyl benzene afforded 4-phenyl-1-(3-phenylpropyl)-1,2,3-triazole **8** as a white solid, m.p. 238 °C (70%). IR (ATR)  $\nu_{\text{max}}$  1380, 1450, 1696,

3071  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.54 (s, 1H), 7.85 (d,  $J = 7.5$  Hz, 4H), 7.43 (m, 4H), 7.31 (m, 2H), 5.81 (s, 2H), 4.62 (d,  $J = 9$  Hz, 2H), 4.43 (s, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  146.4, 130.8, 128.9, 127.8, 125.1, 122.5, 58.3, 53.2. MS [EI+]  $m/z$  (%): 263 [M]<sup>+</sup> (100), 116 [M – C<sub>9</sub>H<sub>11</sub>N<sub>2</sub>]<sup>+</sup> (30).

### 3.1.7. (1-Benzyl-1,2,3-triazol-4-yl)-methanol **9**

Propargyl alcohol and benzyl azide afforded 1-Benzyl-1,2,3-triazol-4-yl)-methanol **9** as a white solid, m.p. 76–77 °C (65%). IR (ATR)  $\nu_{\text{max}}$  3330, 1572, 1475, 1276  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.47(s, 1H), 7.32 (m, 3H), 7.21 (m, 2H), 5.44 (s, 2H), 4.68 (s, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  148.3, 134.5, 129.0, 128.7, 128.0, 121.9, 55.9, 54.0. MS [EI+]  $m/z$  (%): 189 [M]<sup>+</sup> (40), 91 [C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>]<sup>+</sup> (100).

### 3.1.8. 3-(1-Benzyl-1,2,3-triazol-4-yl)-propan-1-ol **10**

Pent-4-yn-1-ol and benzyl azide afforded 3-(1-Benzyl-1,2,3-triazol-4-yl)-propan-1-ol **10** as a white solid, m.p. 90 °C (53%). IR (ATR)  $\nu_{\text{max}}$  3350, 3050, 1600  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.33 (s, 1H), 7.31 (m, 3H), 7.23 (m, 2H), 5.44 (s, 2H), 4.32 (s, 1H), 3.61 (t, 2H), 2.75 (t, 2H), 1.86 (m, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  148.1, 134.9, 129.0, 128.5, 127.9, 121.3, 61.1, 53.9, 32.0, 21.90. MS [EI+]  $m/z$  (%): 217 [M + 1]<sup>+</sup> (5), 91 [C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>]<sup>+</sup> (100).

### 3.1.9. 2-(1-Benzyl-1,2,3-triazol-4-ylmethyl)-isoindole-1,3-dione **11**

2-Prop-2-ynyl-isoindole-1,3-dione and benzyl azide afforded 2-(1-Benzyl-1,2,3-triazol-4-ylmethyl)-isoindole-1,3-dione **11** as a white solid, m.p. 140 °C (41%). IR (ATR)  $\nu_{\text{max}}$  3112, 3072, 3043, 1702  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.85 (m, 4H), 7.51 (s, 1H), 7.36 (m, 5H), 129.1, 5.48 (s, 2H), 4.97 (s, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  167.6, 143.2, 134.4, 33.06, 134.1, 132.0, 129.1, 128.7, 128.1, 123.4, 122.7, 54.2, 33.1. MS [EI+]  $m/z$  (%): 318 [M]<sup>+</sup> (5), 91 [C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>]<sup>+</sup> (100).

## 4. Conclusions

Diverse 1,2,3-triazoles were obtained as the only reaction product when using glucose/Benedict reagent system as a catalyst through a mild synthetic protocol which did not require other additives, with a high functional group tolerance developed from easily available reagents. The simplicity of this synthetic method suggests that this route to obtaining 1,2,3-triazoles will enjoy widespread application.

**Supplementary Materials:** The following are available online at <https://susy.mdpi.com/user/submission/video/1f5c1b194d7028852dff3553de16303>, Video: Azide-Alkyne Cycloaddition Catalyzed by a Glucose/Benedict Reagent System.

**Author Contributions:** Conceptualization, E.C.-Y.; methodology, J.G.G.T., A.F.B.-B.; formal analysis, L.C.G.-S., M.A.G.-E., M.V.B.U. and E.C.-Y.; investigation, J.G.G.T. and A.F.B.-B.; resources, L.C.G.-S. and E.C.-Y.; data curation, M.A.G.-E., and M.V.B.U.; writing—original draft preparation, E.C.-Y.; writing—review and editing, E.C.-Y.; visualization, E.C.-Y.; supervision, L.C.G.-S. and E.C.-Y.; project administration, L.C.G.-S. and E.C.-Y.; funding acquisition, L.C.G.-S. and E.C.-Y. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by CONACYT-Mexico, project No. A1-S-18230 and fellowship for J.G.G.T.; as well as Centro de Investigaciones y Desarrollo Científico (CIDC)—Universidad Distrital Francisco José de Caldas, project No. 4-50-598-19 and fellowship for A.F.B.-B.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** Financial support from CONACYT is gratefully acknowledged. The authors would like to thank N. Zavala, A. Nuñez, L. Triana, and M. C. Martínez for the technical support.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Kolb, H.C.; Finn, M.G.; Sharpless, K.B. Click Chemistry: Diverse Chemical Function from a Few Good Reactions. *Angew. Chem. Int. Ed.* **2001**, *40*, 2004–2021. [[CrossRef](#)]
2. Christian, W.; Tornøe, C.W.; Christensen, C.; Meldal, M. Peptidotriazoles on Solid Phase: [1,2,3]-Triazoles by Regiospecific Copper(I)-Catalyzed 1,3-Dipolar Cycloadditions of Terminal Alkynes to Azides. *J. Org. Chem.* **2002**, *67*, 3057–3064.
3. Chen, Y.; Tong, Z.R. *Click Chemistry: Approaches, Applications and Challenges*; Nova Science Publishers: New York, NY, USA, 2017.
4. Chandrasekaran, S. *Click Reactions in Organic Synthesis*; Wiley-VCH: Weinheim, Germany, 2016.
5. El-Azab, A.S.; Abdel-Aziz, A.A.M. *Click Chemistry and Applications*; LAP Lambert Academic Publishing: Saarbrücken, Germany, 2014.
6. Lahann, J. *Click Chemistry for Biotechnology and Materials Science*; John Wiley & Sons: Chichester, UK, 2009.
7. Rostovtsev, V.V.; Luke, G.; Green, L.G.; Fokin, V.V.; Sharpless, K.B. A Stepwise Huisgen Cycloaddition Process: Copper(I)-Catalyzed Regioselective “Ligation” of Azides and Terminal Alkynes. *Angew. Chem. Int. Ed.* **2002**, *41*, 2596–2599. [[CrossRef](#)]
8. Konwar, M.; Ali, A.A.; Chetia, M.; Saikia, P.J.; Sarma, D. Fehling solution/DIPEA/hydrazine: An alternative catalytic medium for regioselective synthesis of 1,4-disubstituted-1H-1,2,3-triazoles using azide–alkyne cycloaddition reaction. *Tetrahedron Lett.* **2016**, *57*, 4473–4476. [[CrossRef](#)]
9. García, M.A.; Ríos, Z.G.; González, J.; Pérez, V.M.; Lara, N.; Fuentes, A.; González, C.; Corona, D.; Cuevas-Yañez, E. The Use of Glucose as Alternative Reducing Agent in Copper-Catalyzed Alkyne-Azide Cycloaddition. *Lett. Org. Chem.* **2011**, *8*, 701–706. [[CrossRef](#)]
10. Santana-Martínez, I.; Ramírez-Palma, M.T.; Sánchez-Escalera, J.; Martínez-Otero, D.; García-Eleno, M.A.; Dorazco-González, A.; Cuevas-Yañez, E. Synthesis, structural analysis, and photophysical properties of bi-1,2,3-triazoles. *Struct. Chem.* **2020**, *31*, 191–201. [[CrossRef](#)]
11. Konwar, M.; Hazarika, R.; Ali, A.A.; Chetia, M.; Khupse, N.D.; Saikia, P.J.; Sarma, D. Benedict’s solution/vitamin C: An alternative catalytic protocol for the synthesis of regioselective-1,4-disubstituted-1H-1,2,3-triazoles at room temperature. *Appl. Organomet. Chem.* **2018**, *32*, e4425. [[CrossRef](#)]
12. González, J.; Pérez, V.M.; Jiménez, D.O.; López-Valdez, G.; Corona, D.; Cuevas-Yañez, E. Effect of temperature on triazole and bistriazole formation through copper-catalyzed alkyne–azide cycloaddition. *Tetrahedron Lett.* **2011**, *52*, 3514–3517. [[CrossRef](#)]