

Abstract

Comparative Evaluation of Artificial Neural Networks and Data Analysis in Predicting Liposomes' Size in a Periodic Disturbance Micromixer [†]

Ixchel Ocampo ¹, Rubén R. López ², Vahée Nerguizian ³, Ion Stiharu ^{4,*} and Sergio Camacho León ⁵

¹ School of Engineering and Sciences, Tecnológico de Monterrey, Av. Eugenio Garza Sada 2501 Sur, Monterrey 64849, NL, México; a00800468@itesm.mx

² Departments of Oncology & Pathology, McGill University, Cancer Research Program, MUHC-RI, 1001 Décarie, Montreal, QC H4A 3J1, Canada; ruben.lopezsalazar@mail.mcgill.ca

³ Department of Electrical Engineering, École de Technologie Supérieure, 1100 Notre Dame West, Montreal, QC H3C 1K3, Canada; vahe.nerguizian@ens.etsmtl.ca

⁴ Department of Mechanical and Industrial Engineering, Concordia University, 1455 de Maisonneuve Blvd. West, Montreal, QC H3G 1M8, Canada

⁵ School of Engineering and Sciences, Tecnológico de Monterrey, Av. Eugenio Garza Sada 2501 Sur, Monterrey 64849, NL, México; sergio.camacho@tec.mx

* Correspondence: ion.stiharu@concordia.ca

[†] Presented at the 1st International Conference on Micromachines and Applications, 15–30 April 2021; Available online: <https://micromachines2021.sciforum.net/>.

Abstract: Artificial Neural Networks (ANN) and Data analysis are powerful tools used for supporting decision-making. They have been employed in diverse fields and one of them is nanotechnology used, for example, in predicting particles size. Liposomes are nanoparticles used in different biomedical applications that can be produced in Dean Forces-based Periodic Disturbance Micromixers (PDM). In this work, ANN and data analysis techniques are used to build a liposome size prediction model by using the most relevant variables in a PDM, i.e., Flow Rate Ratio (FRR) and Total Flow Rate (TFR). The ANN was designed in MATLAB and fed data from 60 experiments, which were 70% training, 15% validation and 15% testing. For data analysis, regression analysis was used. The model was evaluated; it showed 98.147% of regression number for training and 97.247% in total data compared with 78.89% regression number obtained by data analysis. These results demonstrate that liposomes' size can be better predicted by ANN with just FRR and TFR as inputs, compared with data analysis techniques when the temperature, solvents, and concentrations are kept constant.

Keywords: artificial neural network; data analysis; prediction model; liposomes; Periodic Disturbance Micromixers

Citation: Ocampo, I.; López, R.R.; Nerguizian, V.; Stiharu, I.; Camacho-León, S. Comparative Evaluation of Artificial Neural Networks and Data Analysis in Predicting Liposomes Size in a Periodic Disturbance. *Eng. Proc.* **2021**, *4*, 42. <https://doi.org/10.3390/Micromachines2021-09549>

Academic Editor: Ion Stiharu

Published: 17 February 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>)

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.