

Antimicrobial Activity Screening of *Camellia japonica* Flowers (var. Carolyn Tuttle) for Potential Drug Development [†]

Antia G. Pereira ¹, Aurora Silva ^{1,2}, Clara Grosso ², Javier Echave ¹, Franklin Chamorro ¹, Sepidar Seyyedi-Mansour ¹, Pauline Donn ¹, Maria Fraga-Corral ¹, Maria Fátima Barroso ^{2,*} and Miguel A. Prieto ^{1,*}

¹ Nutrition and Bromatology Group, Department of Analytical Chemistry and Food Science, Faculty of Science, University of Vigo, E32004 Ourense, Spain; antia.gonzalez.pereira@uvigo.es (A.G.P.); mass@isep.ipp.pt (A.S.); javier.echave@uvigo.es (J.E.); franklin.noel.chamorro@uvigo.es (F.C.); sepidar.seyyedi@uvigo.es (S.S.-M.); donn.pauline@uvigo.es (P.D.); mfraga@uvigo.es (M.F.-C.)

² REQUIMTE/LAQV, Instituto Superior de Engenharia do Porto, Instituto Politécnico do Porto, Rua Dr António Bernardino de Almeida 431, 4249-015 Porto, Portugal; claragrosso@isep.ipp.pt

* Correspondence: mfb@isep.ipp.pt (M.F.B.); mprieto@uvigo.es (M.A.P.)

[†] Presented at the 4th International Electronic Conference on Applied Sciences, 27 October–10 November 2023; Available online: <https://asec2023.sciforum.net/>.

Abstract: The escalating global problem of antibiotic resistance among pathogenic microorganisms necessitates the exploration of effective alternatives to combat multi-resistance. Consumer demand for organic products has stimulated research on natural-origin matrices, such as plants, to develop antimicrobial additives. *Camellia japonica* flowers have demonstrated remarkable biological properties, making them a potential source of bioactive molecules for use as bio-preservatives. This study evaluated the antimicrobial activity of *C. japonica* flowers (var. Carolyn Tuttle) against food-related microorganisms using an agar diffusion assay. Extracts were obtained via a conventional and cost-effective maceration method (50 °C, 1 h) using 60% methanol as the solvent. The results revealed significant antimicrobial activity of *C. japonica* flowers (var. Carolyn Tuttle) against *Staphylococcus aureus* (10.29 mm), *Pseudomonas aeruginosa* (9.24 mm), and *Salmonella enteritidis* (6.95 mm). However, they did not exhibit activity against *Escherichia coli*, *Staphylococcus epidermidis*, and *Bacillus cereus*, unlike other varieties of *C. japonica* which displayed activity against these microorganisms. In conclusion, *C. japonica* flowers (var. Carolyn Tuttle) demonstrated potential as antimicrobial agents with promising applications in the food and pharmaceutical industries. This research contributes to developing natural and organic additives to combat antimicrobial resistance and meet consumer demands for safer and more sustainable products.

Keywords: *Camellia japonica*; flowers; bioactive compounds; antimicrobial



Citation: Pereira, A.G.; Silva, A.; Grosso, C.; Echave, J.; Chamorro, F.; Seyyedi-Mansour, S.; Donn, P.; Fraga-Corral, M.; Barroso, M.F.; Prieto, M.A. Antimicrobial Activity Screening of *Camellia japonica* Flowers (var. Carolyn Tuttle) for Potential Drug Development. *Eng. Proc.* **2023**, *56*, 314. <https://doi.org/10.3390/ASEC2023-15909>

Academic Editor: Alessandra Durazzo

Published: 7 November 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The increasing worldwide challenge posed by antibiotic resistance among pathogenic microorganisms has encouraged the search for viable alternatives to address multi-resistance issues [1]. In response to growing consumer preferences for organic products, there has been a surge in research exploring natural-origin matrices, particularly plants, as potential sources of antimicrobial additives [2]. Among these natural resources is *Camellia japonica*. *C. japonica* is a flowering evergreen shrub native to East Asia, particularly Japan and Korea. This remarkable plant species has captured the attention of researchers for its multifaceted applications, which extend beyond its ornamental uses [3].

C. japonica flowers have demonstrated remarkable biological properties, making them a potential source of bioactive molecules for use as bio-preservatives [4]. They have been traditionally used in various cosmetic and skincare products due to their potential skin-

enhancing qualities [5]. However, beyond their cosmetic utility, *C. japonica* flowers possess inherent qualities that make them an interesting raw material for scientific investigations.

This study aims to assess the antimicrobial activity of *C. japonica* flower extracts, particularly from the 'Carolyn Tuttle' variety, against a range of food-related microorganisms using an agar diffusion assay. By harnessing the natural antimicrobial properties of *C. japonica* flowers, we hope to contribute to the development of sustainable and eco-friendly solutions to combat the pressing issue of antibiotic resistance, thereby safeguarding food safety and quality.

2. Material and Methods

2.1. Chemicals and Reagents

Dimethyl sulfoxide (DMSO), lactic acid, and Mueller Hinton broth (MHB) were procured from Sigma-Aldrich located in Steinheim, Germany. The culture medium, Mulher Hinton Agar II, was obtained from Biolife in Milan, Italy. Strains of *Staphylococcus aureus* (ATCC 25923), *Bacillus cereus* (ATCC 14579), *Pseudomonas aeruginosa* (ATCC 10145), and *Salmonella enteritidis* (ATCC 13076) were generously supplied by Selectrol in Buckingham, UK. Additionally, *Escherichia coli* (NCTC 9001) and *Staphylococcus epidermidis* (NCTC 11047) were sourced from Microbiologics in Minnesota, USA.

2.2. Raw Material and Extraction Protocol

C. japonica petals (var. Carolyn Tuttle) underwent botanical identification through official germplasm banks and reference materials. They were gathered in Northwestern Spain (coordinates 42.431° N, 8.6444° W) in January 2020. Following collection, the samples were subjected to lyophilization using a LyoAlfa10/15 freeze dryer from Telstar, ThermoFisher Scientific, based in Waltham, MA, USA. After lyophilization, the samples were finely pulverized using a blender and subsequently stored at a temperature of −20 °C until the extraction process. The extraction procedure was the same as that previously reported in the literature [4].

2.3. Agar Diffusion Assay

The evaluation of flower extract antimicrobial activity encompassed Gram-positive bacterial strains, including *S. aureus* (ATCC 25923), *S. epidermidis* (NCTC 11047), and *B. cereus* (ATCC 14579), as well as Gram-negative strains, such as *P. aeruginosa* (ATCC 10145), *S. enteritidis* (ATCC 13676), and *E. coli* (NCTC 9001). Samples were dissolved in DMSO to reach a final concentration of 20 mg/mL and subsequently sterilized by filtration using a 0.2 µm syringe filter. To ensure consistency, the initial colony-forming units were normalized to a 0.5 McFarland scale, as determined by measuring turbidity at 600 nm [6].

The assessment of antimicrobial activity followed the methodology established by Paz et al. [7]. The diameters of the inhibition zones were determined using a digital caliper rule. Experimental data were collected in triplicate and are presented as the mean ± standard deviation (SD).

3. Discussion

Table 1 illustrates the antimicrobial activity of *C. japonica* (var. Carolyn Tuttle) against a range of Gram-positive and Gram-negative bacteria, as evaluated through the agar diffusion test. These particular microorganisms were chosen due to their prevalence in food-related contexts. *S. aureus* and *S. epidermidis* were selected because of their propensity to promote opportunistic infections.

The results revealed that *C. japonica* var. Carolyn Tuttle extract showed a greater antimicrobial effect against *S. aureus*, with an inhibition zone of 10.29 mm. Additionally, *P. aeruginosa* and *S. enteritidis* were also sensitive to this *C. japonica* flower extract. In contrast, *E. coli*, *S. epidermidis*, and *B. cereus* were resistant to *C. japonica* var. Carolyn Tuttle, as no inhibition zone was observed against any of these strains. In previous articles, it has been observed that the flowers of *C. japonica* also exhibited antimicrobial activity against

some of these strains, as well as others not considered in this study [8,9]. These variations in activity could be attributed to the use of different solvents, as demonstrated in prior research [9–11]. However, it is worth noting that more comprehensive studies are essential to elucidate the specific compounds responsible for each bioactivity. This will enable a better understanding of the variations among different varieties, linking them to their chemical characterization. Hence, future investigations should prioritize the exploration of potential synergistic and/or antagonistic interactions among the constituents of *C. japonica*. This approach will facilitate a deeper comprehension of the reported bioactivities in flower extracts, ultimately paving the way for their potential industrial applications.

Table 1. Average diameter of inhibition zone $\pm$ standard deviation (mm).

	Microorganism	Inhibition Zone
Gram-negative	<i>P. aeruginosa</i>	9.24 $\pm$ 0.46
	<i>S. enteritidis</i>	6.95 $\pm$ 1.00
	<i>E. coli</i>	Nd
Gram-positive	<i>S. aureus</i>	10.29 $\pm$ 0.46
	<i>S. epidermidis</i>	Nd
	<i>B. cereus</i>	Nd

Nd: Not detected.

Author Contributions: Conceptualization, A.G.P., A.S., C.G., J.E., F.C., S.S.-M., P.D., M.F.-C., M.F.B. and M.A.P.; methodology, A.G.P., A.S., C.G., J.E., F.C., S.S.-M., P.D., M.F.-C. and M.F.B.; software, A.G.P., A.S. and M.F.B.; validation, M.F.B. and M.A.P.; formal analysis, A.S.; investigation, A.G.P., A.S. and M.F.B.; writing—original draft preparation, A.G.P.; writing—review and editing, A.G.P. and M.F.-C.; visualization, A.G.P. and M.F.B.; supervision, A.S., M.F.B. and M.A.P. All authors have read and agreed to the published version of the manuscript.

Funding: The project SYSTEMIC Knowledge hub on Nutrition and Food Security, has received funding from national research funding parties in Belgium (FWO), France (INRA), Germany (BLE), Italy (MIPAAF), Latvia (IZM), Norway (RCN), Portugal (FCT), and Spain (AEI) in a due to the joint action of JPI HDHL, JPI-OCEANS, and FACCE-JPI launched in 2019 under the ERA-NET ERA-HDHL (n° 696295). The authors would like to thank the EU and FCT for funding through the programs UIDB/50006/2020; UIDP/50006/2020; and LA/P/0008/2020 and also to the Ibero-American Program on Science and Technology (CYTED—GENOPSYSN, P223RT0141).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Acknowledgments: The research leading to these results was supported by MICINN supporting the Ramón y Cajal grant for M.A. Prieto (RYC-2017-22891) and by Xunta de Galicia supporting the program EXCELENCIA-ED431F 2020/12 that supports the work of F. Chamorro, the post-doctoral grants of M. Fraga-Corral (ED481B-2019/096) and the pre-doctoral grant of M. Carpena (ED481A 2021/313). The JU receives support from the European Union’s Horizon 2020 research and innovation program and the Bio-Based Industries Consortium. MFB (2020.03107.CEECIND) and Clara Grosso (CEECIND/03436/2020) thank FCT for the FCT Investigator grant.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Vladkova, T.; Georgieva, N.; Staneva, A.; Gospodinova, D. Recent Progress in Antioxidant Active Substances from Marine Biota. *Antioxidants* **2022**, *11*, 439. [CrossRef] [PubMed]
2. Pereira, A.G.; Fraga-Corral, M.; García-Oliveira, P.; Jimenez-Lopez, C.; Lourenço-Lopes, C.; Carpena, M.; Otero, P.; Gullón, P.; Prieto, M.A.; Simal-Gandara, J. Culinary and nutritional value of edible wild plants from northern Spain rich in phenolic compounds with potential health benefits. *Food Funct.* **2020**, *11*, 8493–8515. [CrossRef] [PubMed]

3. Pereira, A.G.; Garcia-Perez, P.; Cassani, L.; Chamorro, F.; Cao, H.; Barba, F.J.; Simal-Gandara, J.; Prieto, M.A. *Camellia japonica*: A phytochemical perspective and current applications facing its industrial exploitation. *Food Chem. X* **2022**, *13*, 100258. [[CrossRef](#)] [[PubMed](#)]
4. Pereira, A.G.; Cassani, L.; Oludemi, T.; Chamorro, F.; Calhelha, R.C.; Prieto, M.A.; Barros, L.; Simal-Gandara, J.; Lucini, L.; Garcia-Perez, P. Untargeted metabolomics and in vitro functional analysis unravel the intraspecific bioactive potential of flowers from underexplored *Camellia japonica* cultivars facing their industrial application. *Ind. Crops Prod.* **2023**, *204*, 117389. [[CrossRef](#)]
5. Shin, S.; Kim, M.; Jung, E. Anti-aging Effects of *Camellia Japonica* flower extract on a pollutant-induced stress. *J. Dermatol. Sci.* **2016**, *84*, e138–e139. [[CrossRef](#)]
6. Silva, A.; Rodrigues, C.; Garcia-Oliveira, P.; Lourenço-Lopes, C.; Silva, S.A.; Garcia-Perez, P.; Carvalho, A.P.; Domingues, V.F.; Barroso, M.F.; Delerue-Matos, C.; et al. Screening of Bioactive Properties in Brown Algae from the Northwest Iberian Peninsula. *Foods* **2021**, *10*, 1915. [[CrossRef](#)] [[PubMed](#)]
7. Paz, M.; Gúllon, P.; Barroso, M.F.; Carvalho, A.P.; Domingues, V.F.; Gomes, A.M.; Becker, H.; Longhinotti, E.; Delerue-Matos, C. Brazilian fruit pulps as functional foods and additives: Evaluation of bioactive compounds. *Food Chem.* **2015**, *172*, 462–468. [[CrossRef](#)] [[PubMed](#)]
8. Pereira, A.G.; Silva, A.; Barral-Martinez, M.; Echave, J.; Chamorro, F.; Mansour, S.S.; Cassani, L.; Otero, P.; Xiao, J.; Barroso, F.; et al. Antimicrobial Activity Screening of *Camellia japonica* Flowers (var. Conde de la Torre). *Med. Sci. Forum* **2022**, *12*, 15.
9. Teixeira, A.M.; Sousa, C. A review on the biological activity of camellia species. *Molecules* **2021**, *26*, 2178. [[CrossRef](#)] [[PubMed](#)]
10. Choi, M.-H.; Min, M.-J.; Oh, D.-S.; Shin, H.-J. Antimicrobial and Antioxidant Activity of *Camellia japonica* Extracts for Cosmetic Applications. *KSBB J.* **2013**, *28*, 99–105. [[CrossRef](#)]
11. Jeon, H.; Kim, J.; Choi, J.; Han, E.; Song, C.-L.; Lee, J.; Cho, Y. Effects of the Extracts from Fruit and Stem of *Camellia japonica* on Induced Pluripotency and Wound Healing. *J. Clin. Med.* **2018**, *7*, 449. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.