








Abstract

Fluorescent Carbon Nanodots as Sensors of Toxic Metal Ions and Pesticides [†]

Federico Bruno ¹, Alice Sciortino ¹, Gianpiero Buscarino ^{1,2}, Marco Cannas ¹, Franco Mario Gelardi ¹, Simonpietro Agnello ^{1,2,*} and Fabrizio Messina ^{1,2,*}

¹ Department of Physics and Chemistry Emilio Segrè, University of Palermo, Via Archirafi 36, 90143 Palermo, Italy; federico.bruno@unipa.it (F.B.); alicesciortino@gmail.com (A.S.); gianpiero.buscarino@unipa.it (G.B.); marco.cannas@unipa.it (M.C.); franco.gelardi@unipa.it (F.M.G.)

² Aten Center, University of Palermo, Viale delle Scienze, Edificio 18, 90128 Palermo, Italy

* Correspondence: simonpietro.agnello@unipa.it (S.A.); fabrizio.messina@unipa.it (F.M.)

[†] Presented at the 8th International Symposium on Sensor Science, 17–28 May 2021; Available online: <https://i3s2021dresden.sciforum.net/>.

Abstract: Carbon nanodots (CDs) are a new class of fluorescent carbon-based nanomaterials characterized by a plethora of morphologies and sizes. Among these, we can include two different types of CDs, namely, graphitic and diamond-like. This wide range of structures opens up the possibility to design different CDs, with tunable optical properties accordingly to the synthesis method and precursors used. We prepared two different CDs following a bottom-up approach by thermally induced decomposition of organic precursors (namely, citric acid and urea in different molar ratios), and using purification by Size Exclusion Chromatography (SEC). Obtained CDs were characterized by Raman, absorption and fluorescence (PL) spectroscopies to understand structural and optical properties, and by atomic force microscopy (AFM) to elucidate morphology. They feature graphitic and diamond-like carbon structures with highly efficient visible emissions. Their sensing towards Cd and Hg heavy metals has been tested by PL experiments. We found a PL quenching in the presence of concentrations of metal salts starting from 0.5 μM and a selectivity towards the interacting ions, depending on the CDs structure, enabling using them for sensing. Furthermore, preliminary experiments suggest that these dots can also be used in principle as sensors of common pesticides. Considering the advantages of carbon dots with respect to other nanomaterials, such as non-toxicity, low cost and ease of synthesis, we consider these results to be very promising in view of exploiting the optical response of carbon dots to fabricate in the near future a variety of pollutant-sensing devices.

Keywords: carbon dots; quenching; fluorescence; pollutants



Citation: Bruno, F.; Sciortino, A.; Buscarino, G.; Cannas, M.; Gelardi, F.M.; Agnello, S.; Messina, F. Fluorescent Carbon Nanodots as Sensors of Toxic Metal Ions and Pesticides. *Eng. Proc.* **2021**, *6*, 21. <https://doi.org/10.3390/I3S2021Dresden-10096>

Academic Editors: Gianaurelio Cuniberti and Larysa Baraban

Published: 17 May 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: No data reported.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).