

Contribution to the Ecophysiological Study of Four Mediterranean Forest Species (*Quercus suber*, *Ceratonia siliqua*, *Tetraclinis articulata*, *Cedrus atlantica*)[†]

Mohamed Mouafik^{1,*} , Mohamed Ouajdi², Oumaima Ninich¹, Jalila Aoujda² and Ahmed El Aboudi¹

¹ Botany, Mycology and Environment Laboratory, Department of Biology, Faculty of Sciences, Mohammed V University, Rabat 10050, Morocco

² Forest Research Center, Silviculture and Forest Health Service, Rabat 10050, Morocco

* Correspondence: mohamed960@gmail.com; Tel.: +212-650586856

† Presented at the 3rd International Electronic Conference on Forests—Exploring New Discoveries and New Directions in Forests, 15–31 October 2022; Available online: <https://iecf2022.sciforum.net/>.

Abstract: Forest degradation has been accentuated in recent years by climate change and drought. This study consists of producing quality plants and understanding their ecophysiological behavior in the face of water stress. The plants of two broadleaves (*Quercus suber*, *Ceratonia siliqua*) and two conifers (*Tetraclinis articulata*, *Cedrus atlantica*) aged 6 months were subjected to water stress. The basic (Ψ_b) and minimum (Ψ_m) leaf water potential was measured and processed according to climatic factors for the 4 forest species. The results show that the leaf water potential in *Quercus suber* presents more negative values (−0.42 MPa) for basic leaf water potential and (−1.43 MPa) for minimum leaf water potential than the other forest species studied. While *Cedrus atlantica* presents the least negative values for minimum leaf water potential (−0.89 MPa). These results allowed us to define the species that is more resistant to water stress and climate change. However, they refer us to more general questions concerning the mechanisms of water use in forest plants.

Keywords: *Quercus suber*; *Ceratonia siliqua*; *Cedrus atlantica*; *Tetraclinis articulata*; climatic data; water stress; leaf water potential



Citation: Mouafik, M.; Ouajdi, M.; Ninich, O.; Aoujda, J.; Aboudi, A.E. Contribution to the Ecophysiological Study of Four Mediterranean Forest Species (*Quercus suber*, *Ceratonia siliqua*, *Tetraclinis articulata*, *Cedrus atlantica*). *Environ. Sci. Proc.* **2022**, *22*, 52. <https://doi.org/10.3390/IECF2022-13125>

Academic Editor: Rodolfo Picchio

Published: 31 October 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Forest ecosystems are terrestrial environments that contain several forms of life where forests play an essential role in the major biogeochemical cycles, the natural cycles of carbon, nitrogen, phosphorus and contribute to the purification of air and water. In Morocco, forest ecosystems are located in semi-arid, sub-humid and humid climates, covering an area of 5,719,000 ha. They consist of 66% hardwood species, 18% coniferous species, 9% artificial plantations and 7% are occupied by low formations [1], with a geographical position between the two seas, which gives it an original Mediterranean climate influenced by the Atlantic Ocean; this gives it a real range of types of bioclimates, and therefore a wealth of biodiversity.

In order to ensure the sustainability of the forest, the Forest Research Center conducts research to identify the hardwood and coniferous species more resistant to drought stress for the regeneration and reforestation of degraded species. In this perspective, we studied the ecophysiological mechanisms of four forest species (*Quercus suber*, *Ceratonia siliqua*, *Tetraclinis articulata*, *Cedrus atlantica*) according to climatic parameters.

2. Materials and Methods

For this study we used seedlings of four forest species aged six months, two hardwoods (*Quercus suber*; *Ceratonia siliqua*) and two conifers (*Tetraclinis articulata*; *Cedrus atlantica*), which were subjected to severe water stress by cessation of irrigation for 30 days under

greenhouse conditions. The physiological measurements were collected every four days interval from three replications per treatment/species.

For physiological parameters, we determined basic (Ψ_b) and minimum (Ψ_m) leaf water potential. The basic leaf water potential (Ψ_b) is the leaf water potential of the plant, measured before sunrise, when the plant is considered to be in equilibrium with soil. The minimum leaf water potential (Ψ_m) is the leaf water potential of the plant, measured in the middle of the day. The study of the evolution of the leaf water potential of our forest plants was made using the *Scholander* pressure chamber; this is a principle which consists of applying pressure to the sample (stem, branch) introduced into the chamber, via the nitrogen gas which is in the bottle, until the sap contained in the sample exudes through the section. The pressure exerted is read on a manometer, then corresponds to the water potential of the leaves in the sample.

For the climatic parameters, such as temperature, humidity, vapor pressure deficit (VPD) and solar radiation were noted with each hour of measurements.

The results obtained were statistically analyzed by the correlation coefficients in order to determine the relationship between the two variables. All these statistical analyzes were carried out using Excel software, version 2208.

3. Results and Discussion

The results of the average evolution in the daily leaf water potential of the four species studied in this work showed differences according to climatic parameters and time (Tables 1 and 2). The study of the evolution of the daily leaf water potential during the water stress cycle revealed differences between the species studied. At the end of the cycle, the average daily leaf water potential of *Quercus suber* reached the most negative values, with -0.42 MPa for the basic leaf water potential. On the other hand, the other species have an almost equal value for the basic leaf water potential (-0.2 MPa, -0.19 MPa and -0.17 MPa), respectively, for *Ceratonia siliqua*, *Cedrus atlantica* and *Tetraclinis articulata*. The minimum values of leaf water potential depend on the evapotranspiration of the plant, which is related to temperature, VPD, and solar radiation. At maximum temperature, the VPD and solar radiation reach their maximum as well as the minimum leaf water potential. The most negative value was recorded for *Quercus suber* which is about -1.44 MPa; similar results have been reported by other authors [2]. In addition, with almost equivalent values for *Tetraclinis articulata* (-1.09 MPa) and for *Ceratonia siliqua* (-1.01 MPa), the leaf water potential does not exceed -1.10 MPa for these two species [3,4]. *Cedrus atlantica* has the least negative value of -0.89 MPa, which fluctuates between -0.5 and -1.7 MPa [5].

Table 1. The average measurements of leaf water potential (Ψ_L).

Hours	Leaf Water Potential (Ψ_L)			
	<i>Quercus suber</i>	<i>Ceratonia siliqua</i>	<i>Tetraclinis articulata</i>	<i>Cedrus atlantica</i>
5 h	-0.42	-0.20	-0.17	-0.19
8 h	-0.75	-0.38	-0.56	-0.40
10 h	-0.91	-0.64	-0.85	-0.78
12 h	-1.44	-1.01	-1.09	-0.89
14 h	-1.03	-0.67	-0.81	-0.62

Table 2. The mean measurements of climatological parameters.

Hours	Temperature (°C)	Relative Humidity (%)	VPD (kPa)	Solar Radiation (w/m ²)
5 h	12.67	82.83	0.27	0
8 h	14.17	83.17	0.31	83.33
10 h	19.67	63.83	0.87	479.33
12 h	22.17	56.33	1.22	732.33
14 h	21.67	55.17	1.02	706.33

According to the Table 3 of leaf water potential and climatic parameters, it can be seen that the leaf water potential of forest species is negatively correlated with VPD and solar radiation. The negative correlation between water potential and climatic parameters (radiation, VPD) is explained by the decrease in water potential and the increase in climatic parameters, with opposite variation and similar intensity. When stress becomes stronger, stomatal closure in response to an increase in VPD has often been described [6,7]. It is a physiological response to water deficit.

Table 3. The correlation coefficient measurements of leaf water potential with VPD and Solar Radiation.

Species	Correlation Coefficient (r)	
	VPD	Solar Radiation
<i>Quercus suber</i>	0.83	0.79
<i>Ceratonia siliqua</i>	0.90	0.84
<i>Tetraclinis articulata</i>	0.83	0.81
<i>Cedrus atlantica</i>	0.84	0.78

4. Conclusions

The present study highlighted the behavior and adaptation of four Mediterranean forest species (*Quercus suber*, *Ceratonia siliqua*, *Tetraclinis articulata* and *Cedrus atlantica*) to water stress. The evolution of daily leaf water potential differs according to forest species, with a strong negative correlation between leaf water potential and climatic parameters (VPD and solar radiation). Moreover, the results obtained revealed that conifers tolerate water stress better than hardwoods.

Author Contributions: Conceptualization, A.E.A. and M.O.; methodology, M.M. and O.N.; validation, M.O. and J.A.; formal analysis, M.M.; writing—original draft preparation, M.M. and O.N.; writing—review and editing, M.M.; supervision, A.E.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author. The data are not publicly available due to privacy restrictions.

Acknowledgments: The authors want to thank all members and staff of the Forest Research Center in Rabat for technical support and guidance.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Mhirit, O.; Et-Tobi, M. *Les écosystèmes forestiers face au changement climatique: Situation et perspectives d'adaptation au Maroc. Programme d'études: Changement climatique: Impacts sur le Maroc et options d'adaptation globales*; Institut Royal des Etudes Stratégiques (IRES): Rabat, Morocco, 2010; 260p.
2. Acherar, M.; Rambal, S.; Lepar, J. Evolution du potentiel hydrique foliaire et de la conductance stomatique de quatre chênes méditerranéens lors d'une période de dessèchement. *Ann. For. Sci.* **1991**, *48*, 561–570. [\[CrossRef\]](#)
3. Rejeb, M.; Laffray, D.; Louguet, P. Modification de la conductance stomatique de diverses origines tunisiennes de caroubier (*Ceratonia siliqua* L.) soumises à une contrainte hydrique prolongée. In *Dans L'amélioration de Plantes Pour L'adaptation Aux Milieux Arides*; John Libbey Eurotext: Paris, France, 1991; pp. 149–158.
4. Aussenac, G.; Oleffe, P. Elements on the comparative water behavior of *Pseudotsuga menziesii*, *Thuja plicata* and *Tsuga heterophylla*. *Revue Forestière Française* **1984**, *36*, 122–126. [\[CrossRef\]](#)
5. Zine El Abidine, A.; Lamhamedi, M.S.; Taoufik, A. Relation hydrique des arbres sains et dépérissants de *Cedrus atlantica* M. au Moyen Atlas tabulaire au Maroc. *Géo-Eco-Trop* **2013**, *37*, 157–176.

6. Froux, F. Caractéristiques hydrauliques, régulation stomatique et efficience d'utilisation de l'eau de quatre espèces de conifères méditerranéens (*Cedrus atlantica*, *Cupressus sempervirens*, *Pinus halepensis* et *Pinus nigra*). Ph.D. Thesis, Université Henri Poincaré-Nancy, Nancy, France, 2002; pp. 7–8.
7. El Aboudi, A. Typologie des Arganeraies Inframéditerranéennes et écophysologie de l'Arganier (*Argania spinosa* (L.) Skeels) Dans le Souss (Maroc). Ph.D. Thesis, The Univ Joseph Fourier, Grenoble, France, 1990; p. 133.