



Abstract Estimating Fire Hazard in a Protected Area of Central Spain (Cabañeros National Park) by a Full Characterization of Vegetation Using LiDAR⁺

Olga Viedma *^D, Victor Cuevas ^D, Ivan Torres and Jose Manuel Moreno *^D

* Correspondence: olga.viedma@uclm.es (O.V.); josem.moreno@uclm.es (J.M.M.)
* Presented at the 3rd International Electronic Conference on Forests—Exploring New Discoveries and New

Directions in Forests, 15–31 October 2022; Available online: https://iecf2022.sciforum.net.

Abstract: The hazardousness of Mediterranean landscapes has increased since the second half of the 20th century, and fuel loads of highly flammable vegetation types have increased throughout the region. Moreover, under the context of more severe fire weather, large fires of high intensity may cause losses in ecosystem services. Accordingly, fire prevention tools to monitor when and where a fire will have the most negative effects through increases in fire severity are required. Fuel characterization is key to wildfire prevention as fuel is one of the primary factors affecting wildfire risk and behavior. Here, we characterized the valuable natural vegetation in Cabañeros National Park (Central Spain) (part of the Natura-2000 network), composed of typical Mediterranean ecosystems, by using LiDAR and other auxiliary data. LiDAR data were obtained from the first Spanish National LiDAR flight, carried out over the study area in 2009–2010. LiDAR data were pre-processed and ground returns were classified using the progressive TIN filter algorithm, carrying out a sensitivity analysis using different settings. Later, the height of the points above the ground were normalized and the Canopy Height Model (CHM) was calculated. Fuel models were derived using the Prometheus fuel classification framework, and they were determined using several LiDAR height metrics and some compositional metrics (i.e., percentage covered by different height ranges) [<0.6 m, 0.6–1 m, 1–2 m, 2–4 m and \geq 4 m] at 30 m. All these metrics allowed for the estimation of fractional canopy cover, fuel height, and vertical continuity. Moreover, tree-tops and crowns were delimited and standard height metrics, as well as vertical profiles, were obtained. All these data, combined with information about the flammability of dominant species and the vulnerability to fire based on functional traits, allowed for the identification of which resource values were most severely exposed to wildfires.

Keywords: fire hazard; fuel complexes; LiDAR; flammability; vulnerability; Natura-2000 network

Published: 27 October 2022

Academic Editor: Rodolfo Picchio

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). **Supplementary Materials:** The following are available online at https://www.mdpi.com/article/10 .3390/IECF2022-13114/s1.

Author Contributions: Conceptualization, O.V., I.T. and J.M.M.; methodology, O.V. and J.M.M.; formal analysis, V.C. and O.V.; investigation, V.C. and O.V.; writing—original draft preparation, V.C. and O.V.; writing—review and editing, I.T., O.V. and J.M.M.; supervision, J.M.M., O.V. and I.T.; funding acquisition, O.V. and J.M.M. All authors have read and agreed to the published version of the manuscript.

Funding: This research is part of the project INFORICAM (PID2020-119402RB-I00), funded by the Spanish MCIN/AEI/10.13039/501100011033 and by the "European Union NextGenerationEU/PRTR".

Informed Consent Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.



Citation: Viedma, O.; Cuevas, V.; Torres, I.; Moreno, J.M. Estimating Fire Hazard in a Protected Area of Central Spain (Cabañeros National Park) by a Full Characterization of Vegetation Using LiDAR. *Environ. Sci. Proc.* 2022, 22, 66. https://doi.org/ 10.3390/IECF2022-13114

Department of Environmental Science, University of Castilla-La Mancha, Avda. Carlos III, 45071 Toledo, Spain