

Article

# Structure of the Charge-Transfer State in PM6/Y6 and PM6/Y6:YT Composites Studied by Electron Spin Echo Technique


Ekaterina A. Lukina <sup>1</sup>, Aina V. Kulikova <sup>1</sup>, Mikhail N. Uvarov <sup>1</sup>, Alexander A. Popov <sup>2</sup>, Ming Liu <sup>3</sup>, Yong Zhang <sup>3</sup> and Leonid V. Kulik <sup>1,\*</sup>

<sup>1</sup> Voevodsky Institute of Chemical Kinetics and Combustion of the Siberian Branch of the Russian Academy of Sciences, 630090 Novosibirsk, Russia


<sup>2</sup> Physical Department of Novosibirsk State University, 630090 Novosibirsk, Russia

<sup>3</sup> School of Materials Science and Engineering, Harbin Institute of Technology, Harbin 150001, China; yongzhang@hit.edu.cn (Y.Z.)

\* Correspondence: chemphy@kinetics.nsc.ru



**Figure S1.** Echo-detected EPR spectrum of the spin-polarized species in PM6/Y-T composite taken at temperature 80 K with DAF = 1  $\mu$ s. Black lines correspond to in-phase component of the spectrum; red lines correspond to out-of-phase component of the spectrum. The number of scans for signal averaging is 10 times larger than that for PM6/Y6 and PM6/Y6:Y-T composites (Fig. 2 in the main text).



**Figure S2.** Field-averaged out-of-phase ESEEM trace for PM6/Y6 composite spin-coated on glass plate. Temperature 80 K with DAF = 200 ns.