

Dental Patients' Perspective on COVID-19: A Systematic Review

Stefano Salgarello , Elisabetta Audino, Paolo Bertoletti, Matteo Salvadori and Maria Luisa Garo *

Department of Medical and Surgery Specialties, Dental School, University of Brescia, Radiological Sciences and Public Health, 25123 Brescia, Italy; stefano.salgarello@unibs.it (S.S.); bettaz92@gmail.com (E.A.); paolobertoletti1993@gmail.com (P.B.); salvaz.dori@gmail.com (M.S.)

* Correspondence: marilu.garo@gmail.com; Tel.: +39-030-383424

Definition: The COVID-19 epidemic has changed patients' approach to dental treatments. While dentists worldwide have shown an excellent level of adaptability to face the new challenges presented by the unprecedented situation due to the rapid spread of COVID-19, dental patients have witnessed a sudden suspension of elective treatments and a slow resumption of dental care activities after several national lockdowns. In addition, the general climate of anxiety and fear due to the high COVID-19 risk and the high level of mortality has influenced the perception and attitudes of people towards dental activity, inducing many dental patients to avoid appointments to the dentist if not highly urgent. We present an overview of the current state of knowledge about dental patients' perception, perspective, attitude, and expectations towards a full resumption of regular dental treatments.

Keywords: perception; dental patient; dental treatment; safety; COVID-19



Citation: Salgarello, S.; Audino, E.; Bertoletti, P.; Salvadori, M.; Garo, M.L. Dental Patients' Perspective on COVID-19: A Systematic Review. *Encyclopedia* **2022**, *2*, 365–382. <https://doi.org/10.3390/encyclopedia2010022>

Academic Editors: Stephen Bustin and Raffaele Barretta

Received: 17 January 2022

Accepted: 26 January 2022

Published: 1 February 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Since the beginning of the COVID-19 pandemic, dental professionals have been exposed to a high level of anxiety and stress due to several changes that have affected dentistry in many ways. To date, many international and national guidelines and recommendations are available to support dentists in their everyday dental activities [1,2]. In addition, many studies have investigated dentists' perceptions, problems, attitudes, and expectations worldwide to understand how dentists have faced the unprecedented situation due to the COVID-19 outbreak [3–5].

The management of the emergencies and urgencies in public and private dental clinics during the various national lockdown, the handling of COVID-19 patients in need of urgent dental treatments, the optimization of the personal protective equipment (PPE), and the appropriate approach to resume the regular dental activity with a high level of safety for dental staff and patients have been deeply analyzed [6–10].

In the first COVID-19 outbreak period, many countries adopted prudential measures to reduce the risk of contagion, such as suspending elective dental treatments and recommending executing only non-postponable dental treatments. However, after the first waves of COVID-19, the need to restart private and public dental activities has become the first and foremost goal of many dental professionals [11].

2. Synthesis

From the end of 2020, the availability of COVID-19 vaccines has supported a complete restart of dental activity, allowing the medical staff to resume their work with a certain level of safety and confidence [12,13].

Therefore, if, on the one hand, COVID-19 has created an unprecedented situation for all health services, on the other hand, this is also an unknown situation for patients, who have to handle fear of COVID-19 contagion and discomfort situations due to the suspension of planned dental treatments [14,15].

In this context, the psychological consequence of COVID-19 should not be neglected [16]. As emerged in a recent Chinese study, people living in a pandemic can suffer from different levels of psychological distress, experiencing increased fear of becoming infected [17]. In dentistry, this higher level of stress and anxiety can be translated into a series of ugly behaviors, such as avoidance of dental appointments, treatment delays, and adoption of therapies based on their experience and common sense without the support of their dentists [18].

This work aimed to review the current literature about the role of COVID-19 in the perceptions, perspectives, attitudes, and expectations of patients toward dental treatments and the risk of COVID-19 infection within dental clinics.

2.1. Methods

A systematic review of the current literature was performed following PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines [19]. On the basis of the inclusion criteria, we selected original studies reporting dental patients' perceptions or perspectives during the COVID-19 pandemic and their attitudes and expectations about the safe restart of dental clinics and dental offices activities.

PubMed and Web of Science databases were searched using the following search terms: (COVID-19 [Title/Abstract]) AND (dental patient [Title/Abstract]) AND (perspective or perception [Title/Abstract]).

The search strategy was carried out from September to October 2021, without time and language restriction. First, the resultant citations were screened by title, followed by abstracts and then full texts. In addition to the initial search, references in the selected sources were checked manually, and further studies were included if relevant.

Additionally, hand searches were performed to identify possible articles other than those found in the electronic databases. Two reviewers (MLG and EA) performed the first (title/abstract screening) and second (full-text assessment) steps of the search process. Any disagreement was discussed and then solved by consensus.

All original peer-reviewed research publications were considered. Inclusion criteria for eligible studies were: (1) studies carried out on COVID-19 impact in dentistry; (2) studies reporting dental patients' perceptions, perspectives, or attitudes toward dental treatment during the COVID-19 pandemic; (3) studies analyzing dental patients' anxiety or fear.

Data extraction was organized in tables containing the following information:

- Study characteristics: first author, year, country.
- Study design, period, and setting.
- Sample size and patients' characteristics (gender and age).
- Types of dental emergency/urgency.
- Questionnaire structure.
- Main outcomes.

No numerical information was extracted from the figures reported in the study publications.

The risk of bias was assessed through the CHERRIES (*C*hecklist for *R*eporting *R*esults of *I*nternet *E*-*S*urveys) or SURge (*S*urvey *R*eporting *G*uideline), as suggested by Weir et al. [20].

Given the lack of a validated checklist to evaluate the surveys, we adapted the CHERRIES checklist for web-based surveys and SURge for self-administered surveys non-online administered. For both checklists, we applied the following procedure. For each checklist item, we assigned a value of 1 if the topic was reported in the study, and a value of 0 if it was not registered. Each survey's final score was determined by calculating the sum of all valid items divided by their number. Low-, moderate-, or high-quality studies were assessed according to the following evaluation: final score < 35% = low quality; 35% ≤ final score < 75% = moderate quality; final score ≥ 75% = high quality.

2.2. Study Selection

The literature search identified 205 studies. After removing duplicates and screening titles and abstracts, 15 full-text studies were reviewed. Ten studies between 2020 and 2021

met the inclusion and exclusion criteria. A flow diagram of the search strategy results is presented in Figure 1.




Figure 1. PRISMA Flow-chart.

2.3. Risk of Bias

Given the lack of a specific tool for evaluating bias in surveys, we assessed the risk of bias using two checklists. The overall risk of bias assessment of the included studies is presented in Tables S1 and S2 in Supplementary Materials. Five studies reported a moderate level of bias [21–25], while the remaining five studies showed low quality [26–30].

In surveys administered through online platforms (Table S1 in Supplementary Materials), all studies reported the survey design describing the target population or sample frame. For IRB approval and informed consent process, IRB approval was obtained by seven out of eight studies [21–24,26,29,30], informed consent was obtained by four studies [22,25,28,29], and data protection was reported by only one study [22], while it was not applicable by four surveys [21,23,27,29]. Development and pre-testing were performed by only two studies [22,30]. For the recruitment process and description of the sample having access to the questionnaire, a closed survey was applied by all studies; only one indicated that the contact mode was not through the Internet [21], while advertising to announce the survey was carried out only by Moffat et al. [22]. All surveys were administered through the web or by email; six out of eight surveys described the context [21–24,28,30], and only two studies were not mandatory surveys [28,30]. Only one study used incentives [22], although this item was applicable for only two studies [21,22]. The timeframe was identified in five studies [21–24,28]. All included studies did not report the number of items, number of screens, and completeness check, while review steps were only adopted by Peloso et al. [29]. The unique visitor ID was assessed by two surveys [21,29], while none of the studies reported view rate, participation rate, and completion rate. For preventing multiple entries from the same individual, i.e., using cookies, IP check, log file analysis, and registration, no precautions were adopted by all surveys. Information about the handling of incomplete questionnaires was not described. Statistical corrections were adopted by five studies [21–24,28].

Two studies were evaluated through SURge [25,27]. Although with slight differences, both studies lacked information regarding many items. Mainly, neither study reported the procedures used to develop, pre-test, and validate the new survey, sample size calculation and its representativeness, the type of contact, and incentives. Furthermore, the statistical analysis did not provide information about non-response error, response rate, and handling of missing data. In addition, neither study reported considerations about the non-response rate and possible differences in respondents' samples and discussed the external validity of the findings. The complete results of risk of bias for both studies are reported in Table S2 (Supplementary Materials).

2.4. Study Characteristics

We provide a descriptive summary of the study design, period, setting, sample size, gender, age, types of dental emergency/urgency treated during the period of interest, and the questionnaire structure in Table 1. We included a total of 10 studies: 2 carried out in Brazil [26,29], 2 in the U.S.A. [22,25], 1 in Spain [21], Turkey [27], Italy [28], Saudi Arabia [23], China [24], and Nigeria [30], respectively.

Table 1. Studies characteristics.

Study	Country	Study Design	Period	Setting	Sample Size	Gender	Age	Types of Dental Emergency/Urgency	Questionnaire Structure	Main Outcomes
Cotrin et al. (2020) [26]	Brasil	Survey	NR *	Private Dental Clinics	354	231 females and 123 males	35.49 ± 13.93	Orthodontic treatments	<ul style="list-style-type: none"> • Socio-Demographics • Quarantine • Level of anxiety (NRS): 0 no anxiety—10 extreme anxiety 	<ul style="list-style-type: none"> • 78.2% go out just when needed; 13% not leave home; 8.8% go out as usual without respecting the quarantine • Feelings about COVID-19: 44.7% is calm; 23.4% is afraid; 22.9% is anxious; 5.6% is indifferent; 3.4% is in panic • Level of anxiety: 4.98 ± 2.42 • Dental appointment attendance: 60.2% would go to an appointment; 25.1% would go only in case of urgency/emergency; 14.7% would not go • Females more anxious, afraid, and in panic than men in relation to quarantine and the COVID-19 pandemic • Males were more willing to go a dental appointment than females • Males were more concerned about a delay in treatment than females • PPE fundamental: surgical mask (88.7%), medical head cap (81.1%), lab coat (55.9%), face shield (55.1%) • Good practices: avoiding crossing other patients at reception (78%), PPE for patients (35%), alcohol gel available at reception (91.8%)

Table 1. Cont.

Study	Country	Study Design	Period	Setting	Sample Size	Gender	Age	Types of Dental Emergency/Urgency	Questionnaire Structure	Main Outcomes
Moffat et al. (2020) [22]	USA	Survey	May 2020	Web-survey through Amazon Mechanical Turk (MTurk)	452	198 females, 250 males, and 4 NR	40 years	None	<ul style="list-style-type: none"> Socio-Demographics Perceived susceptibility to contract COVID-19 during dental appointments Attitudes and beliefs regarding the risk of contracting COVID-19 during dental appointments Events that need to occur for patients to feel comfortable returning to the dental office 	<ul style="list-style-type: none"> Concerned about contracting COVID-19 from dental professionals (3.7 ± 1.7) and other patients in the dental office (3.3 ± 1.8) Aware of the government recommendations to dental professionals for appropriate dental treatment during the COVID-19 outbreak (3.8 ± 1.9) Trust that dental offices follow government recommendations to dental professionals for appropriate dental treatment during the COVID-19 outbreak (2.2 ± 1.2) Concerned that visiting dental providers will cause a shortage in PPE available for healthcare providers fighting COVID-19 (4.5 ± 1.9) The risk of getting COVID-19 in the dental office outweighs the risk of me not getting necessary dental treatment completed (3.6 ± 1.7)
Peloso et al. (2020) [29]	Brazil	Survey	NR *	Private Dental Clinics	595	412 females and 183 males	38.21 ± 13.94	74.3% orthodontic 18.4% oral rehabilitation 7.3% restorative or other	<ul style="list-style-type: none"> Socio-Demographics Presence of symptoms related to COVID-19 Dental treatments received during the quarantine Contacts with own dentist to schedule an appointments Willing to go to dentist for an emergency Concerns about attending a dental appointment PPE used by dentist and in the dental office 	<ul style="list-style-type: none"> Concerned about attending a dental appointment included the risk of getting infected and/or contaminating the family (18.5%) 5% afraid because dentists are a group at high risk of contamination Dental treatments can wait (20.2%) Females more anxious and afraid than males Males more willing to go a dental appointment Females not worried about how quarantine might affect their dental treatment More anxious, afraid, or panicked would attend an appointment in the case of emergency (47.1%, 46.4%, and 62.5%) Importance of disposable lab coat (63.9%), disposable mask (67.4%), having alcohol gel at reception (85.4%), avoiding close contact with other patients at reception (63.5%), having PPE for patients (41.8%)

Table 1. Cont.

Study	Country	Study Design	Period	Setting	Sample Size	Gender	Age	Types of Dental Emergency/Urgency	Questionnaire Structure	Main Outcomes
Arqub et al. (2021) [25]	USA	Survey	July–October 2020	Dental Clinic	154	95 females and 59 males	29.30 ± 12.01	Orthodontic treatments	<ul style="list-style-type: none"> • Socio-Demographics • Physiological aspects relevant to clinical orthodontics • Emergencies • Preferences regarding the urgency on following up treatment during the pandemic and preferred communication means • Patients' insight on postponing or discontinuing treatment, schedules appointment intervals due to safety concerns • Clinic facilities, screening protocols, pretesting for COVID-19 • Phycological symptoms (Kessler Psychological Distress Scale) 	<ul style="list-style-type: none"> • Extremely pleased with the restrictive protocols at the clinic (80.51%) and the idea that telephone and pre-clinical appointment screening is an effective tool to prevent the spread of infection (68%) • Useful tools: pretesting patients (81.81%), the staff, and orthodontists (85.06%) during the pandemic • Dentist should wear full PPE and N95 mask (45.5%)

Table 1. Cont.

Study	Country	Study Design	Period	Setting	Sample Size	Gender	Age	Types of Dental Emergency/Urgency	Questionnaire Structure	Main Outcomes
Gonzales-Olmo et al. (2021) [21]	Spain	Survey—Repeated measures design (before and after Spain lockdown)	March–June 2020	Web-survey addressed to community of Madrid	961	559 females and 402 males	38.4 ± 16.1	None	<ul style="list-style-type: none"> • Socio-Demographics • Perceived vulnerability to disease (PVD) • Scale of fear of COVID-19 • Structured questions about avoidance behavior towards the dental clinic • Question covering whether the participant had been ill with COVID-19 	<ul style="list-style-type: none"> • Significantly higher scores on T1 than T0, both in infectability ($p < 0.01$) and in aversion to germs ($p < 0.01$) • Significant positive correlation between COVID-19 fear scale and the subscales of infectability and germ aversion in T0 and T1 ($p < 0.01$) • Strong positive association between fear of COVID-19 in T1 and aversion to germs in T0 ($p < 0.01$) • Higher score for women for infectability and germ aversion in T1 ($p < 0.01$) • Those who contracted COVID-19 reported a higher COVID fear score ($p < 0.01$) • 30.9% afraid of going to the dentist because of the possibility of contagion with COVID-19, although 56.3% continues to go to the dentist • 23.5% would go to the dentist because they had not finished their treatments, but 42.5% would not start esthetic, orthodontic, or implant treatments • 43.7% would not go to the dentist, 24.5% for fear of COVID-19, 16% because of financial problems, 3.2% for other reasons—more than half would maintain this decision until the disease is eradicated or an effective treatment is found • 20.7% would not go to the dentist even in case of gum problems; 20.2% would not go even if they suspected cavities; 16.3% would not go even if filling or teeth were fractured • Dentist avoidance: high perceived infectability (OR = 4.21, 95%CI: 2.87–5.64), high fear of COVID-19 (OR = 5.18, 95%CI: 2.96–9.4), older than 60 years (OR = 7.63, 95%CI: 3.56–15.35)

Table 1. Cont.

Study	Country	Study Design	Period	Setting	Sample Size	Gender	Age	Types of Dental Emergency/Urgency	Questionnaire Structure	Main Outcomes
Karagözoğlu et al. (2021) [27]	Turkey	Survey	NR	Dentistry Faculty	300	145 females and 155 males	18–38: 53%, 39–59: 31.7%; 60–80: 15.3%	Patients receiving dental treatment	<ul style="list-style-type: none"> Socio-Demographics COVID-19 and patient-related questions Spielberg’s STAT-S anxiety scale 	<ul style="list-style-type: none"> Mean anxiety score: 43.38 ± 8.34 Higher anxiety score in those who think that COVID-19 can be transmitted through dental treatment ($p < 0.001$), would not seek dental treatment if their condition was not urgent ($p < 0.003$), think that dentists have a high risk of contamination, thought that dentist can transmit COVID-19 during dental treatment, and believe they could be infected by the other patients in the waiting area Lower anxiety score in those who thought that adequate measures were taken in the dental unit ($p < 0.001$), the measures taken by dentists while treating patients were sufficient, feel safe because healthcare professionals received the COVID-19 vaccine, and would have had the dental treatment safely if vaccinated
Martina et al. (2021) [28]	Italy	Survey	May–November 2020	Web Survey	1566	852 females, 698 males, 16 NR	18–29: 31.3%; 30–39: 29.8%; 40–49: 18.3%; 50–59: 10.6%; 60–69: 6.8%; >70.3%	Orthodontic care	<ul style="list-style-type: none"> Socio-Demographics Anxiety over going to the dentist Presence of TMD symptoms Orthodontic treatment and the perception of the risk of continuing orthodontic treatment 	<ul style="list-style-type: none"> 55.3% believed that there is a greater risk of contracting COVID-19 at a dental practice Men less afraid to go to a dentist clinic Younger people less scared to go to a dental clinic than people over 40 ($p < 0.001$) Higher level of anxiety was associated with a fear of going to a dental practice 79.5% thought than in the dental practice there was a greater risk of contracting COVID-19 For 84.6%, COVID-19 increased the fear of going to the dentist

Table 1. Cont.

Study	Country	Study Design	Period	Setting	Sample Size	Gender	Age	Types of Dental Emergency/Urgency	Questionnaire Structure	Main Outcomes
Nazir et al. (2021) [23]	Saudi Arabia	Survey	June–July 2020	Scientific research unit at the College of Dentistry	606	246 females and 360 males	30.49 ± 12.01	Patients receiving dental treatment	<ul style="list-style-type: none"> • Socio-Demographics • Dental Fear Survey (DFS) and Modified Dental Anxiety Scale (MDAS) • Dental attendance patterns • Visiting a dental office in case of emergency during the COVID-19 pandemic 	<ul style="list-style-type: none"> • Higher DFS in females (41.17 ± 17.94) than in males (37.65 ± 20.34) ($p = 0.029$)
Quan et al. (2021) [24]	China	Survey	February–March 2020	Dental Clinic	1078	786 females and 292 males	22.59 ± 8.28	Orthodontic patients	<ul style="list-style-type: none"> • Socio-Demographics • Self-management ability of oral health and compliance and cooperation • Orthodontic problems and emergencies during the pandemic • Risk perception about the COVID-19 pandemic • Self-Rating Anxiety Scale 	<ul style="list-style-type: none"> • Patients who faced orthodontic problems: 55.1% contacted their attending doctor; 33.3% thought the problem was tolerable and did not take any measure; 23.1% indicated they could solve it by themselves according to their experiences or common sense. • Difficulties of orthodontic follow-up visits: 67.9% worried about the risk of infection

Table 1. Cont.

Study	Country	Study Design	Period	Setting	Sample Size	Gender	Age	Types of Dental Emergency/Urgency	Questionnaire Structure	Main Outcomes
Umeh et al. (2021) [30]	Nigeria	Survey		Public and private dental facilities	304	221 females and 83 males	35.6	Orthodontic patients receive treatment	<ul style="list-style-type: none"> • Socio-Demographics • Perception of COVID-19 risk in orthodontic treatment • Knowledge about COVID-19 infection • Risk perception and attitude toward the COVID-19 infection 	<ul style="list-style-type: none"> • Orthodontic patients were severely (40.8%) or moderately (30.6%) vulnerable to contracting the disease but were willing regardless to carry on with their orthodontic treatment during the pandemic (72.4%) • Fear of contracting the virus during a routine appointment: 39.8% • Missed routine appointment: 27.3% • Increased treatment time: 35.5% • Increase in cost of treatment: 8.6% • Patients have a significant role in infection control in the orthodontic practice setting (81.3% social distancing in the clinic, 82.9% temperature screening before entering the clinic; 69.7%; asked relevant screening question; 78.9% limiting child's company; 80.9%; wearing facemasks in the clinic; 88.5% handwashing in the dental clinic)

Studies were carried out in 2020–2021: Quan et al. from February to March 2020, Gonzales et al. from March to June 2020, Moffat et al. in May 2020, Martina et al. from May to November 2020, Arqub from July to October 2020, and Nazir et al. from June to July 2020. Two studies [26,29] did not report the period of questionnaire administration but referred to that period being before the COVID-19 peak in Brazil: we presume before June 2020. Respondents were chosen in different settings: for four studies in public dental clinics or however related to academic scientific research units [24,26,28,29], three studies administered the questionnaires through private dental clinics [21,23,27], and two through web-survey addressed to local communities [21,22]. Only one study distributed the questionnaires to public and private dental clinics [30].

Six thousand three hundred and seventy (6370) patients were included in the studies. Three thousand seven hundred and forty-five (3745) respondents were female (female-male ratio 1.44). The mean age varied between 22.59 ± 8.28 [24] and 40 years [22]. Two studies reported age ranges instead of mean age [27,28], showing a high prevalence of young people (18–38 years) and a mild prevalence of older people (older than 60 years). Six studies involved mainly patients with orthodontic problems [21,23,24,27,29,30], two studies involved patients who received dental treatments after the COVID-19 outbreak [23,27], and the other two studies involved respondents who did not necessarily receive dental treatments after the COVID-19 outbreak [21,22].

All questionnaires were structured to gather data about socio-demographic questions (e.g., age, gender, education background, ethnicity, income, and city of residence). Three investigated COVID-19 and patient-related problems such as quarantine, COVID-19 symptoms, and COVID-19 infection [21,23,26]. Six studies investigated patients' anxiety toward dental treatments through different validated or not-validated psychometric scales [21,23–27]. Nine studies investigated the risk perceptions, attitudes, avoidance behaviors, and beliefs toward the COVID-19 infection during dental treatments [21–26,28–30]. Five studies investigated protocols and procedures adopted by dental clinics to safeguard patients and allow them to feel comfortable about returning to the dental offices [22,25,26,29,30]. One study analyzed preferences regarding the urgency of following up treatments during the pandemic and preferred communication methods with the dentist [25]. Gonzales et al. only investigated possible reasons to avoid dental treatments [21].

2.5. Anxiety, Risk Perception, and Attitudes toward Dental Treatments during the Pandemic

Cotrin et al. [26], in a survey carried out presumably from March to June 2020, including 354 Brazilian respondents, reported a high level of anxiety (4.98 ± 2.42) among quarantine patients, with a substantial prevalence among females. Feelings about COVID-19 ranged from a relaxed (44.7%) or indifferent (5.6%) attitude to feelings of pain (3.4%), fear (23.4%), or anxiety (22.9%). Males were more willing to go to a dental appointment and more concerned about the delay in dental treatments due to COVID-19 restrictions. Dental appointment attendance continued to represent an important factor in everyday life, because 60.2% of the respondents would go to dental appointments for elective treatments. In comparison, only 25.1% of them would go only in case of urgency or emergency. About 15% of the respondents would not go to a dental appointment due to fear of contagion.

The same level of anxiety and fear emerged in Gonzales et al. study [21], which involved people living in Madrid. In an accurate comparison of the attitudes and perceptions of people toward dental treatment during the Spanish lockdown (March–June 2020), they showed that 30.9% of the respondents were afraid of going to the dentist because of the possibility of contagion with COVID-19. However, 56.3% of them continued to go to the dentist. More than 23% of the respondents decided to continue to go to the dentist because the planned dental treatment was not finished. The fear of COVID-19 impacted the start of new dental treatments such as orthodontics or implants: 42.5% of the Spanish respondents reported no intention to start a new treatment because of financial problems (16%) or fear of COVID-19 (24.5%). More than half of the interviewees would not go to the dentist until the disease was eradicated or an effective treatment was found. About 21% of the

respondents stated they would not go to the dentist even in case of gum problems, 20.2% would not go even if they suspected cavities, and 16.3% would not go even if filling or teeth were fractured. The probability of dental avoidance due to the fear of COVID-19 infection was equal to 84% (OR = 5.18, 95%CI: 2.96–9.4) and increased to 88% in people older than 60 years (OR = 7.63, 95%CI: 3.56–15.35). Comparing infectability perception and aversion to germs at the beginning and at the end of the Spanish lockdown, Gonzales et al. showed an increase in the score of perceived vulnerability to disease, with a significant positive correlation between COVID-19 fear scale and infectability perception or aversion to germs in the post-lockdown period ($p < 0.01$).

In a study involving 300 patients who received dental treatments after the COVID-19 outbreak, Karagözoğlu et al. [27] reported a high level of anxiety (43.38 ± 8.34), especially in those who thought that COVID-19 could be transmitted through dental treatment ($p < 0.001$). Moreover, they showed that dental clinics were considered places with a high risk of contamination, given the close contact with dentists and other patients. Patients who would not seek dental treatment if their condition was not urgent also showed a higher level of anxiety ($p < 0.003$). On the contrary, a lower anxiety score was reported in patients who thought that adequate measures were taken in the dental unit ($p < 0.001$) and the measures taken by the dentists while treating patients were sufficient. A lower level of anxiety was also registered in patients who felt safe because healthcare professionals received the COVID-19 vaccine.

In a large study conducted in China involving 1078 orthodontic patients, Quan et al. [24] reported some difficulties for follow-up visits in 67.9% of respondents because of the risk of COVID-19 infection. However, during the pandemic crisis, 55.1% of the respondents contacted their attending doctors; on the contrary, 33.3% did not take any measure, and 23.1% even preferred to solve the problem according to their experience or common sense.

Two studies [23,25] did not report a significant level of dental anxiety related to the COVID-19 outbreak. However, Nazir et al. registered a higher level of dental fear among females (41.17 ± 17.94) than among males (37.65 ± 20.34) ($p = 0.029$).

In an American survey conducted in May 2020 through Amazon Mechanical Turk, Moffat et al. [22] interviewed 448 people. Many concerns were reported about contracting COVID-19 from dental professionals (3.7 ± 1.7) and other patients (3.3 ± 1.8) in the dental offices. However, the risk of getting COVID-19 in a dental office outweighed the risk of not getting necessary dental treatments.

Peloso et al. [29] in their study about patients' perceptions in private dental clinics, that involved 595 Brazilians, showed that some interviewees were concerned about attending dental appointments because of the possibility of getting infected and/or of contaminating their family (18.5%), sometimes because dentists were considered at high risk of contamination. About 47% and 63% of the respondents reported increased feelings of anxiety, fear, or pain when the dental appointment was not postponable because of an emergency. About 20% of the respondents said that dental treatment could wait. In this study also, females were more worried about COVID-19 infection and less willing to go to dental appointments than males.

In a large study conducted on 1566 Italian orthodontic patients interviewed after the Italian lockdown (March–May 2020), Martina et al. [28] showed that 55.3% of the respondents believed there was a greater risk of COVID-19 infection during dental treatments. Furthermore, for 84.6% of patients, COVID-19 increased the fear of going to the dentist. Younger people and males were less scared to go to the dentist, while females and people older than 40 showed a high level of fear.

In a Nigerian survey conducted from 2020 until 2021 in public and private dental facilities involving 304 orthodontic patients, Umeh et al. [30] showed that people perceived themselves severely or moderately vulnerable to contracting COVID-19 (40.8% and 30.5%, respectively) but were willing to carry on with their dental treatments during the pandemic (72.4%). However, approximately 40% of the interviewees reported a certain level of fear of

contracting the virus during dental appointments. In addition, the patients perceived the increase in treatment time and cost.

2.6. Protocols and Procedures for COVID-19 Infection Control in the Dental Setting

Five studies investigated the patients' perception of protocols and procedures adopted in dental clinics to reduce the risk of COVID-19 infection [22,25,26,29,30].

Two studies [26,29], both carried out in Brazil, showed that wearing PPE improved patients' confidence in dental professionals. In Cotrin et al. [26], more than 80% of the patients reported reduced anxiety and fear when dentists wore surgical masks and medical head caps during dental treatments; over half of the respondents reported a positive reduction of fear when dentists and medical staff wore face shields. In addition, some safety procedures, such as alcohol gel for patients and avoidance of crossing patients in the waiting areas, showed a positive effect in reducing the fear of COVID-19 infection in dental offices. The same trend, although with some percentage reductions, also emerged in Peloso et al. [29]

Moffat et al. [22] focused on patients' confidence and awareness about the appropriate use of government recommendations within dental offices. The respondents reported low levels of trust that dental offices follow government recommendations to dental professionals during the COVID-19 pandemic (2.2 ± 1.2). At the same time, they perceived themselves as aware of the government recommendation to dental professionals for appropriate dental treatments during the COVID-19 outbreaks (3.8 ± 1.9).

In Arqub et al. [25], approximately 80% of the respondents believed that pretesting patients and medical staff during the pandemic reduced patients' fear of COVID-19 infection in dental offices; wearing full PPE or N95 masks by dentists during dental treatments helped to increase patients' trust in the safety of dental treatments.

Finally, Umeh et al. [30] reported that patients attributed a crucial role to many safety procedures to access dental offices: social distancing, temperature screening before access, relevant screening questions, limiting the number of adults accompanying a child, wearing facemasks, and handwashing were found to play a crucial role in enhancing patients' trust.

3. Current Status

This work aimed to understand dental patients' perspectives, perceptions, attitudes, and expectations toward dental treatments after the first year of the COVID-19 pandemic. In the last two years, many studies have investigated the role of dentists in reducing COVID-19 risk infection and resuming their daily activity keeping a high level of safety [31]. Although dental professionals have been considered at high risk due to the close contact with patients' mouth and the upper respiratory tract [2] and the use of aerosol-generating procedures [32–35], many recommendations and guidelines have provided support to restart dental activity safely.

For the patients, returning to good oral health habits without increasing the risk of COVID-19 infection, and therefore returning to routine dental care, is very important as it provides extensive benefits to the population [22].

Many encouraging findings have emerged from our work. Patients are aware of the recommendation to reduce the risk of contagion and stringently follow them. Social distancing, handwashing with alcohol gel, wearing face masks, and avoiding close contact with other patients were perceived as safe procedures to access dental clinics. The same positive perception emerged when dentists wore all PPE and used N95 or FFP2 masks or face shields during dental treatments. The adoption of these procedures by dentists was not considered a hindrance to dental treatments. Still, it was an appropriate medical staff's behavior to assure the optimal resumption of elective dental treatments. Assurances of public health officials and institutions of the safety restart of routine dental care have also been reported as determinants in enhancing trust in this resumption [22].

About anxiety and fear of COVID-19 infection during dental treatments, we observed an increased sense of pain and nervousness during the first COVID-19 outbreak. Due to

stringent lockdowns, people closed in their homes suffered a higher level of anxiety [17], which necessarily had repercussions on behaviors and perceptions toward dental treatments. This perception should be considered as multifactorial. While a high level of dental anxiety was observed after confinement in different studies, with a prevalence ranging from 27.2 to 38.7% [36,37], it should be observed that dental treatments require patients to avoid, albeit for a short time, the use of personal protection, like masks and social distancing, tools that continue to be indicated as the first and essential barriers to prevent infection, even now that vaccines are available in many countries and large populations have already received two vaccine doses [38].

Females demonstrated a significantly higher dental anxiety and fear than males. This can be partly explained as the consequence of females' higher level of anxiety toward dental treatments [39,40] and, in the last two years, toward the COVID-19 pandemic [41]. During the COVID-19 pandemic, males were more willing to attend a dental appointment, while females were strongly prevented from going to the dentist by their fear of infection, and a natural aversion to germs made them prefer to postpone non-urgent dental appointments [21].

About the behaviors of younger people, we report different findings. In Cotrin et al., the younger showed a higher level of anxiety and depressive symptoms toward dental treatments than older people [26]. On the contrary, Martina et al. reported a higher level of fear among younger people than among people over 40 years [28]. These contrasting results should be due to the different times of observation. During the evolution of the COVID-19 pandemic, scientific evidence has proved a higher risk of infection for older people that could have translated in a significant growing attention of the adults to avoid situations that could increase the risk of contagion. Thus, dentist avoidance emerged as a protective measure during quarantine, especially for patients who perceive a high risk of infection, are afraid of COVID-19, and are older than 60 years [21].

This study has several limitations. First, significant heterogeneity among the included studies was observed, preventing us from determining a quantitative prevalence of dental anxiety associated with the COVID-19 pandemic. Second, the sample sizes varied from hundreds of patients to more than 1500. Thus, in some cases, the sample size could not be representative of the reference population. Third, some studies reported dental anxiety and fear of patients not needing dental care and who would not schedule dental care at that time, so that the lack of a concrete need could have altered the perception. Fourth, the prevalence of COVID-19 infection and mortality in the interviewed area was not considered. Finally, all included studies had a cross-sectional design and were executed by adopting a self-reported questionnaire that, in some cases, used not-validated tools for psychometric measurements.

4. Conclusions and Prospects

The dental profession has always faced complex challenges related to viral pandemics. However, as demonstrated during the COVID-19 outbreak, dentists have promptly responded to new scenarios, improving safety. According to the COVID-19 guidelines, dentists change their workflows patterns or re-configure their clinic settings to ensure safety. Many precautions consist in increasing the use of protective materials and/or in prolonging the treatment time. If, on the one hand, these behaviors assure high safety standards for patients and dental treatments, on the other hand, they increase the cost of dental treatments. To reduce the impact of such new costs, some dentists have added significant charges for PPE to their bills, especially in the case of aerosol-generating procedures. In these cases, the cost of safety is entirely absorbed by patients who may be induced to go to the dentist only in cases of urgency. On the contrary, the lack of national policies to reduce the impact of the cost related to the proper application of the recommended guidelines, also in private dental offices, can increase the pressure on dentists and may set the stage for increased errors to avoid a significant rise of patients' cost.

How to prevent such negative scenarios? These issues spotlight the need for shared national policies to reduce the negative impact of the COVID-19 crisis. As done in many countries to support national and local economies, in the new normalcy of dentistry after the COVID-19 outbreak, sharing costs among patients, dentists, and national policies may help restore the regular dental activity, helping patients not to renounce dental treatments.

In conclusion:

- The availability of vaccines for the population and the awareness of patients about recommended guidelines to prevent COVID-19 contagion support a safe and lasting resumption of routine dental activities
- Communication with patients about adopted precautions in dental clinics plays a fundamental role in reducing patients' dental anxiety and fear related to the risk of COVID-19 infection within dental clinics.
- Compliance with dentist appointments should be encouraged, helping patients re-discover the confidential relationship of trust with their dentist and return to regular dental visits.
- Identifying economic policies to support the current financial pressure due to the high cost of dental treatments may speed up the return to normalcy for dentists and patients.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/encyclopedia2010022/s1>. Table S1: Risk of bias-CHERRIES Checklist. Table S2: Risk of bias-SURge Checklist.

Author Contributions: Conceptualization, S.S. and M.L.G.; methodology, M.L.G. and E.A.; software, M.L.G. and P.B.; validation, S.S., M.L.G., E.A., and M.S.; writing—original draft preparation, M.L.G., E.A., and P.B.; writing—review and editing, S.S. and M.S.; supervision, M.L.G. and E.A.; project administration, M.L.G. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Informed Consent Statement: Not Applicable.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

DFS	Dental Fear Survey
MDAS	Modified Dental Anxiety Scale
MTurk	Mechanical Turk
NRS	Numerical Rating Scale
PPE	Personal Protective Equipment
PVD	Perceived Vulnerability to disease
TMD	Temporomandibular Disorders

References

1. American Dental Association. ADA Recommending Dentists Postpone Elective Procedures. Available online: <https://www.ada.org/en/publications/ada-news/2020-archive/march/ada-recommending-dentists-postpone-elective-procedures> (accessed on 29 October 2021).
2. Center for Disease Control and Prevention, SARS-CoV-2 and Transmission. Available online: <https://www.cdc.gov/coronavirus/2019-ncov/science/science-briefs/sars-cov-2-transmission.html> (accessed on 29 October 2021).
3. Bakaeen, L.G.; Masri, R.; AlTarawneh, S.; Garcia, L.T.; AlHadidi, A.; Khamis, A.H.; Hamdan, A.M.; Baqain, Z.H. Dentists' knowledge, attitudes, and professional behavior toward the COVID-19 pandemic: A multisite survey of dentists' perspectives. *J. Am. Dent. Assoc.* **2021**, *152*, 16–24. [CrossRef] [PubMed]
4. Beltran-Aguilar, E.; Benzian, H.; Niederman, R. Rational perspectives on risk and certainty for dentistry during the COVID-19 pandemic. *Am. J. Infect. Control.* **2021**, *49*, 131–133. [CrossRef] [PubMed]
5. Salgarello, S.; Salvadori, M.; Mazzoleni, F.; Salvalai, V.; Francinelli, J.; Bertolotti, P.; Lorenzi, D.; Audino, E.; Garo, M.L. Urgent Dental Care During Italian Lockdown: A Cross-sectional Survey. *J. Endod.* **2021**, *47*, 204–214. [CrossRef] [PubMed]

6. Ball, M.; Akintola, D.; Harrington, Z.; Djemal, S. Emergency dental care triage during the COVID-19 pandemic. *Br. Dent. J.* **2021**, *1–5*. [\[CrossRef\]](#) [\[PubMed\]](#)
7. Eggmann, F.; Haschemi, A.A.; Doukoudis, D.; Filippi, A.; Verna, C.; Walter, C.; Weiger, R.; Zitzmann, N.U.; Bornstein, M.M. Impact of the COVID-19 pandemic on urgent dental care delivery in a Swiss university center for dental medicine. *Clin. Oral Investig.* **2021**, *25*, 5711–5721. [\[CrossRef\]](#) [\[PubMed\]](#)
8. Ostrc, T.; Pavlovic, K.; Fidler, A. Urgent dental care on a national level during the COVID-19 epidemic. *Clin. Exp. Dent. Res.* **2021**, *7*, 271–278. [\[CrossRef\]](#) [\[PubMed\]](#)
9. Pajpani, M.; Patel, K.; Bendkowski, A.; Stenhouse, P. Rapid response: Activity from a hospital based Urgent Dental Care Centre during the COVID-19 pandemic. *Br. J. Oral Maxillofac. Surg.* **2020**, *58*, e98–e103. [\[CrossRef\]](#)
10. Walter, E.; von Bronk, L.; Hickel, R.; Huth, K.C. Impact of COVID-19 on Dental Care during a National Lockdown: A Retrospective Observational Study. *Int. J. Environ. Res. Public Health* **2021**, *18*, 7963. [\[CrossRef\]](#)
11. Salgarello, S.; Salvadori, M.; Mazzoleni, F.; Francinelli, J.; Bertolotti, P.; Audino, E.; Garo, M.L. The New Normalcy in Dentistry after the COVID-19 Pandemic: An Italian Cross-Sectional Survey. *Dent. J.* **2021**, *9*, 86. [\[CrossRef\]](#)
12. American Dental Association. ADA ‘Strongly’ Encouraging Dental Professionals to be Vaccinated for COVID-19. Available online: <https://www.ada.org/en/publications/ada-news/2021-archive/july/ada-strongly-encouraging-dental-professionals-to-be-vaccinated> (accessed on 29 October 2021).
13. Zigron, A.; Dror, A.A.; Morozov, N.G.; Shani, T.; Haj Khalil, T.; Eisenbach, N.; Rayan, D.; Daoud, A.; Kablan, F.; Marei, H.; et al. COVID-19 Vaccine Acceptance Among Dental Professionals Based on Employment Status During the Pandemic. *Front Med.* **2021**, *8*, 618403. [\[CrossRef\]](#)
14. Menhadji, P.; Patel, R.; Asimakopoulou, K.; Quinn, B.; Khoshkhounejad, G.; Pasha, P.; Garcia Sanchez, R.; Ide, M.; Kalsi, P.; Nibali, L. Patients’ and dentists’ perceptions of tele-dentistry at the time of COVID-19. A questionnaire-based study. *J. Dent.* **2021**, *113*, 103782. [\[CrossRef\]](#) [\[PubMed\]](#)
15. Yavan, M.A. Effects of the COVID-19 pandemic on new patient visits for orthodontic treatment: A comparison of 2020 and the previous 3 years. *J. World Fed. Orthod.* **2021**, *10*, 127–131. [\[CrossRef\]](#) [\[PubMed\]](#)
16. Guo, H.; Zhou, Y.; Liu, X.; Tan, J. The impact of the COVID-19 epidemic on the utilization of emergency dental services. *J. Dent. Sci.* **2020**, *15*, 564–567. [\[CrossRef\]](#) [\[PubMed\]](#)
17. Huang, Y.; Zhao, N. Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: A web-based cross-sectional survey. *Psychiatry Res.* **2020**, *288*, 112954. [\[CrossRef\]](#) [\[PubMed\]](#)
18. Lei, L.; Huang, X.; Zhang, S.; Yang, J.; Yang, L.; Xu, M. Comparison of Prevalence and Associated Factors of Anxiety and Depression Among People Affected by versus People Unaffected by Quarantine During the COVID-19 Epidemic in Southwestern China. *Med. Sci. Monit.* **2020**, *26*, e924609. [\[CrossRef\]](#)
19. Moher, D.; Liberati, A.; Tetzlaff, J.; Altman, D.G.; Group, P. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med.* **2009**, *6*, e1000097. [\[CrossRef\]](#)
20. Weir, K.R.; Ailabouni, N.J.; Schneider, C.R.; Hilmer, S.N.; Reeve, E. Considerations for systematic reviews of quantitative surveys: Learnings from a systematic review of the patients’ Attitudes Towards Deprescribing questionnaire. *Res. Soc. Adm. Pharm.* **2021**, *18*, 2345–2349. [\[CrossRef\]](#)
21. Gonzalez-Olmo, M.J.; Delgado-Ramos, B.; Ortega-Martinez, A.R.; Romero-Maroto, M.; Carrillo-Diaz, M. Fear of COVID-19 in Madrid. Will patients avoid dental care? *Int. Dent. J.* **2022**, *72*, 76–82. [\[CrossRef\]](#)
22. Moffat, R.C.; Yentes, C.T.; Crookston, B.T.; West, J.H. Patient Perceptions about Professional Dental Services during the COVID-19 Pandemic. *JDR Clin. Trans. Res.* **2021**, *6*, 15–23. [\[CrossRef\]](#)
23. Nazir, M.; Almulhim, K.S.; AlDaamah, Z.; Bubshait, S.; Sallout, M.; AlGhamdi, S.; Alhumaid, J. Dental Fear and Patient Preference for Emergency Dental Treatment Among Adults in COVID-19 Quarantine Centers in Dammam, Saudi Arabia. *Patient Prefer. Adherence* **2021**, *15*, 1707–1715. [\[CrossRef\]](#)
24. Quan, S.; Guo, Y.; Zhou, J.; Zhang, G.; Xing, K.; Mei, H.; Li, J. Orthodontic emergencies and mental state of Chinese orthodontic patients during the COVID-19 pandemic. *BMC Oral Health* **2021**, *21*, 477. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Arqub, S.A.; Voldman, R.; Ahmida, A.; Kuo, C.L.; Godoy, L.D.C.; Nasrawi, Y.; Al-Khateeb, S.N.; Uribe, F. Patients’ perceptions of orthodontic treatment experiences during COVID-19: A cross-sectional study. *Prog. Orthod.* **2021**, *22*, 17. [\[CrossRef\]](#) [\[PubMed\]](#)
26. Cotrin, P.; Peloso, R.M.; Oliveira, R.C.; de Oliveira, R.C.G.; Pini, N.I.P.; Valarelli, F.P.; Freitas, K.M.S. Impact of coronavirus pandemic in appointments and anxiety/concerns of patients regarding orthodontic treatment. *Orthod. Craniofac. Res.* **2020**, *23*, 455–461. [\[CrossRef\]](#) [\[PubMed\]](#)
27. KaragÖzoğlu, İ.; Parlar ÖZ, Ö. Investigation of the patients’ perception on dental treatment and their anxiety levels during the COVID-19 pandemic process. *J. Health Sci. Med.* **2021**, *4*, 710–715. [\[CrossRef\]](#)
28. Martina, S.; Amato, A.; Faccioni, P.; Iandolo, A.; Amato, M.; Rongo, R. The perception of COVID-19 among Italian dental patients: An orthodontic point of view. *Prog. Orthod.* **2021**, *22*, 11. [\[CrossRef\]](#)
29. Peloso, R.M.; Pini, N.I.P.; Sundfeld Neto, D.; Mori, A.A.; Oliveira, R.C.G.; Valarelli, F.P.; Freitas, K.M.S. How does the quarantine resulting from COVID-19 impact dental appointments and patient anxiety levels? *Braz. Oral Res.* **2020**, *34*, e84. [\[CrossRef\]](#)
30. Umeh, O.D.; Utomi, I.L.; Isiekwe, I.G.; Aladenika, E.T. Impact of the coronavirus disease 2019 pandemic on orthodontic patients and their attitude to orthodontic treatment. *Am. J. Orthod. Dentofac. Orthop.* **2021**, *159*, e399–e409. [\[CrossRef\]](#)

31. Ather, A.; Patel, B.; Ruparel, N.B.; Diogenes, A.; Hargreaves, K.M. Coronavirus Disease 19 (COVID-19): Implications for Clinical Dental Care. *J. Endod.* **2020**, *46*, 584–595. [[CrossRef](#)]
32. Lanmark 360. Survey of COVID-19 Dental Patients' Perceptions and Attitudes Toward Returning to Dental Practices. Available online: <https://lanmark360.com/research/Lanmark360-Dental-Patients-COVID-19-Survey-Results.pdf> (accessed on 29 October 2021).
33. Innes, N.; Johnson, I.G.; Al-Yaseen, W.; Harris, R.; Jones, R.; Kc, S.; McGregor, S.; Robertson, M.; Wade, W.G.; Gallagher, J.E. A systematic review of droplet and aerosol generation in dentistry. *J. Dent.* **2021**, *105*, 103556. [[CrossRef](#)]
34. Peng, X.; Xu, X.; Li, Y.; Cheng, L.; Zhou, X.; Ren, B. Transmission routes of 2019-nCoV and controls in dental practice. *Int. J. Oral Sci.* **2020**, *12*, 9. [[CrossRef](#)]
35. Zimmermann, M.; Nkenke, E. Approaches to the management of patients in oral and maxillofacial surgery during COVID-19 pandemic. *J. Craniomaxillofac. Surg.* **2020**, *48*, 521–526. [[CrossRef](#)] [[PubMed](#)]
36. Mazza, C.; Ricci, E.; Biondi, S.; Colasanti, M.; Ferracuti, S.; Napoli, C.; Roma, P. A Nationwide Survey of Psychological Distress among Italian People during the COVID-19 Pandemic: Immediate Psychological Responses and Associated Factors. *Int. J. Environ. Res. Public Health* **2020**, *17*, 3165. [[CrossRef](#)] [[PubMed](#)]
37. Roy, D.; Tripathy, S.; Kar, S.K.; Sharma, N.; Verma, S.K.; Kaushal, V. Study of knowledge, attitude, anxiety & perceived mental healthcare need in Indian population during COVID-19 pandemic. *Asian J. Psychiatr.* **2020**, *51*, 102083. [[CrossRef](#)] [[PubMed](#)]
38. Our World in Data, Coronavirus (COVID-19) Vaccinations. Available online: <https://ourworldindata.org/covid-vaccinations> (accessed on 15 October 2021).
39. Jeddy, N.; Nithya, S.; Radhika, T.; Jeddy, N. Dental anxiety and influencing factors: A cross-sectional questionnaire-based survey. *Indian J. Dent. Res.* **2018**, *29*, 10–15. [[CrossRef](#)]
40. Lin, C.S.; Wu, S.Y.; Yi, C.A. Association between Anxiety and Pain in Dental Treatment: A Systematic Review and Meta-analysis. *J. Dent. Res.* **2017**, *96*, 153–162. [[CrossRef](#)]
41. Wang, C.; Pan, R.; Wan, X.; Tan, Y.; Xu, L.; Ho, C.S.; Ho, R.C. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *Int. J. Environ. Res. Public Health* **2020**, *17*, 1729. [[CrossRef](#)]