## SUPPLEMENTARY FILES

**Supplementary Table S1.** Relevant characteristics of the study sample.

| Variables | Study sample (N=65) |
|---|---|
| **Presence of symptoms** | 62 (95.4%) |
| **Subfebrility** | 28 (43.1%) |
| **Febrility** | 29 (44.6%) |
| **Cough** | 36 (55.4%) |
| **Dyspnea** | 15 (23.1%) |
| **Headache** | 16 (24.6%) |
| **Sore throat** | 13 (20.0%) |
| **Malaise** | 49 (75.4%) |
| **Rhinorrhea** | 3 (4.6%) |
| **Anosmia** | 4 (6.2%) |
| **Diarrhea** | 13 (20.0%) |
| **Venous thromboembolism** | 1 (1.5%) |
| **X-ray confirmed pneumonia** | 46 (70.8%) |
| **X-ray confirmed pleural effusion** | 2 (3.1%) |
| **Positive urine culture** | 9 (13.8%) |
| **Positive blood culture** | 4 (6.2%) |

**Abbreviations:** None.

**Supplementary Table S2.** ROC curve analysis of ImAgeS risk score with 30-day post-discharge mortality.

| Variables | C-statistic (95% CI) | p-value | Sensitivity / specificity | Youden index |
|---|---|---|---|---|
| **ImAgeS risk score** | 0.679 (0.519-0.840) | 0.027 | 58.0/85.0 | 69.15 |

**Note:** ImAgeS – novel clinical risk score.

**Supplementary Table S3.** Association of different patient characteristics with 30-day post-discharge mortality.

| Variables | aOR (95% CI)* |
|---|---|
| **Age** | 1.04 (0.95-1.14), p=0.440 |
| **Male sex** | 1.66 (0.31-9.06), p=0.558 |
| **Diabetes mellitus** | 3.16 (0.64-15.65), p=0.159 |
| **Arterial hypertension** | 0.64 (0.10-4.17), p=0.640 |
| **Chronic heart failure** | 6.70 (0.75-60.10), p=0.089 |
| **Transplant duration (years)** | 1.10 (0.98-1.23), p=0.092 |

*Binomial multivariable logistic regression model.
**Abbreviations:** aOR – adjusted odds ratios; CI – confidence intervals.

**Supplementary Figure S1.** Flow diagram.

**Note:** ImAgeS – novel clinical risk score.

**Legend:** COVID-19 – Coronavirus disease 2019.

**Supplementary Figure S2.** Calibration plot

**Note:** ImAgeS – novel clinical risk score.