

# Technology for the Production of Somatic Seeds Useful for the Storage of Valuable Genotypes of *Salvia officinalis* †

Aleksandra Deja 

Biotechnology Department, Institute of Natural Fibres & Medicinal Plants National Research Institute, Wojska Polskiego str. 71b, 60-630 Poznan, Poland; aleksandra.deja@iwnirz.pl

† Presented at the 2nd International Electronic Conference on Plant Sciences—10th Anniversary of Journal Plants, 1–15 December 2021; Available online: <https://iecps2021.sciforum.net/>.

**Abstract:** Current research has attempted to produce and convert somatic seeds that would enable the long-term preservation of valuable sage genotypes. The creation of artificial seeds consists of placing explants capable of regenerating into plants in a protective casing. The method of obtaining somatic seeds used in this study allowed us to obtain a high level of conversion of seeds into plants using apical buds (85%), which was slightly lower in the case of axillary buds (62.5%). The results concerning the formation and conversion of somatic seeds allowed us to obtain a high level of plant viability, which may prove the usefulness of the method of storing valuable *Salvia officinalis* genotypes.

**Keywords:** somatic seeds; *Salvia officinalis*; in vitro cultures

## 1. Introduction

Biotechnological methods are being increasingly used to protect biodiversity. The protection of genetic resources is necessary not only for future generations, but also for obtaining new valuable varieties. Currently, gene banks and in vitro cultures enable medium- and long-term storage of plant material. Somatic seeds are also used to preserve genetic resources in gene banks. Artificial seeds contain any regenerative plant fragment in a protective bead in preparation for long-term storage or for commercial use. The definition of artificial seeds implies their use in the storage and mass multiplication of genetically identical and healthy plants. The aim of this research was to produce and convert somatic seeds that would enable the long- or medium-term storage of valuable genotypes of *Salvia officinalis* (sage). The development of a method to store valuable genotypes is very important for sage, which is an insect-pollinated plant and usually blooms in the second year of cultivation. Moreover, somatic seeds can ensure the storage of plant line with a guarantee of genetic homogeneity and propagation of variable genotypes such as transgenic plants. In this context, research into the improvement of biotechnological methods for the production of somatic sage seeds is highly relevant.

## 2. Materials and Methods

### 2.1. Plant Material and Production of Somatic Seeds

The apical and axillary buds were used as explants for the production of somatic seeds. The explants ( $\pm 1.5$  cm) were dissected from the 2-week-old multi-shoot cultures, then encapsulated with 1.2% sodium alginate solution, and then dripped into the 200 mM  $\text{CaCl}_2$  solution. The occurring reaction of exchanging calcium ions with sodium ions leads to the formation of a beads with a hard polymer coat. As a result of mixing, the capsules take a spherical form containing a single explant. The somatic seeds were stored for 1 month at 4 °C.

The somatic seeds were converted on MS medium containing 0.3 mg L<sup>-1</sup> BAP. In the study, we used 20 genotypes in 4 replicates. After 2 and 4 weeks, the following data



**Citation:** Deja, A. Technology for the Production of Somatic Seeds Useful for the Storage of Valuable Genotypes of *Salvia officinalis*. *Biol. Life Sci. Forum* **2022**, *11*, 43. <https://doi.org/10.3390/IECPS2021-12057>

Academic Editor: Feibo Wu

Published: 7 December 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).


were recorded: frequency of shoot formation, morphological data (color, vitrification, necrose etc.). The regenerated shoots were rooted on  $\frac{1}{2}$  MS medium containing  $0.3 \text{ mg L}^{-1}$  IAA in order to obtain the acclimatized plantlets. Conversion and rooting proceeded at  $25 \pm 2 \text{ }^\circ\text{C}$  with a 16 h photoperiod under a photosynthetic flux of  $120 \mu \text{ mol m}^2 \text{ s}^{-1}$  (daylight fluorescent tubes). Experiments were performed for both types of explants in 4 replicates of 20 seeds.

## 2.2. Statistical Analysis

The results were expressed as the percentage (%) of germinated seeds of three replicates each of 20 seeds. The data were statistically analyzed using one-way analysis of variance (ANOVA) and the statistical significance was determined. The data was compared by the least significant difference ( $p \leq 0.05$ ) test using GenStat15 System.

## 3. Results

As the result of the conducted experiments, the conversion ability of somatic seeds of valuable *S. officinalis* genotypes was verified. Somatic seeds were produced using both the apical and axillary buds (Figure 1).



**Figure 1.** Somatic seeds of *Salvia officinalis*: (a) using apical buds; (b) using axillary buds; (c) after production in sterile conditions; (d) single somatic seed.

The obtained results of the observation during cultivation on MS medium containing  $0.3 \text{ mg L}^{-1}$  BAP are presented in Table 1.

The method of obtaining somatic seeds used in this study allowed to obtain a high level of conversion of seeds into plants using apical buds (85%). The slightly lower percentage of regenerated plants was obtained for axillary buds (62.5%). However, the statistical significance of seed viability in the ANOVA analysis of variance was at the level of  $p < 0.01$  (Table 2). Therefore, it can be assumed that the usefulness of the developed method applies to both initial explants.

For complete regeneration of plants, shoots were rooted in  $\frac{1}{2}$  MS medium containing  $0.3 \text{ mg L}^{-1}$  IAA. The results indicate that four explants out of five developed roots ( $4.067 \pm 0.3914$ ). Regenerated sage plants were characterized by proper growth and development at each stage of culture (Figure 2).

**Table 1.** In vitro cultures of somatic seed of *Salvia officinalis*.

Somatic Seed	Explant			Observation		
	W	B	L.S.D <sub>0.05</sub>	1	2	L.S.D <sub>0.05</sub>
	Mean Values (%)			Mean Values (%)		
Green	77.5 <sup>a</sup>	75.0 <sup>a</sup>	17.51	100.0 <sup>a</sup>	52.5 <sup>b</sup>	17.51
Brown-green	22.5 <sup>a</sup>	25.0 <sup>a</sup>	17.51	0.0 <sup>b</sup>	47.5 <sup>a</sup>	17.51
Brown	0.0	0.0	-	0.0	0.0	-
Growing	85.0 <sup>a</sup>	62.5 <sup>b</sup>	12.18	47.5 <sup>b</sup>	100.0 <sup>a</sup>	12.18
Non-vitality	15.0 <sup>b</sup>	37.5 <sup>a</sup>	12.18	52.5 <sup>a</sup>	0 <sup>b</sup>	12.18
Necrosed	0.0	0.0	-	0.0	0.0	-
Vittificated	30.0 <sup>a</sup>	32.5 <sup>a</sup>	13.71	0.0 <sup>b</sup>	62.5 <sup>a</sup>	13.71
Contaminated	0.0	0.0	-	0.0	0.0	-

W—apical; B—axillary. Values represent percentage (of germinated seeds) of three replicates each of 20 seeds. Different lowercase letters (a, b) in the same column indicate a significant difference at  $p \leq 0.05$ .

**Table 2.** ANOVA analysis of the traits observed in in vitro cultures of somatic seeds.

Source of Variation	D.F.	Somatic Seed					
		Green	Brown-Green	Brown	Growing	Non-Vitality	Vittificated
Observation	1	9025 ***	9025 ***	0	11,025 ***	11,025 ***	15,625 ***
Explant	1	25	25	0	2025 **	2025 **	25
Residual	12	258.3	258.3	0	125	125	158.3

\*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$ .



**Figure 2.** In vitro culture of *Salvia officinalis*: (a) somatic seeds placed on MS medium containing  $0.3 \text{ mg L}^{-1}$  BAP; (b) shoots regenerated from somatic seeds; (c) multi-shoot cultures; (d) regenerated plants on  $\frac{1}{2}$  MS medium containing  $0.3 \text{ mg L}^{-1}$  IAA.

#### 4. Discussion

Since Kitto and Janick [1] began their initial research on artificial seeds by working on carrots, there has been continual interest in this topic. This led to a general definition of an artificial seed, also called a synthetic seed or synseed, seed analog or manufactured seed, which includes a range of plant structures, including somatic embryos, buds, shoots or other meristematic tissues inside a coating that can be sown in the same way as a conventional seed to produce a new plant [2]. The results obtained in this study have shown that somatic seeds of sage were capable of conversion and regeneration. The developed protocol was effective and enabled the complete regeneration of plants from apical and axillary buds. The use of this type of explant for the production of artificial seeds is increasingly becoming an alternative to the use of somatic embryos [3]. There is only one report devoted to the production of artificial seeds of *S. officinalis*, which confirms the usefulness of this

technology for high medical- and commercial-value plants such as sage [4]. Similar results regarding the formation of shoots from somatic seeds were obtained. However, in the present study, MS medium, sucrose and gibberellic acid were not used as the components of the encapsulation gel, which significantly simplified the procedure. Therefore, our simplified procedure for producing artificial seeds and high conversion and regeneration efficiency of somatic seeds is a novelty in the study on the application of the technique of medium- and long-term storage genotypes of *Salvia officinalis*.

The potential benefits of using somatic seeds include rapid multiplication and long-term storage of a given plant line with a guarantee of genetic homogeneity. In addition, somatic seeds can provide for the delivery and propagation of variable genotypes such as transgenic plants with high genetic instability. Often, artificial seeds can be the only way to reproduce genetically modified plants [5]. In this context, the method obtained in our own work could be developed and optimized in the future for the storage of valuable sage genotypes after genetic transformation [6].

**Funding:** The presented work was financially supported by The Ministry of Science and Higher Education in Poland, Technological Initiative I, Grant number MNiSzW IT-1-13910: "Obtaining of salvia (*Salvia officinalis*) with anti carries properties".

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Conflicts of Interest:** The author declares no conflict of interest.

## References

1. Kitto, S.; Janick, J. Hardening treatments increase survival of synthetically-coated asexual embryos of carrot. *J. Am. Soc. Hortic. Sci.* **1985**, *110*, 283–286.
2. Pond, S.; Cameron, S. Artificial Seeds. In *Encyclopedia of Applied Plant Sciences*, 2nd ed.; Academic Press, Elsevier Ltd.: Amsterdam, The Netherlands, 2017; Volume 2, pp. 419–427. ISBN 978-0-12-394808-3.
3. Makowczyńska, J.; Andrzejewska-Golec, E. Somatic seeds of *Plantago asiatica* L. *Acta Soc. Bot. Pol.* **2006**, *75*, 17–21.
4. Grzegorzczak, I.; Wysokińska, H. A protocol for synthetic seeds from *Salvia officinalis* L. shoot tips. *Acta Biol. Crac. Ser. Bot.* **2011**, *53*, 80–85. [[CrossRef](#)]
5. Kopcewicz, J.; Lewak, S. *Fizjologia Roślin*; Wydawnictwo Naukowe PWN Warszawa: Wrocław, Poland, 2002; pp. 194–195. ISBN 9788301145491.
6. Luwańska, A.; Wielgus, K.; Seidler-Łożykowska, K.; Lipiński, D.; Słomski, R. Evaluation of *Agrobacterium tumefaciens* Usefulness for the Transformation of Sage (*Salvia officinalis* L.). In *Transgenesis and Secondary Metabolism*; Jha, S., Ed.; Springer International Publishing: Cham, Switzerland, 2017; pp. 153–176. [[CrossRef](#)]