



Abstract Modeling and Simulation of Magnetoliposome Formation by Encapsulation of Core-Shell, Magnetite-Chitosan Nanoparticles in Liposomes Enabled by a Low-Cost Microfluidic System [†]

Andres Mantilla-Orozco ¹, Cristian Felipe Rodriguez ¹, Isabella Quiroz ¹, Juan S. Bermudez ¹, Daniel Felipe Forero ¹, Maria Camila Monsalve ¹, Carolina Muñoz-Camargo ¹, Luis H. Reyes ², Johann F. Osma ³, Valentina Quezada ^{1,*} and Juan C. Cruz ^{1,*}

- ¹ Department of Biomedical Engineering, School of Engineering, Universidad de Los Andes, Carrera 1 No. 18A-12, Bogota 111711, Colombia
- ² Department of Chemical and Food Engineering, School of Engineering, Universidad de Los Andes, Carrera 1 No. 18A-12, Bogota 111711, Colombia
- ³ Department of Electrical and Electronic Engineering, School of Engineering, Universidad de Los Andes, Carrera 1 No. 18A-12, Bogota 111711, Colombia
- * Correspondence: v.quezada@uniandes.edu.co (V.Q.); jc.cruz@uniandes.edu.co (J.C.C.); Tel.: +57-1-3394949 (ext. 1789) (J.C.C.)
- + Presented at the 2nd International Electronic Conference on Biomolecules: Biomacromolecules and the Modern World Challenges, 1–15 November 2022; Available online: https://iecbm2022.sciforum.net/.

Abstract: Research in nanostructured materials has led to the development of different applications of relevance in the fields of medicine and biomedical engineering. In this regard, the field of drug delivery has probably benefited the most due to the possibility to engineer vehicles of high potency and increased activity and selectivity toward selected intracellular targets. Such vehicles can therefore potentially address one of the major cornerstones of modern pharmacology, which is increasing the bioavailability of drugs of low permeability. Our research group has developed cell-penetration nanobioconjugates by interfacing several nanomaterials (e.g., chitosan, gelatin nanoparticles, graphene oxide, and magnetite) with translocating peptides. The obtained nanobioconjugates have demonstrated facilitated cell internalization and endosomal escape abilities. To improve cell penetration even further, we encapsulated the magnetite-based nanobioconjugates into liposomes (to form magnetoliposomes) with very appealing results. Our plan is to expand the available nanoplatforms by combining the attributes of magnetite and polymeric nanoparticles through a core-shell system comprised of magnetite (core) and chitosan (shell). The encapsulation process has been successfully accomplished with the aid of passive micromixers with different channel geometries to favor intimate contact between the dispersed phase (nanoparticles) and the continuous phase (phospholipid solution). To model the encapsulation process, we implemented an Eulerian simulation in the software COMSOL Multiphysics[®] 6.0 (COMSOL Inc, Stockholm, Sweden) where mixing required the Navier-Stokes equations as governing equations of momentum transport, turbulence, eddy viscosity, and damping functions to approximate turbulence using the κ - ϵ turbulence model near the walls. The simulation was conducted for the different geometries (i.e., SARS, chambers, and serpentine) and for Reynolds numbers ranging from 0.2 to 10 Also, we tested a low Reynolds turbulent model using the κ - ε model given in the Euler-Euler module. The Euler-Euler approach showed that the encapsulation reaches higher encapsulation efficiency (EE%) values compared with the previously implemented mixture model. Our encapsulation results indicate that including the κ - ε turbulence model with a low Reynolds turbulence model near the walls provides a higher agreement between in-silico and experimental approaches. Future work will be dedicated to evaluating the performance of our previously tested magnetophoretic separators with the newly developed encapsulates, to assure sufficient purity for further biocompatibility testing.

Keywords: Microfluidics; MNP's; liposome; drug delivery systems; Euler-Euler approach



Citation: Mantilla-Orozco, A.; Rodriguez, C.F.; Quiroz, I.; Bermudez, J.S.; Forero, D.F.; Monsalve, M.C.; Muñoz-Camargo, C.; Reyes, L.H.; Osma, J.F.; Quezada, V.; et al. Modeling and Simulation of Magnetoliposome Formation by Encapsulation of Core-Shell, Magnetite-Chitosan Nanoparticles in Liposomes Enabled by a Low-Cost Microfluidic System. *Biol. Life Sci. Forum* 2022, 20, 29. https://doi.org/ 10.3390/IECBM2022-13398

Academic Editor: Robert Stahelin

Published: 1 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). **Supplementary Materials:** The presentation material of this work is available online at https://www.mdpi.com/article/10.3390/IECBM2022-13398/s1.

Author Contributions: Conceptualization, J.C.C., L.H.R., J.F.O. and C.M.-C.; methodology, data curation and data analysis A.M.-O., C.F.R., and J.C.C.; software, C.F.R., A.M.-O., J.S.B., and I.Q.; validation, J.C.C., L.H.R., J.F.O., C.M.-C. and V.Q.; formal analysis and investigation, A.M.-O., C.F.R., and I.Q.; resources, A.M.-O., C.F.R., V.Q., J.F.O., L.H.R. and J.C.C.; writing—original draft preparation, A.M.-O., C.F.R., I.Q., D.F.F., and M.C.M.; writing— review and editing, J.C.C., L.H.R., J.F.O., C.M.-C. and V.Q.; visualization, A.M.-O., C.F.R., and I.Q.; supervision, J.C.C., L.H.R., J.F.O., C.M.-C. and V.Q.; project administration, J.F.O., L.H.R. and J.C.C.; funding acquisition, J.F.O. and L.H.R. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Minciencias, grant ID 120484467244.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.