



Bioaccumulation and Human Health Risk of Heavy Metals from Pesticides in Some Crops Grown in Plateau State, Nigeria [†]

Bawa Usman ^{1,*} , AbdulHameed Ahmad ² , Nayaya A. Jibrin ², Ezra A. Gaya ² and Maryam Jibrin ³

¹ Department of Biological Sciences, Bayero University, Kano, P.M.B., Kano 3011, Nigeria

² Department of Applied Ecology, Abubakar Tafawa Balewa University, P.M.B., Bauchi 0248, Nigeria; gesunnam.aa@gmail.com (A.A.); nayayaaj2001@yahoo.com (N.A.J.); agezra2006@gmail.com (E.A.G.)

³ Department of Biology Education, Aminu Saleh College of Education, P.M.B., 944 Azare, Bauchi 751101, Nigeria; mayan.adamu@gmail.com

* Correspondence: bawa.usman@yahoo.com

[†] Presented at the 1st International Electronic Conference on Plant Science, 1–15 December 2020; Available online: <https://iecps2020.sciforum.net/>.

Abstract: The health risk assessment of heavy metals as the source of metal contamination in food crops fumigated exclusively with pesticides is mostly overlooked. This study determined the concentrations of heavy metals (Cd, Pb, Cr, Cu, and Zn) in some food crops fumigated with pesticides and their health risk to humans. The mean concentrations of heavy metals in different parts of the studied crops ranged from 0.12 to 2.03 for Zn, 1.73 to 23.34 for Pb, 1.60 to 1150.50 for Cu, 0.67 to 19.50 for Cr, and 0.09 to 6.14 mg/kg for Cd. The concentrations of Cd, Pb, and Cr in the investigated crops were above the WHO (2011) permissible limits and in decreasing trend of Cu > Pb > Cr > Cd > Zn. The bioaccumulation factor (BAF) > 1 values for Cd, Pb, and Zn and the BAF value were highest for copper (141.75) in *Oryza sativa*. Pollution indices showed all crops were contaminated with Cd, Pb, and Cr and were likely to pose a potential health risk to humans. The estimated daily intake of Cd and Pb from all the studied crops exceeded the USEPA (2006) oral reference dose daily limit. A hazard quotient > 1 was observed only from the consumption of *Oryza sativa* (3.504) for Cu and could likely cause potential health risk in human. The hazard index indicated a health risk through the consumption of *Oryza sativa* (4.666), *Zea mays* (1.475), and *Capsicum annum* (1.132) for all the studied metals. Therefore, there is a need for regular screening and monitoring of heavy metals from pesticides sources in food crops.

Keywords: bioaccumulation factor; hazard quotient; hazard index; heavy metals; pesticides



Citation: Usman, B.; Ahmad, A.; Jibrin, N.A.; Gaya, E.A.; Jibrin, M. Bioaccumulation and Human Health Risk of Heavy Metals from Pesticides in Some Crops Grown in Plateau State, Nigeria. *Biol. Life Sci. Forum* **2021**, *4*, 12. <https://doi.org/10.3390/IECPS2020-08737>

Academic Editor: Yoselin Benitez-Alfonso

Published: 1 December 2020

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Pesticides are extensively employed in agriculture to kill pests or unwanted organisms that may reduce crop yield and increase agricultural production [1]. Farmers in northern Nigeria have largely depended on pesticides for the control of pests, weeds, and diseases [2]. This has led to a proliferation in the importation of new pesticide products into Nigeria whose chemical contents are not known or are mostly concealed by the manufacturer [1]. The use of pesticides has been on the increase and these pesticides have been shown to contain heavy metals [1,3]. However, despite the banning of heavy metals in pesticides globally, a recent study revealed the presence of heavy metals in pesticides at levels above the recommended dilution rate in Europe [4]. Soil–plant heavy metal transfer is the main pathway for pollutants to enter the human body through the food chain.

There is a paucity of studies on heavy metal contamination of food crops exclusively fumigated with pesticides and the health risk to humans. Therefore this study was designed with the aim of determining the concentrations of heavy metals (Cd, Pb, Cr, Cu, and Zn) from pesticides in some crops and soils in Jos, Plateau State, Nigeria and their associated human health risk.

2. Materials and Methods

Samples of leaves, stems, roots, and fruits of tomato, pepper, onion, cabbage, carrot, cucumber, spinach, lettuce, and maize and corresponding soils, were collected from Naraguta Farm (A) in Plateau State, Nigeria (N 09°58.586, E 008°53.820) and Naraguta Farm B (N 09°58.562, E 008°53.230). Soils collected from some locations outside the agricultural farms that had had no pesticides application were used as controls. All samples were collected in a clean brown envelope, labelled, and transported to the ATBU Biology Laboratory and analyzed for Cr, Cu, Cd, Zn, and Pb using an atomic absorption spectrophotometer. Health risk assessment [5–7], hazard quotient [8], hazard index [8], estimated daily intake [7,8], and pollution index [8,9] were determined and results statistically analyzed by SPSS version 8.1. and two-way analysis of variance.

3. Results

Heavy Metals in the Plants and Soil and Their Factors

There was significant variation ($p < 0.05$) in the concentration of heavy metals in different parts of most of the studied crops (Tables 1 and 2). The trend of heavy metals in the studied crops was in the decreasing order of $\text{Cu} > \text{Pb} > \text{Cr} > \text{Cd} > \text{Zn}$ (Tables 1 and 2). Cadmium, chromium, and lead concentrations in the all the studied crops were above the permissible limits except in *Allium cepa* (root, leaf, and bulb), *Daucus carota* (root and stem), *Cucumis sativus* (fruit), and *Lactuca sativa* (root, leaf) (Tables 1 and 2). The concentrations of zinc in all the investigated crops were below the permissible limit. Copper was also below the permissible limit except in *Cucumis sativus* (stem, leaf, and fruit), *Zea mays* (root, leaf, and fruit), and *Oryza sativa* (root, stem, and fruit) (Tables 1 and 2).

Table 1. Mean concentration of heavy metals in crops grown in Plateau State (2018).

Sampling Site	Name of Sample	Botanical Name	Hausa Name	Heavy Metals mg/kg				
				Cd	Pb	Cr	Cu	Zn
Jos	Tomato	<i>Solanum lycopersicum</i>	Tomatur					
	Root			4.47 ^a	2.32 ^a	10.17 ^{ab}	28.37 ^b	1.02 ^c
	Stem			5.66 ^b	4.48 ^a	7.58 ^a	18.94 ^a	0.15 ^a
	Leaf			5.14 ^{ab}	3.26 ^a	9.83 ^{ab}	39.25 ^c	0.47 ^b
	Fruit			5.08 ^{ab}	1.73 ^a	10.92 ^b	35.26 ^{bc}	1.40 ^d
	Pepper	<i>Capsicum annum</i>	Attarugu					
	Root			3.11 ^a	14.88 ^b	2.67 ^a	12.28 ^a	1.72 ^a
	Stem			3.25 ^a	15.97 ^b	5.58 ^b	18.44 ^{ab}	0.12 ^a
	Leaf			3.76 ^a	9.51 ^a	4.25 ^b	20.71 ^{ab}	1.01 ^a
	Fruit			3.33 ^a	21.25 ^b	5.00 ^b	25.55 ^c	0.97 ^a
	Onion	<i>Allium cepa</i>	Albasa					
	Root			2.58 ^a	14.98 ^a	1.67 ^a	6.30 ^a	1.90 ^b
	Stem			ND	ND	ND	ND	ND
	Leaf			3.95 ^c	18.60 ^a	1.83 ^a	12.42 ^b	1.42 ^{ab}
	Bulb			2.72 ^a	17.25 ^a	2.00 ^a	6.67 ^a	0.58 ^a
	Carrot	<i>Daucus carota</i>	Karas					
	Root			ND	ND	ND	ND	ND
	Stem			4.87 ^a	18.33 ^b	0.67 ^a	12.08 ^b	2.03 ^b
	Leaf			4.55 ^a	19.78 ^b	1.17 ^a	23.43 ^c	0.72 ^a
	Fruit			4.63 ^a	3.77 ^a	3.00 ^b	1.60 ^a	ND
	Spinach	<i>Spinacia oleracea</i>	Alayyaho					
	Root			3.57 ^a	14.91 ^a	3.58 ^a	12.21 ^b	ND
	Stem			3.91 ^a	14.20 ^a	3.75 ^{ab}	10.68 ^a	0.84
	Leaf			3.21 ^a	16.12 ^a	4.17 ^c	15.68 ^c	ND
Safe limits ^a				0.2	0.3	2.3	40	60

^a Source: FAO/WHO (2001). Mean followed with same letter across the column are not significantly different ($p > 0.05$).

Table 2. Mean concentration of heavy metals in crops grown in Plateau State (2018).

Sampling Site	Name of Sample	Botanical Name	Hausa Name	Heavy Metals mg/kg				
				Cd	Pb	Cr	Cu	Zn
Jos	Lettuce	<i>Lactuca sativa</i>	Salad					
	Root			0.66 ^b	17.03 ^c	1.25 ^a	8.34 ^a	ND
	Stem			0.09 ^a	15.13 ^a	2.25 ^c	6.75 ^a	ND
	Leaf			1.43 ^c	17.21 ^c	1.92 ^{ab}	14.61 ^b	ND
	Cabbage	<i>Brassica oleracea</i>	Kabeji					
	Root			2.87 ^b	22.52 ^c	1.83 ^a	11.45 ^b	0.27
	Stem			0.55 ^a	15.38 ^b	3.17 ^a	3.07 ^a	ND
	Leaf			5.03 ^c	5.50 ^a	2.67 ^a	0.38 ^a	ND
	Cucumber	<i>Cucumis sativus</i>	Kwawamba					
	Root			ND	ND	ND	ND	ND
	Stem			1.43 ^a	15.75 ^{ab}	3.67 ^c	214.48 ^c	ND
	Leaf			1.92 ^a	16.13 ^c	2.83 ^b	16.52 ^a	ND
	Fruit			1.38 ^a	12.78 ^a	1.83 ^a	26.08 ^b	ND
	Maize	<i>Zea mays</i>	Masara					
	Root			3.53 ^a	12.42 ^a	18.92 ^b	111.80 ^b	ND
	Stem			5.53 ^b	15.68 ^a	9.00 ^a	1.78 ^a	ND
	Leaf			6.14 ^b	15.37 ^a	9.08 ^a	30.78 ^a	ND
	Fruit			5.53 ^b	15.68 ^a	8.08 ^a	105.80 ^b	ND
	Rice	<i>Oryza sativa</i>	Shinkafa					
	Root			3.68 ^a	13.55 ^a	12.42 ^a	92.55 ^a	1.00 ^a
	Stem			3.92 ^a	18.70 ^{ab}	6.17 ^a	37.13 ^a	0.97 ^a
	leaf			ND	ND	ND	ND	ND
	Fruit			3.68 ^a	23.34 ^c	19.50 ^a	1150.50 ^b	0.93 ^a
Safe limits ^a				0.2	0.3	2.3	40	60

^a Source: FAO/WHO (2001). Mean followed with same letter across the column are not significantly different ($p > 0.05$).

The bioaccumulation factor (BAF) of heavy metals was $BAF > 1$ for Cd, Pb, and Zn and the BAF was in the decreasing order of $Cu > Zn > Pb > Cd > Cr$ (Figure 1). Pollution indices (PI) were >1 for Cd and Pb in all crops and in most crops for Cr (Figure 2). The estimated daily intake of metal (EDI) for adults exceeded the USEPA, ref. [10] oral reference dose daily limit in all the crops for Cd and Pb (Table 3). The EDI for Cr, Zn, and Cu were below the USEPA [10] except in *Solanum lycopersicum*, *Brassica oleracea*, and *Oryza sativa* for Cu. (Table 3). EDI values were in decreasing order of $Cu > Pb > Cr > Cd > Zn$. Hazard quotient (HQ) values were not detected for Zn and >1 for *Oryza sativa*. (Table 4). The HI values for all crops were >1 (4.666) in *Oryza sativa*, (1.475) *Zea mays*, (1.132), and *capsicum annum*.




Figure 1. Bioaccumulation factor (BAF) of heavy metals in the edible parts of crops grown in Plateau State (2018).




Figure 2. Pollution indices of heavy metals in the edible parts of crops grown in Plateau State (2018).

Table 3. Estimated daily intake of metals (EDI) (mg/kg/bw/day) through consumption of crops grown in Plateau State (2018).

Name of Sample	Botanical Name	Hausa Name	Estimated Daily Intake				
			Cd	Pb	Cr	Cu	Zn
Tomato	<i>Solanum lycopersicum</i>	Tomatur	0.007	0.053	0.016	0.051	0.002
Pepper	<i>Capsicum annum</i>	Attarugu	0.005	0.647	0.007	0.037	0.001
Onion	<i>Allium cepa</i>	Albasa	0.004	0.525	0.003	0.010	0.001
Carrot	<i>Daucus carota</i>	Karas	0.007	0.115	0.004	0.002	0.000
Spinach	<i>Spinacia oleracea</i>	Alayyaho	0.005	0.491	0.006	0.022	0.000
Lettuce	<i>Lactuca sativa</i>	Salad	0.002	0.524	0.003	0.021	0.000
Cabbage	<i>Brassica oleracea</i>	Kabeji	0.007	0.168	0.004	0.001	0.000
Cucumber	<i>Cucumis sativus</i>	Kokwamba	0.002	0.389	0.003	0.037	0.000
Maize	<i>Zea mays</i>	Masara	0.008	0.478	0.012	0.152	0.000
Rice	<i>Oryza sativa</i>	Shinkafa	0.005	0.711	0.028	1.649	0.001
RfD ^a			0.001	0.004	1.5	0.04	0.30

^a Source: USEPA, (2006).

Table 4. Hazard quotient and hazard index for adult population through the consumption of crops grown in Plateau State (2018).

Name of Sample	Botanical Name	Hausa Name	Hazard Quotient (HQ)					Hazard Index (HI)
			Cd	Pb	Cr	Cu	Zn	
Tomato	<i>Solanum lycopersicum</i>	Tomatur	0.619	0.053	0.001	0.107	ND	0.780
Pepper	<i>Capsicum annum</i>	Attarugu	0.406	0.647	ND	0.078	ND	1.132
Onion	<i>Allium cepa</i>	Albasa	0.331	0.525	ND	0.020	ND	0.877
Carrot	<i>Daucus carota</i>	Karas	0.564	0.115	ND	0.005	ND	0.684
Spinach	<i>Spinacia oleracea</i>	Alayyaho	0.391	0.491	ND	0.048	ND	0.930
Lettuce	<i>Lactuca sativa</i>	Salad	0.174	0.524	ND	0.044	ND	0.742
Cabbage	<i>Brassica oleracea</i>	Kabeji	0.613	0.168	ND	0.001	ND	0.782
Cucumber	<i>Cucumis sativus</i>	Kokwamba	0.169	0.389	ND	0.079	ND	0.637
Maize	<i>Zea mays</i>	Masara	0.674	0.478	0.001	0.322	ND	1.475
Rice	<i>Oryza sativa</i>	Shinkafa	0.449	0.711	0.002	3.504	ND	4.666

4. Discussion and Conclusions

The contamination of food crops by heavy metals from pesticides sources is a major concern for food quality and safety. The concentrations of Cd, Pb, and Cr in all the studied crops fumigated with pesticides as the only source of contamination exceeded the WHO [11] permissible limits, while the concentration of heavy metals in the corresponding soils of all the studied crops were below the UNEP [12] limits for agricultural soils. Most of the studied crops showed BAF > 1 for Cd, Pb, and Zn and the BAF was in decreasing order of Cu > Zn > Pb > Cd > Cr. Pollution indices indicated that most of the studied crops were contaminated for Pb, Cd, and Cr. The estimated daily intake of metals showed that all the studied crops exceeded the daily oral reference dose limit and could cause risk to humans. The hazard quotient showed all the studied crops were safe for human consumption except *Oryza sativa* for Cu which may cause risk to humans. However, people may be experiencing a severe adverse health risk (HI) from all the studied metals for the consumption of *Oryza sativa*, *Zea mays*, and *Capsicum annum*. Similar reports relating to this work include but are not limited to [13–18]. Thus, there is need for regular screening of heavy metals in pesticides. The predominant use of metal-based pesticides with high Cd, Pb, and Zn in the study areas could be responsible for the BAF > 1 values observed [3].

References

- Barau, B.W.; Abdulhamed, A.; Ezra, A.G.; Muhammed, M.; Bawa, U.; Yuguda, A.U.; Kyari, E.M. Heavy metal contamination of some vegetables from pesticides and potential Health risk in Bauchi, Northern Nigeria. *Int. J. Sci. Technol.* **2018**, *7*, 1–11. [\[CrossRef\]](#)
- Desalu, O.; Busari, O.; Adeoti, A. Respiratory Symptoms among Crop Farmers Exposed to Agricultural Pesticide in Three Rural Communities in South Western Nigeria: A Preliminary Study. *Ann. Med. Health Sci. Res.* **2014**, *4*, 662–666. [\[CrossRef\]](#) [\[PubMed\]](#)
- Yuguda, A.U.; Abubakar, Z.A.; Jibo, A.U.; AbdulHameed, A.; Nayaya, A.J. Assessment of Toxicity of Some Agricultural pesticides on Earthworm (*Lumbriscus Terrestris*). *Am. Eurasian J. Sustain. Agric.* **2015**, *9*, 49–59.
- Defarge, N.; De Vendômois, J.S.; Séralini, G.-E. Toxicity of formulants and heavy metals in glyphosate-based herbicides and other pesticides. *Toxicol. Rep.* **2018**, *5*, 156–163. [\[CrossRef\]](#) [\[PubMed\]](#)
- Zhong, T.; Xue, D.; Zhao, L.; Zhang, X. Concentration of heavy metals in vegetables and potential health risk assessment in China. *Environ. Geochem. Health* **2018**, *40*, 313–322. [\[CrossRef\]](#) [\[PubMed\]](#)
- Rattan, R.; Datta, S.P.; Chhonkar, P.K.; Suribabau, K.; Singh, A.K. Longterm impact of irrigation with sewage effluents on heavy metal content in soils, crops and ground Water (a case study). *Agric. Ecosyst. Environ.* **2005**, *109*, 310–322. [\[CrossRef\]](#)
- Hart, A.D.; Azubuike, C.U.; Barimalaa, I.S.; Achinewhu, S.C. Vegetable consumption Pattern of households in selected areas of the old Rivers state in Nigeria. *Afr. J. Food Agric. Nutr. Dev.* **2015**, *5*.
- Chukwuma, C. Evaluating baseline data for lead (Pb) and cadmium (Cd) in rice, yam, cassava and guinea grass from cultivated soils in Nigeria. *Toxicol. Environ. Chem.* **1994**, *45*, 45–56. [\[CrossRef\]](#)
- Mahmood, A.; Malik, R.N. Human health risk assessment of heavy metals through consumption of contaminated vegetables collected from different irrigation sources in Lahore, Pakistan. *Arab. J. Chem.* **2014**, *7*, 91–99. [\[CrossRef\]](#)

10. US-EPA. *Reference Dose (RFD). Description and Use in Health Risk Assessments Background Document 1A, Integrated Risk Information System (IRIS) 2013*; United States Environmental Protection Agency: Washington, DC, USA, 2013. Available online: <http://www.epa.gov/iris/rfd.htm> (accessed on 9 June 2021).
11. FAO/WHO Codex Alimentarius International Food Standards. General Standards for Contaminants and Toxins in Food and Feed. Available online: <http://www.fao.org/fao-whocodexalimentarius/en> (accessed on 9 June 2021).
12. UNEP. Environmental risk challenges of anthropogenic metals flows and cycles. In *A Report of the Working Group on the Global Metal Flow to the International Resource Panel*; Vandervoet, E., Salminen, R., Eckelman, M., Mudd, G., Norgate, T., Hirschier, R., Eds.; UNEP: Athens, Greece, 2013; p. 231.
13. Liang, G.; Gong, W.; Li, B.; Zuo, J.; Pan, L.; Liu, X. Analysis of Heavy Metals in Foodstuffs and an Assessment of the Health Risks to the General Public via Consumption in Beijing, China. *Int. J. Environ. Res. Public Health* **2019**, *16*, 909. [[CrossRef](#)] [[PubMed](#)]
14. Njuguna, S.M.; Makokha, V.A.; Yan, X.; Gituru, R.W.; Wang, Q.; Wang, J. Health risk assessment by consumption of vegetables irrigated with reclaimed waste water: A case study in Thika (Kenya). *J. Environ. Manag.* **2019**, *231*, 576–581. [[CrossRef](#)] [[PubMed](#)]
15. Peters, D.E.; Eebu, C.; Nkpaa, K.W. Potential Human Health Risk Assessment of Heavy Metals via Consumption of Root Tubers from Ogoniland, Rivers State, Nigeria. *Biol. Trace Elem. Res.* **2018**, *186*, 568–578. [[CrossRef](#)] [[PubMed](#)]
16. Eliku, T.; Leta, S. Heavy metals bioconcentration from soil to vegetables and appraisal of health risk in Koka and Wnji, Ethiopia. *Environ. Sci. Pollut. Res.* **2017**, *24*, 11807–11815. [[CrossRef](#)] [[PubMed](#)]
17. Proshad, R.; Kormoker, T.; Islam, S.; Chandra, K. Potential health risk of heavy metals via consumption of rice and vegetables grown in the industrial areas of Bangladesh. *Hum. Ecol. Risk Assess. Int. J.* **2019**, *26*, 921–943. [[CrossRef](#)]
18. Khan, S.; Cao, Q.; Zheng, Y.; Huang, Y.; Zhu, Y.-G. Health risks of heavy metals in contaminated soils and food crops irrigated with wastewater in Beijing, China. *Environ. Pollut.* **2008**, *152*, 686–692. [[CrossRef](#)] [[PubMed](#)]