



Abstract

# Recovery of Polyphenols and Compounds with Antioxidant Activity from Spirulina (*Arthrospira platensis*) through the Use of Different Organic Solvents Assisted by Pulsed Electric Fields <sup>†</sup>

Francisco J. Martí-Quijal \* , Francisco J. Barba and Francesc Ramon-Mascarell

Nutrition and Food Science Area, Preventive Medicine and Public Health, Food Science, Toxicology and Forensic Medicine Department, Faculty of Pharmacy, Universitat de València, Avda. Vicent Andrés Estellés, Burjassot, 46100 València, Spain; francisco.barba@uv.es (F.J.B.); ramas@alumni.uv.es (F.R.-M.)

\* Correspondence: francisco.j.marti@uv.es

<sup>†</sup> Presented at the 2nd International Electronic Conference on Foods—“Future Foods and Food Technologies for a Sustainable World”, 15–30 October 2021; Available online: <https://foods2021.sciforum.net/>.

**Abstract:** Currently, there is a growing nutritional interest in microalgae because they are a good source of nutrients and bioactive compounds. These bioactive compounds include polyphenols, which are known to be beneficial to health due to their antioxidant capacity. However, their extraction by conventional methods requires time and uses organic solvents, which are harmful to the environment. To reduce both time and environmental impact, the use of pulsed electric fields technology as a pre-treatment prior to extraction by stirring has been studied. For this purpose, a treatment of 100 kJ/kg at 3 kV/cm was applied to a suspension of 2% spirulina in water. Afterwards, an extraction was carried out using ethanol (EtOH) 50% or dimethyl sulfoxide (DMSO) 50% in water (*v/v*) for 3 h. In order to evaluate both the polyphenol extraction and the antioxidant capacity, a kinetic curve for each was established at different extraction times with the different solvents. It was observed that pre-treatment with PEF had a significant ( $p < 0.05$ ) positive effect on the extraction at all times with respect to the conventional treatment. The greatest differences were observed in the first 5–15 min of the extraction. The highest antioxidant capacity measured by ORAC and TEAC was obtained using EtOH as solvent. When DMSO was used, PEF pre-treatment allowed a higher recovery of polyphenols after 5 min ( $12.53 \pm 0.31$  mg gallic acid equivalents (GAE)/g dry weight) than the control sample after 180 min ( $4.84 \pm 0.48$  mg/g). For both solvents, the highest value of total phenolic compounds (TPC) was obtained after 120 min of extraction. Finally, PEF increased the extraction of polyphenols by 408% using EtOH 50% as solvent. It can be concluded that PEF technology increases the extraction of polyphenols from microalgae, reducing the process time and the consumption of organic solvents.

**Keywords:** spirulina (*Arthrospira platensis*); polyphenols; antioxidant activity; pulsed electric fields



**Citation:** Martí-Quijal, F.J.; Barba, F.J.; Ramon-Mascarell, F. Recovery of Polyphenols and Compounds with Antioxidant Activity from Spirulina (*Arthrospira platensis*) through the Use of Different Organic Solvents Assisted by Pulsed Electric Fields. *Biol. Life Sci. Forum* **2021**, *6*, 17. <https://doi.org/10.3390/Foods2021-10963>

Academic Editor: Diego Moreno-Fernandez

Published: 13 October 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/Foods2021-10963/s1>, Poster: Recovery of Polyphenols and Compounds with Antioxidant Activity from Spirulina (*Arthrospira platensis*) through the Use of Different Organic Solvents Assisted by Pulsed Electric Fields.

**Author Contributions:** Conceptualization, F.J.B.; methodology, F.J.M.-Q. and F.R.-M.; formal analysis, F.J.M.-Q.; investigation, F.J.M.-Q., F.J.B. and F.R.-M.; resources, F.J.B.; writing—original draft preparation, F.J.M.-Q. and F.R.-M.; writing—review and editing, F.J.M.-Q. and F.J.B.; supervision, F.J.B.; project administration, F.J.B.; funding acquisition, F.J.B. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the EU Commission and BBI-JU Horizon H2020, through AQUABIOPRO-FIT project (Aquaculture and agriculture biomass side stream proteins and bioactives for feed, fitness and health promoting nutritional supplements) grant number 790956. This research was also funded by Generalitat Valenciana (IDIFEDER/2018/046—Procesos innovadores de extracción y conservación: pulsos eléctricos y fluidos supercríticos) through the European Union ERDF funds (European Regional Development Fund).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** F.J.M-Q. would like to acknowledge the pre-PhD scholarship program of University of Valencia “Atracció de Talent”.

**Conflicts of Interest:** The authors declare no conflict of interest.