



Proceeding Paper

# Reconstruction of Models with Variable Cosmological Parameter in $f(R, T)$ Theory <sup>†</sup>

Rishi Kumar Tiwari <sup>1,‡</sup>, Aroonkumar Beesham <sup>2,\*</sup>  and Bhupendra Kumar Shukla <sup>3,‡</sup>

<sup>1</sup> Department of Mathematics, Govt Model Science College, Rewa 486001, MP, India; rishitiwari59@rediffmail.com

<sup>2</sup> Department of Mathematical Sciences, University of Zululand, P Bag X1001, Kwa-Dlangezwa 3886, South Africa

<sup>3</sup> Department of Mathematics, Pandit S.N. Shukla University, Shahdol 484001, MP, India; bhupendrashukla121@gmail.com

\* Correspondence: abeesham@yahoo.com

† Presented at the 1st Electronic Conference on Universe, 22–28 February 2021; Available online: <https://ecu2021.sciforum.net/>.

‡ These authors contributed equally to this work.

§ Alternate address: Faculty of Natural Sciences, Mangosuthu University of Technology, Jacobs 4026, South Africa.

**Abstract:** The standard  $\Lambda$ CDM model is reasonably successful in describing the universe, and is the most widely acceptable model in cosmology. However, there are several theoretical issues, such as the initial singularity, the cosmological constant problem, the particle nature of dark matter, the existence of anomalies in the cosmic microwave background radiation and on small scales, the predictions and tests of the inflationary scenario and whether general relativity is valid on the largest possible scales. Hence, there is growing interest in looking at modified theories. In this presentation, a reconstruction is made of the Friedmann–Lemaître–Robertson–Walker models with a dynamic cosmological parameter in  $f(R, T)$  modified gravity. This theory has a number of pleasing features, such as the avoidance of the initial Big Bang singularity and a variable cosmological parameter. A dynamic cosmological parameter, which arises naturally in this theory, can solve the cosmological constant problem, and it is also a candidate for dark energy. In addition, a variable cosmological parameter fits observations better than the standard  $\Lambda$ CDM model. The model exhibits a transition from deceleration to acceleration. The time evolution of the physical parameters such as energy density, pressure and equation of state are analyzed.

**Keywords:** dark energy; variable cosmological parameter;  $f(R, T)$  gravity



**Citation:** Tiwari, R.K.; Beesham, A.; Shukla, B.K. Reconstruction of Models with Variable Cosmological Parameter in  $f(R, T)$  Theory. *Phys. Sci. Forum* **2021**, *2*, 59. <https://doi.org/10.3390/ECU2021-09372>

Academic Editor: Yifu Cai

Published: 1 March 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The most widely accepted theory to study the evolution of the universe is undoubtedly Einstein's general theory of relativity, which predicts that the universe was condensed into a very small, hot and dense state initially, and then it expanded. In recent times, observations of Type Ia supernovae indicate that the current rate of expansion of the universe is accelerating. After this, many observations have supported the idea of an accelerated expansion [1–4]. In order to explain this acceleration, a new form of energy has been postulated that has a repulsive effect called dark energy (DE) [5–7]. According to the Planck mission team's best estimate [6], the universe is made up of three forms of matter/energy, viz., 68.5% DE, 26.6% dark matter (DM) and only 4.9% baryonic matter.

The most favored explanation for DE is Einstein's cosmological constant which is obtained by adding a cosmological constant to the equations of Einstein's standard Friedmann–Lemaître–Robertson–Walker (FLRW) equations and leads to the  $\Lambda$ CDM model [6]. This has a solution that includes accelerated expansion. Despite the fine tuning and coincidence

problems in the  $\Lambda$ CDM model, it is widely accepted as the best solution to the DE problem [8]. While this DE is constant, another consideration is a changing field, in which the equation of state (EOS) varies, viz., quintessence, phantom DE and scalar field models. For excellent reviews of DE, we refer to the works in [8,9].

In another direction, the reason for the acceleration of the universe can be sought in modified theories of gravitation. One of these is  $f(R, T)$  gravity [10], where  $R$  is the Ricci scalar and  $T$  is the trace of the energy momentum tensor. This theory allows for an explanation of accelerated expansion without DE and the avoidance of the initial singularity. A very interesting feature of the theory is that, in some sense, it may be thought of as general relativity with a modified matter part, thus allowing for a dynamic cosmological parameter [11]. This allows for the possibility of solving the cosmological constant problem. In this paper, we reconstruct the Friedman–Lemaître–Robertson–Walker (FLRW) model in  $f(R, T)$  theory, focussing on a variable cosmological parameter. This correspondence has not been fully studied. We find a model that exhibits a transition from deceleration in the past, to current acceleration. The behavior of the physical parameters is analyzed, as are the energy conditions. The cosmological parameter varies, being large at early times and decreasing to a small value today.

## 2. Basic Equations

In  $f(R, T)$  gravity, the action is:

$$S = \int \left( \frac{1}{16\pi G} f(R, T) + S_m \right) \sqrt{-g} dx^4. \tag{1}$$

We choose [10]

$$f(R, T) = f_1(R) + f_2(T), \tag{2}$$

i.e., a sum of two independent functions of  $R$  and  $T$ , respectively. With this condition, Equation (1) can be written as

$$f_1'(R)R_{ij} - \frac{1}{2}(f_1(R) + f_2(T))g_{ij} + (g_{ij}\square - \nabla_i\nabla_j)f_1'(R) = 8\pi T_{ij} - f_2'(T)T_{ij} - f_2'(T)\Theta_{ij}, \tag{3}$$

where  $\square \equiv \nabla^i\nabla_j$  is the D'Alembertian operator. The tensor  $\Theta_{ij}$  is defined as

$$\Theta_{ij} = g^{lm} \frac{\delta T_{lm}}{\delta g^{ij}}. \tag{4}$$

A prime denotes a derivative with respect to its argument.

In particular [12], we assume the forms

$$f_1(R) = \lambda_1 R, f_2(T) = \lambda_2 T \tag{5}$$

where  $\lambda_1$  and  $\lambda_2$  are arbitrary coupling constants of  $f(R, T)$  gravity. We take the matter content in the universe to be a perfect fluid with energy momentum tensor:

$$T_{ij} = (\rho + p)u_i u_j - pg_{ij}, \tag{6}$$

where  $\rho$  and  $p$  are energy density and pressure of the fluid, respectively, and  $u^i$  is four velocity vector satisfying the condition  $u^i u_i = 1$ . Then, Equation (4) yields:

$$\Theta_{ij} = -2T_{ij} - pg_{ij}, \tag{7}$$

Using Equations (4)–(7), Equation (3) becomes:

$$R_{ij} - \frac{1}{2}Rg_{ij} = \left( \frac{8\pi + \lambda_2}{\lambda_1} \right) T_{ij} + \frac{\lambda_2}{\lambda_1} \left( p + \frac{1}{2}T \right) g_{ij}. \tag{8}$$

Note that this equation reduces to the general relativistic limit when we put  $\lambda_1 = 1$ ,  $\lambda_2 = 0$ .

Now, in the general theory of relativity, the Einstein field equations with cosmological constant are (in units  $G = c = 1$ ):

$$R_{ij} - \frac{1}{2}Rg_{ij} = 8\pi T_{ij} + \Lambda g_{ij}, \tag{9}$$

We notice a similarity between Equations (8) and (9). Hence, we may set:

$$\Lambda \equiv \Lambda(T) = \frac{\lambda_2}{\lambda_1} \left( p + \frac{1}{2}T \right), \tag{10}$$

which leads naturally to a varying cosmological parameter  $\Lambda$  as a function of  $T$ . Evaluating the trace  $T$  from Equation (6), we obtain:

$$\Lambda = \frac{\lambda_2}{2\lambda_1} (\rho - p). \tag{11}$$

We consider the flat FLRW metric:

$$ds^2 = dt^2 - \sum_{i=1}^3 a^2(t)(dx_i^2), \tag{12}$$

where  $a(t)$  is the scale factor. The Hubble and deceleration parameters are defined by, respectively:

$$H = \frac{\dot{a}}{a}, \quad q = -\frac{\ddot{a}a}{\dot{a}^2} \tag{13}$$

With the metric (12), the field Equation (8) become, with the aid of Equation (10):

$$3H^2 = \left[ \left( \frac{8\pi + \lambda_2}{\lambda_1} \right) + \frac{\lambda_2}{2\lambda_1} \right] \rho - \frac{\lambda_2}{2\lambda_1} p, \tag{14}$$

$$2\dot{H} + 3H^2 = - \left[ \left( \frac{8\pi + \lambda_2}{\lambda_1} \right) + \frac{\lambda_2}{2\lambda_1} \right] p + \frac{\lambda_2}{2\lambda_1} \rho, \tag{15}$$

We obtain the energy density and pressure from the above two equations as

$$\rho = \frac{\lambda_1}{(8\pi + 2\lambda_2)} \left[ 3 + \frac{\lambda_2}{(8\pi + \lambda_2)}(q + 1) \right] H^2, \tag{16}$$

$$p = \frac{\lambda_1}{(8\pi + 2\lambda_2)} \left[ -3 + \frac{(16\pi + 3\lambda_2)}{(8\pi + \lambda_2)}(q + 1) \right] H^2, \tag{17}$$

and, then, the cosmological parameter from Equation (10) as:

$$\Lambda = \left[ \frac{3\lambda_2}{(8\pi + 2\lambda_2)} - \frac{\lambda_2}{(8\pi + 2\lambda_2)}(q + 1) \right] H^2. \tag{18}$$

The equation of state parameter ( $\omega = p/\rho$ ) is given by

$$\omega = -1 + \frac{(16\pi + 4\lambda_2)(q + 1)}{(24\pi + 4\lambda_2) + \lambda_2 q}. \tag{19}$$

### 3. Solution to Field Equations

To solve the field equations, we assume the following condition for  $q$  [13]:

$$q = -1 + \beta H, \tag{20}$$

where  $\beta$  is a constant. From the definitions of  $H$  and  $q$ , Equation (13), we can write the solution as:

$$a = \exp\left[\frac{1}{\beta}\sqrt{2\beta t + k}\right], \quad H = \frac{1}{\sqrt{2\beta t + k}}, \quad q = -1 + \frac{\beta}{\sqrt{2\beta t + k}} \quad (21)$$

We note from Equation (21) that  $q$  is positive at early times, corresponding to deceleration, and negative at late times, corresponding to acceleration.

We now use observations to estimate the values of the constant  $\beta$  and the integration constant  $k$ . If we evaluate Equation (20) at the present time, we have

$$q_0 = -1 + \beta H_0, \quad (22)$$

Observations tell us that  $q_0 = -0.51$  [14] and that  $H_0 = 75.35 \text{ km} \cdot \text{s}^{-1}\text{Mpc}^{-1}$  [15]. Substituting these values into Equation (22), we get  $\beta = 0.0065$ . From Equation (21), we then get the value of  $k$  as  $k = 0.000175$ . Now, how do we select appropriate values for  $\lambda_1$  and  $\lambda_2$ ? Since the general relativistic limit is given by  $\lambda_1 \rightarrow 1, \lambda_2 \rightarrow 0$ , we choose  $\lambda_1 = 0.9$  and  $\lambda_2 = 0.00016$  to plot all subsequent figures. These values of  $\beta, k, \lambda_1$  and  $\lambda_2$  are used in plotting all figures.

From the observational point of view, it is more useful to express the parameters in term of redshift  $z$ . The average scale factor  $a$  in terms of redshift is given by

$$a(z) = \frac{a_0}{1+z}. \quad (23)$$

From Equation (21), we get

$$\frac{1}{\beta}\sqrt{2\beta t + k} = \ln(a), \quad (24)$$

and, from Equation (23), we get

$$\ln(a) = \ln(a_0) - \ln(1+z). \quad (25)$$

Hence, the Hubble and deceleration parameters in terms of redshift are as follows:

$$H(z) = \frac{1}{\beta[\ln(a_0) - \ln(1+z)]}, \quad (26)$$

$$q(z) = -1 + \frac{1}{[\ln(a_0) - \ln(1+z)]}. \quad (27)$$

We now plot the deceleration parameter  $q$  against the redshift  $z$  in Figure 1.

Using the  $t - z$  relationship, we get the energy density, pressure, cosmological parameter and equation of state parameters as follows:

$$\rho = \frac{\lambda_1}{(8\pi + 2\lambda_2)} \left[ 3 + \frac{\lambda_2}{(8\pi + \lambda_2)[\ln(a_0) - \ln(1+z)]} \right] \frac{1}{\beta^2[\ln(a_0) - \ln(1+z)]^2}, \quad (28)$$

$$p = \frac{\lambda_1}{(8\pi + 2\lambda_2)} \left[ -3 + \frac{(16\pi + 3\lambda_2)}{(8\pi + \lambda_2)[\ln(a_0) - \ln(1+z)]} \right] \frac{1}{\beta^2[\ln(a_0) - \ln(1+z)]^2}, \quad (29)$$

$$\Lambda = \left[ \frac{3\lambda_2}{(8\pi + 2\lambda_2)} - \frac{\lambda_2}{(8\pi + 2\lambda_2)[\ln(a_0) - \ln(1+z)]} \right] \frac{1}{\beta^2[\ln(a_0) - \ln(1+z)]^2}, \quad (30)$$

and

$$\omega = -1 + \frac{(16\pi + 4\lambda_2)}{3(8\pi + \lambda_2)[\ln(a_0) - \ln(1+z)] + \lambda_2}. \quad (31)$$




Figure 1. Deceleration parameter  $q$  against redshift  $z$ .

We can plot these quantities in Figures 2–5.




Figure 2. Energy density against redshift  $z$ .




Figure 3. Pressure against redshift  $z$ .




Figure 4. Cosmological parameter against redshift  $z$ .




Figure 5. EOS parameter against redshift  $z$ .

#### 4. Conclusions

In this paper, we analyze a model with a variable cosmological parameter in flat FLRW space-time in  $f(R, T)$  theory. We graph the various physical parameters against redshift. The vital characteristics of our model are the following. Our model presents decelerating in the past and current accelerating, as shown in Figure 1. The behavior of the parameters  $q, \rho, p, \omega$  and  $\Lambda$  are illustrated in Figures 2–5. We have a variable cosmological parameter, which is large initially and decreases with time. This could help in resolving the cosmological constant problem.

**Author Contributions:** Conceptualization, R.K.T., A.B. and B.K.S.; methodology, R.K.T.; software, A.B. and B.K.S.; validation, R.K.T., A.B. and B.K.S.; formal analysis, R.K.T., A.B. and B.K.S.; investigation, R.K.T., A.B. and B.K.S.; resources, R.K.T. and B.K.S.; data curation, B.K.S.; writing—original draft preparation, R.K.T. and B.K.S.; writing—review and editing, R.K.T., A.B. and B.K.S.; visualization, R.K.T., A.B. and B.K.S.; supervision, R.K.T. and A.B.; project administration, R.K.T.; funding acquisition, A.B. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was based on the research supported wholly/in part by the National Research Foundation of South Africa (Grant Numbers: 118511).

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

## References

1. Perlmutter, S.; Aldering, G.; Goldhaber, G.; Knop, R.A.; Nugent, P.; Castro, P.G.; Deustua, S.; Fabbro, S.; Goobar, A.; Groom, D.E.; et al. Measurements of  $\Omega$  and  $\Lambda$  from 42 High-Redshift Supernovae. *Astrophys. J.* **1999**, *517*, 565–586. [[CrossRef](#)]
2. Riess, A.G.; Filippenko, A.V.; Challis, P.; Clocchiatti, A.; Diercks, A.; Garnavich, P.M.; Gilliland, R.L.; Hogan, C.J.; Jha, S.; Kirshner, R.P.; et al. Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. *Astron. J.* **1998**, *116*, 1009–1038. [[CrossRef](#)]
3. Percival, W.J.; Baugh, C.M.; Bland-Hawthorn, J.; Bridges, T.; Cannon, R.; Cole, S.; Colless, M.; Collins, C.; Couch, W.; Dalton, G.; et al. The 2dF Galaxy Redshift Survey: The power spectrum and the matter content of the universe. *Mon. Not. R. Astron. Soc.* **2001**, *327*, 1297–1306. [[CrossRef](#)]
4. Suwa, M.; Kobayashi, K.; Oshima, H. The Interacting Generalized Ricci Dark Energy Model in Non-Flat Universe. *J. Cosmol. Astropart. Phys.* **2010**, *1002*, 008. [[CrossRef](#)]
5. Spergel, D.N.; Verde, L.; Peiris, H.V.; Komatsu, E.; Nolta, M.R.; Bennett, C.L.; Halpern, M.; Hinshaw, G.; Jarosik, N.; Kogut, A.; et al. First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters. *Astrophys. J. Suppl.* **2003**, *148*, 175–194. [[CrossRef](#)]
6. Aghanim, N.; Akrami, Y.; Ashdown, M.; Aumont, J.; Baccigalupi, C.; Ballardini, M.; Banday, A.J.; Barreiro, R.B.; Bartolo, N.; Basak, S.; et al. Planck 2018 results VI. Cosmological parameters. *Astron. Astrophys.* **2020**, *641*, A6.
7. Eisenstein, D.J.; Zehavi, I.; Hogg, D.W.; Scoccamarro, R.; Blanton, M.R.; Nichol, R.C.; Scranton, R.; Seo, H.J.; Tegmark, M.; Zheng, Z.; et al. Detection of the Baryon Acoustic Peak in the Large-Scale Correlation Function of SDSS Luminous Red Galaxies. *Astrophys. J.* **2005**, *633*, 560–574. [[CrossRef](#)]
8. Bull, P.; Akrami, Y.; Adamek, J.; Baker, T.; Bellini, E.; Jimenez, J.B.; Bentivegna, E.; Camera, S.; Clesse, S.; Davis, J.H.; et al. Beyond  $\Lambda$ CDM: Problems, solutions, and the road ahead. *Phys. Dark Univ.* **2016**, *12*, 56–99. [[CrossRef](#)]
9. Durrer, R.; Maartens, R. Dark Energy and Dark Gravity. *Gen. Relativ. Grav.* **2008**, *40*, 301–328. [[CrossRef](#)]
10. Harko, T.; Lobo, F.S.N.; Nokiri, S.; Odintsov, S.D.  $f(R, T)$  gravity. *Phys. Rev. D* **2011**, *84*, 024020. [[CrossRef](#)]
11. Tiwari, R.K.; Beesham, A. Anisotropic model with decaying cosmological term. *Astrophys. Space Sci.* **2018**, *363*, 234. [[CrossRef](#)]
12. Pasqua, A.; Chattopadhyay, S.; Khomenko, I. A reconstruction of modified holographic Ricci dark energy in  $f(R, T)$  gravity. *Can. J. Phys.* **2013**, *91*, 632–638. [[CrossRef](#)]
13. Tiwari, R.K.; Singh, R.; Shukla, B.K. A Cosmological Model with Variable Deceleration Parameter. *Afr. Rev. Phys.* **2015**, *10*, 48. [[CrossRef](#)]
14. Jusus, J.F. Gaussian process estimation of transition redshift. *J. Cosmol. Astropart. Phys.* **2020**, *4*, 053. [[CrossRef](#)]
15. Camarena, D.; Marra, V. Local determination of the Hubble constant and the deceleration parameter. *Phys. Rev. Res.* **2020**, *2*, 013028. [[CrossRef](#)]